

=> d his full

(FILE 'HOME' ENTERED AT 09:30:05 ON 11 OCT 2007)

FILE 'REGISTRY' ENTERED AT 09:30:15 ON 11 OCT 2007

L1 STRUCTURE UPLOADED
 L2 1 SEA SSS SAM L1
 D SCA
 L3 560 SEA SSS FUL L1
 SAVE TEMP L3 LAO058STR1L/A

FILE 'ZCAPLUS' ENTERED AT 09:35:51 ON 11 OCT 2007

L4 117 SEA ABB=ON PLU=ON L3
 L5 ANALYZE PLU=ON L4 1- RN : 5098 TERMS
 D
 D
 D 1-20

FILE 'REGISTRY' ENTERED AT 09:37:09 ON 11 OCT 2007

L6 1 SEA ABB=ON PLU=ON 152044-54-7
 L7 1 SEA ABB=ON PLU=ON 152044-53-6
 L8 1 SEA ABB=ON PLU=ON 189453-10-9
 L9 1 SEA ABB=ON PLU=ON 186692-73-9
 L10 1 SEA ABB=ON PLU=ON 187527-25-9
 L11 1 SEA ABB=ON PLU=ON 188730-08-7
 L12 1 SEA ABB=ON PLU=ON 20949-84-2
 L13 1 SEA ABB=ON PLU=ON 106921-60-2
 L14 1 SEA ABB=ON PLU=ON 193146-27-9
 L15 0 SEA ABB=ON PLU=ON (L6 OR L7 OR L8 OR L9 OR L10 OR L11 OR L12
 OR L13 OR L14) AND L4
 L16 0 SEA ABB=ON PLU=ON (L6 OR L7 OR L8 OR L9 OR L10 OR L11 OR L12
 OR L13 OR L14) AND L3
 L17 1 SEA ABB=ON PLU=ON 187283-46-1
 L18 1 SEA ABB=ON PLU=ON 188899-14-1
 L*** DEL 0 S S 184246-38-6
 L19 1 SEA ABB=ON PLU=ON 184246-38-6
 L20 1 SEA ABB=ON PLU=ON 189453-35-8
 L21 1 SEA ABB=ON PLU=ON 219989-84-1
 L22 1 SEA ABB=ON PLU=ON 63928-37-0
 L23 1 SEA ABB=ON PLU=ON 52079-23-9
 L24 1 SEA ABB=ON PLU=ON 70113-32-5
 L25 1 SEA ABB=ON PLU=ON 185148-95-2
 L26 1 SEA ABB=ON PLU=ON 186692-84-2
 L27 0 SEA ABB=ON PLU=ON L3 AND (L6 OR L7 OR L8 OR L9 OR L10 OR L11
 OR L12 OR L13 OR L14 OR L15 OR L16 OR L17 OR L18 OR L19 OR L20
 OR L21 OR L22 OR L23 OR L24 OR L25 OR L26)
 L28 191 SEA ABB=ON PLU=ON L3 AND CASREACT/LC

FILE 'ZCAPLUS' ENTERED AT 09:43:45 ON 11 OCT 2007

L29 115 SEA ABB=ON PLU=ON L3/P

FILE 'CASREACT' ENTERED AT 09:53:42 ON 11 OCT 2007

L30 69 SEA ABB=ON PLU=ON L3
 L31 STRUCTURE UPLOADED
 L32 0 SEA SUB=L30 SSS SAM L31 (0 REACTIONS)
 L33 23 SEA SUB=L30 SSS FUL L31 (468 REACTIONS)
 D STAT QUE L33

SN 10/563058 Page 168 of 172 STIC STN SEARCH RESULTS

FILE 'REGISTRY' ENTERED AT 11:21:02 ON 11 OCT 2007

L34 22933 SEA ABB=ON PLU=ON OC15/ESS
 L35 27330 SEA ABB=ON PLU=ON C16/ESS
 L36 726 SEA ABB=ON PLU=ON NC15/ESS
 L37 0 SEA ABB=ON PLU=ON NSC14/ESS
 L38 50989 SEA ABB=ON PLU=ON (L34 OR L35 OR L36 OR L37)
 L39 12165 SEA ABB=ON PLU=ON L38 AND CASREACT/LC

FILE 'CASREACT' ENTERED AT 11:22:32 ON 11 OCT 2007

L40 2534 SEA ABB=ON PLU=ON L39/PRO
 L41 59 SEA ABB=ON PLU=ON L30 (L) L40
 L42 65 SEA ABB=ON PLU=ON L3/RRT
 L43 59 SEA ABB=ON PLU=ON L42 (L) L40
 L44 19 SEA ABB=ON PLU=ON L43 AND L33

FILE 'CAPLUS' ENTERED AT 11:34:47 ON 11 OCT 2007

L45 59 SEA ABB=ON PLU=ON L43
 L46 53 SEA ABB=ON PLU=ON L45 AND PY<2005
 L47 49 SEA ABB=ON PLU=ON L45 AND PY<2004

FILE 'CASREACT' ENTERED AT 11:35:34 ON 11 OCT 2007

D L43

L48 59 SEA ABB=ON PLU=ON L43 AND 1/NS
 L49 50 SEA ABB=ON PLU=ON L43 AND 2/NS
 L50 44 SEA ABB=ON PLU=ON L43 AND 3/NS
 L51 42 SEA ABB=ON PLU=ON L43 AND 4/NS
 L52 9 SEA ABB=ON PLU=ON L48 NOT L49

D SCA

FILE 'CAPLUS' ENTERED AT 11:45:06 ON 11 OCT 2007

L53 45 SEA ABB=ON PLU=ON L45 AND J/DT
 L54 14 SEA ABB=ON PLU=ON L45 AND P/DT
 L55 12 SEA ABB=ON PLU=ON L54 AND PD<20040619
 L56 39 SEA ABB=ON PLU=ON L53 AND ED<20040619
 L57 6 SEA ABB=ON PLU=ON L53 NOT L56
 L58 2 SEA ABB=ON PLU=ON L54 NOT L55
 L59 8 SEA ABB=ON PLU=ON (L57 OR L58)
 SEL AN

FILE 'CASREACT' ENTERED AT 11:47:33 ON 11 OCT 2007

L60 8 SEA ABB=ON PLU=ON ("142:134344"/AN OR "143:211773"/AN OR
 "143:422202"/AN OR "144:170808"/AN OR "145:271524"/AN OR
 "145:397261"/AN OR "146:229070"/AN OR "146:251631"/AN OR
 "2004:985335"/AN OR "2005:1154536"/AN OR "2005:1305128"/AN OR
 "2005:614221"/AN OR "2006:1337456"/AN OR "2006:641138"/AN OR
 "2006:66747"/AN OR "2006:805502"/AN)
 L61 51 SEA ABB=ON PLU=ON L48 NOT L60
 L62 42 SEA ABB=ON PLU=ON L49 NOT L60
 L63 36 SEA ABB=ON PLU=ON L50 NOT L60
 L64 35 SEA ABB=ON PLU=ON L51 NOT L60

FILE 'CAPLUS' ENTERED AT 11:49:27 ON 11 OCT 2007

E US2006-563058 /APPS

L65 1 SEA ABB=ON PLU=ON US2006-563058 /AP
 D SCA
 SEL RN

FILE 'REGISTRY' ENTERED AT 11:50:13 ON 11 OCT 2007

L66 55 SEA ABB=ON PLU=ON (130486-85-0/BI OR 152044-53-6/BI OR
 152044-54-7/BI OR 185148-95-2/BI OR 220367-73-7/BI OR 220774-16

-3/BI OR 220774-19-6/BI OR 220774-20-9/BI OR 220774-21-0/BI OR
 220774-22-1/BI OR 220774-23-2/BI OR 220774-57-2/BI OR 220774-58
 -3/BI OR 220774-59-4/BI OR 220774-60-7/BI OR 220774-61-8/BI OR
 220774-62-9/BI OR 220775-18-8/BI OR 220775-76-8/BI OR 289477-70
 -9/BI OR 289477-71-0/BI OR 289477-72-1/BI OR 289477-73-2/BI OR
 289477-74-3/BI OR 303154-55-4/BI OR 303154-56-5/BI OR 303154-57
 -6/BI OR 303154-58-7/BI OR 303154-59-8/BI OR 303154-60-1/BI OR
 305840-13-5/BI OR 823203-01-6/BI OR 823203-02-7/BI OR 823203-03
 -8/BI OR 823203-04-9/BI OR 823203-05-0/BI OR 823203-06-1/BI OR
 823203-07-2/BI OR 823203-08-3/BI OR 823203-09-4/BI OR 823203-10
 -7/BI OR 823203-11-8/BI OR 823203-12-9/BI OR 823203-13-0/BI OR
 823203-14-1/BI OR 823203-15-2/BI OR 823203-16-3/BI OR 823203-17
 -4/BI OR 823203-18-5/BI OR 823203-19-6/BI OR 823203-20-9/BI OR
 823203-23-2/BI OR 823203-24-3/BI OR 823203-25-4/BI OR 823203-27
 -6/BI)

L67 2 SEA ABB=ON PLU=ON L66 AND L39
 D SCA

FILE 'CAPLUS' ENTERED AT 11:50:53 ON 11 OCT 2007

L68 1 SEA ABB=ON PLU=ON L67 AND L65
 D SCA

FILE 'REGISTRY' ENTERED AT 11:54:40 ON 11 OCT 2007

E EPOTHILONE C/CN

L69 1 SEA ABB=ON PLU=ON EPOTHILONE C/CN
 D SCA

L70 1 SEA ABB=ON PLU=ON EPOTHILONE D/CN
 D SCA
 D RN L67 1-2

FILE 'CASREACT' ENTERED AT 11:56:56 ON 11 OCT 2007

L71 21 SEA ABB=ON PLU=ON 152044-54-7/PRO
 L72 14 SEA ABB=ON PLU=ON 152044-53-6/PRO
 L73 7 SEA ABB=ON PLU=ON L42 (L) L71
 L74 7 SEA ABB=ON PLU=ON L42 (L) L72
 L75 14 SEA ABB=ON PLU=ON (L73 OR L74)
 L76 13 SEA ABB=ON PLU=ON L75 NOT L60

FILE 'CAPLUS' ENTERED AT 11:58:45 ON 11 OCT 2007

L77 11 SEA ABB=ON PLU=ON L45 AND PY<2000
 SEL AN

FILE 'CASREACT' ENTERED AT 12:00:01 ON 11 OCT 2007

L78 11 SEA ABB=ON PLU=ON ("126:251010"/AN OR "127:108793"/AN OR
 "127:293040"/AN OR "128:101936"/AN OR "129:189151"/AN OR
 "131:199535"/AN OR "131:286299"/AN OR "131:31819"/AN OR
 "131:31829"/AN OR "131:351125"/AN OR "132:49832"/AN OR
 "1997:206419"/AN OR "1997:430309"/AN OR "1997:665094"/AN OR
 "1997:787450"/AN OR "1998:378435"/AN OR "1999:176999"/AN OR
 "1999:372044"/AN OR "1999:383492"/AN OR "1999:444724"/AN OR
 "1999:606636"/AN OR "1999:819379"/AN)
 L79 11 SEA ABB=ON PLU=ON L78 AND L43
 L80 16 SEA ABB=ON PLU=ON L79 OR L52
 L81 1 SEA ABB=ON PLU=ON L80 AND L73
 L82 1 SEA ABB=ON PLU=ON L80 AND L74
 L83 27 SEA ABB=ON PLU=ON L52 OR L79 OR (L81 OR L82) OR L76
 L84 15 SEA ABB=ON PLU=ON L61 NOT L63
 D FHIT 7
 L85 59 SEA ABB=ON PLU=ON L43 (L) 1/NS
 L86 45 SEA ABB=ON PLU=ON L43 (L) 2/NS

L87 37 SEA ABB=ON PLU=ON L43 (L) 3/NS
L88 31 SEA ABB=ON PLU=ON L43 (L) 4/NS
L89 21 SEA ABB=ON PLU=ON L43 (L) 5/NS
L90 28 SEA ABB=ON PLU=ON L85 NOT L88

FILE 'CAPLUS' ENTERED AT 12:09:08 ON 11 OCT 2007

L91 80 SEA ABB=ON PLU=ON KLAR U?/AU
L92 116 SEA ABB=ON PLU=ON BUCHMANN B?/AU
L93 60 SEA ABB=ON PLU=ON SCHWEDE W?/AU
L94 186 SEA ABB=ON PLU=ON SKUBALLA W?/AU
L95 32 SEA ABB=ON PLU=ON L91 AND (L92 OR L93 OR L94)
L96 68 SEA ABB=ON PLU=ON L92 AND (L93 OR L94)
L97 24 SEA ABB=ON PLU=ON L93 AND L94
L98 25 SEA ABB=ON PLU=ON L95 AND (L96 OR L97)
L99 24 SEA ABB=ON PLU=ON L96 AND L97
L100 24 SEA ABB=ON PLU=ON L98 AND L99
L101 1 SEA ABB=ON PLU=ON L100 AND L43

FILE 'REGISTRY' ENTERED AT 12:11:27 ON 11 OCT 2007

FILE 'CAPLUS' ENTERED AT 12:11:29 ON 11 OCT 2007

D STAT QUE L100
D IBIB ABS L100 1-24
D IBIB ABS L100 8-24
L102 24 SEA ABB=ON PLU=ON L91 AND L92 AND L93 AND L94
D COST FULL
D IBIB ABS L102 TOT
D IBIB L102 10
D IBIB L102 9
D ABS L102 8
D ABS L102 8
D IBIB ABS L102 9-24

FILE 'REGISTRY' ENTERED AT 12:16:30 ON 11 OCT 2007

FILE 'CASREACT' ENTERED AT 12:16:34 ON 11 OCT 2007

D STAT QUE L33
D IBIB ABS FHIT L33 1-23

FILE 'CASREACT' ENTERED AT 12:18:14 ON 11 OCT 2007

D STAT QUE L90
D IBIB ABS FHIT L90 1-28
L103 3 SEA ABB=ON PLU=ON L77 NOT L90
L104 3 SEA ABB=ON PLU=ON L78 NOT L90

FILE 'CAPLUS' ENTERED AT 12:40:57 ON 11 OCT 2007

L105 21 SEA ABB=ON PLU=ON L45 AND PY<2001
SEL AN

FILE 'CASREACT' ENTERED AT 12:41:26 ON 11 OCT 2007

L106 21 SEA ABB=ON PLU=ON ("126:251010"/AN OR "127:108793"/AN OR
"127:293040"/AN OR "128:101936"/AN OR "129:189151"/AN OR
"131:199535"/AN OR "131:286299"/AN OR "131:31819"/AN OR
"131:31829"/AN OR "131:351125"/AN OR "132:251011"/AN OR
"132:49832"/AN OR "133:266631"/AN OR "133:266634"/AN OR
"133:321737"/AN OR "133:362657"/AN OR "134:178371"/AN OR
"134:29228"/AN OR "134:4795"/AN OR "134:56502"/AN OR "135:37156
6"/AN OR "1997:206419"/AN OR "1997:430309"/AN OR "1997:665094"/
AN OR "1997:787450"/AN OR "1998:378435"/AN OR "1999:176999"/AN
OR "1999:372044"/AN OR "1999:383492"/AN OR "1999:444724"/AN OR

"1999:606636"/AN OR "1999:819379"/AN OR "2000:514132"/AN OR
"2000:52387"/AN OR "2000:597944"/AN OR "2000:624043"/AN OR
"2000:701228"/AN OR "2000:719579"/AN OR "2000:733774"/AN OR
"2000:842116"/AN OR "2000:853645"/AN OR "2001:843887"/AN)
L107 7 SEA ABB=ON PLU=ON L106 NOT L90
L108 21 SEA ABB=ON PLU=ON L106 AND L43
L109 7 SEA ABB=ON PLU=ON L108 NOT L90
D STAT QUE L109
D IBIB ABS FHIT L109 1-7

FILE HOME

FILE REGISTRY

Property values tagged with IC are from the ZIC/VINITI data file
provided by InfoChem.

STRUCTURE FILE UPDATES: 10 OCT 2007 HIGHEST RN 950149-06-1

DICTIONARY FILE UPDATES: 10 OCT 2007 HIGHEST RN 950149-06-1

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH June 29, 2007

Please note that search-term pricing does apply when
conducting SmartSELECT searches.

REGISTRY includes numerically searchable data for experimental and
predicted properties as well as tags indicating availability of
experimental property data in the original document. For information
on property searching in REGISTRY, refer to:

<http://www.cas.org/support/stngen/stndoc/properties.html>

FILE ZCAPLUS

Copyright of the articles to which records in this database refer is
held by the publishers listed in the PUBLISHER (PB) field (available
for records published or updated in Chemical Abstracts after December
26, 1996), unless otherwise indicated in the original publications.
The CA Lexicon is the copyrighted intellectual property of the
American Chemical Society and is provided to assist you in searching
databases on STN. Any dissemination, distribution, copying, or storing
of this information, without the prior written consent of CAS is
strictly prohibited.

FILE COVERS 1907 - 11 Oct 2007 VOL 147 ISS 16

FILE LAST UPDATED: 10 Oct 2007 (20071010/ED)

New CAS Information Use Policies, enter HELP USAGETERMS for details.

This file contains CAS Registry Numbers for easy and accurate
substance identification.

FILE CASREACT

Copyright of the articles to which records in this database refer is
held by the publishers listed in the PUBLISHER (PB) field (available
for records published or updated in Chemical Abstracts after December
26, 1996), unless otherwise indicated in the original publications.

FILE CONTENT:1840 - 6 Oct 2007 VOL 147 ISS 16

New CAS Information Use Policies, enter HELP USAGETERMS for details.

*
* CASREACT now has more than 13.8 million reactions *
* *

Some CASREACT records are derived from the ZIC/VINITI database (1974-1999) provided by InfoChem, INPI data prior to 1986, and Biotransformations database compiled under the direction of Professor Dr. Klaus Kieslich.

This file contains CAS Registry Numbers for easy and accurate substance identification.

FILE CAPLUS

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 11 Oct 2007 VOL 147 ISS 16
FILE LAST UPDATED: 10 Oct 2007 (20071010/ED)

Effective October 17, 2005, revised CAS Information Use Policies apply. They are available for your review at:

<http://www.cas.org/infopolicy.html>

=>

=> d his full

(FILE 'HOME' ENTERED AT 09:30:05 ON 11 OCT 2007)
 FILE 'REGISTRY' ENTERED AT 09:30:15 ON 11 OCT 2007
 STRUCTURE UPLOADED
 1 SEA SSS SAM L1
 D SCA
 560 SEA SSS FUL L1
 SAVE TEMP L3 LA0058STR1L/A
 FILE 'ZCAPLUS' ENTERED AT 09:35:51 ON 11 OCT 2007
 117 SEA ABB-ON PIU-ON L3
 ANALYZE PIU-ON L4 1- RN : 5098 TERMS
 D
 D
 D 1-20

FILE 'REGISTRY' ENTERED AT 09:37:09 ON 11 OCT 2007
 1 SEA ABB-ON PIU-ON 152044-54-7
 1 SEA ABB-ON PIU-ON 152044-53-6
 1 SEA ABB-ON PIU-ON 189453-10-9
 1 SEA ABB-ON PIU-ON 186692-73-9
 1 SEA ABB-ON PIU-ON 187527-25-9
 1 SEA ABB-ON PIU-ON 188730-08-7
 1 SEA ABB-ON PIU-ON 20949-84-2
 1 SEA ABB-ON PIU-ON 106921-60-2
 1 SEA ABB-ON PIU-ON 193146-27-9
 0 SEA ABB-ON PIU-ON (L6 OR L7 OR L8 OR L9 OR L10 OR L11 OR L12 OR L13 OR L14) AND L4
 0 SEA ABB-ON PIU-ON (L6 OR L7 OR L8 OR L9 OR L10 OR L11 OR L12 OR L13 OR L14) AND L3
 1 SEA ABB-ON PIU-ON 187283-46-1
 1 SEA ABB-ON PIU-ON 188999-14-1
 0 S 184246-38-6
 1 SEA ABB-ON PIU-ON 184246-38-6
 1 SEA ABB-ON PIU-ON 189453-35-8
 1 SEA ABB-ON PIU-ON 219989-84-1
 1 SEA ABB-ON PIU-ON 63928-37-0
 1 SEA ABB-ON PIU-ON 52079-23-9
 1 SEA ABB-ON PIU-ON 70113-32-5
 1 SEA ABB-ON PIU-ON 185148-95-2
 1 SEA ABB-ON PIU-ON 186692-84-2
 0 SEA ABB-ON PIU-ON L3 AND (L6 OR L7 OR L8 OR L9 OR L10 OR L11 OR L12 OR L13 OR L14 OR L15 OR L16 OR L17 OR L18 OR L19 OR L20 OR L21 OR L22 OR L23 OR L24 OR L25 OR L26)
 191 SEA ABB-ON PIU-ON L3 AND CASREACT/LC

FILE 'ZCAPLUS' ENTERED AT 09:43:45 ON 11 OCT 2007
 115 SEA ABB-ON PIU-ON L3/P
 FILE 'CASREACT' ENTERED AT 09:53:42 ON 11 OCT 2007
 69 SEA ABB-ON PIU-ON L3
 STRUCTURE UPLOADED
 0 SEA SUB-L30 SSS SAM L31 (0 REACTIONS)
 23 SEA SUB-L30 SSS FUL L31 (468 REACTIONS)
 D STAT QUE L33

FILE 'REGISTRY' ENTERED AT 11:21:02 ON 11 OCT 2007
 22933 SEA ABB-ON PIU-ON OC15/ESS
 27330 SEA ABB-ON PIU-ON C16/ESS
 726 SEA ABB-ON PIU-ON NC15/ESS
 0 SEA ABB-ON PIU-ON NSC14/ESS
 50989 SEA ABB-ON PIU-ON (L34 OR L35 OR L36 OR L37)
 12165 SEA ABB-ON PIU-ON L38 AND CASREACT/LC
 FILE 'CASREACT' ENTERED AT 11:22:32 ON 11 OCT 2007
 2534 SEA ABB-ON PIU-ON L39/PRO
 59 SEA ABB-ON PIU-ON L30 (L) L40
 65 SEA ABB-ON PIU-ON L3/RRT
 59 SEA ABB-ON PIU-ON L42 (L) L40
 19 SEA ABB-ON PIU-ON L43 AND L33
 FILE 'CAPLUS' ENTERED AT 11:34:47 ON 11 OCT 2007
 59 SEA ABB-ON PIU-ON L43
 53 SEA ABB-ON PIU-ON L45 AND PY<2005
 49 SEA ABB-ON PIU-ON L45 AND PY<2004
 FILE 'CASREACT' ENTERED AT 11:35:34 ON 11 OCT 2007
 D L43
 59 SEA ABB-ON PIU-ON L43 AND 1/NS
 50 SEA ABB-ON PIU-ON L43 AND 2/NS
 44 SEA ABB-ON PIU-ON L43 AND 3/NS
 42 SEA ABB-ON PIU-ON L43 AND 4/NS
 9 SEA ABB-ON PIU-ON L48 NOT L49
 D SCA
 FILE 'CAPLUS' ENTERED AT 11:45:06 ON 11 OCT 2007
 45 SEA ABB-ON PIU-ON L45 AND J/DT
 14 SEA ABB-ON PIU-ON L45 AND P/DT
 12 SEA ABB-ON PIU-ON L54 AND PD<20040619
 39 SEA ABB-ON PIU-ON L53 AND ED<20040619
 6 SEA ABB-ON PIU-ON L53 NOT L56
 2 SEA ABB-ON PIU-ON L54 NOT L55
 8 SEA ABB-ON PIU-ON (L57 OR L58)
 SEL AN
 FILE 'CASREACT' ENTERED AT 11:47:33 ON 11 OCT 2007
 8 SEA ABB-ON PIU-ON ("142:134344"/AN OR "143:211773"/AN OR "143:271524"/AN OR "143:422202"/AN OR "144:170808"/AN OR "145:251631"/AN OR "145:397261"/AN OR "146:229070"/AN OR "146:251631"/AN OR "2004:985335"/AN OR "2005:1154536"/AN OR "2005:1305128"/AN OR "2005:614221"/AN OR "2006:1337456"/AN OR "2006:667477"/AN OR "2006:805502"/AN)
 51 SEA ABB-ON PIU-ON L48 NOT L60
 42 SEA ABB-ON PIU-ON L49 NOT L60
 36 SEA ABB-ON PIU-ON L50 NOT L60
 35 SEA ABB-ON PIU-ON L51 NOT L60
 FILE 'CAPLUS' ENTERED AT 11:49:27 ON 11 OCT 2007
 E US2006-563058 /APES
 1 SEA ABB-ON PIU-ON US2006-563058 /AP
 D SCA
 SEL RN
 FILE 'REGISTRY' ENTERED AT 11:50:13 ON 11 OCT 2007
 55 SEA ABB-ON PIU-ON (130486-83-0/BI OR 152044-53-6/BI OR 152044-54-7/BI OR 185148-95-2/BI OR 220367-73-7/BI OR 220774-16

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

RX(3) RCT H 188730-19-0

STAGE(1)

RGT J 4136-95-2 2,4,6-Cl₃C₆H₂COCl, P 121-44-8 Et₃N
SOL 109-99-9 THF

STAGE(2)

RGT L 1122-58-3 4-DMAP
SOL 108-88-3 PhMe

PRO I 186692-84-2

NTE Key step

REFERENCE COUNT: 100

THERE ARE 100 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L90 ANSWER 28 OF 28 CASREACT COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 126:251010 CASREACT Full-text

TITLE: Total synthesis of epothilone A: the

macrolactonization approach

AUTHOR(S): Nicolaou, K. C.; Sarabia, Francisco; Ninkovic, Sacha; Yang, Zhen

CORPORATE SOURCE: Dep. Chem., Skaggs Inst. Chem. Biol., Scripps Res.

Inst., La Jolla, CA, 92037, USA

Angewandte Chemie, International Edition in English

(1997), 36(5), 525-527

CODEN: ACIEAY; ISSN: 0570-0833


VCH

PUBLISHER: Journal

DOCUMENT TYPE: English


LANGUAGE: English

GI



AB Epothilone A (I) was prepared via a highly convergent and flexible route with macrolactonization of hydroxy acid II as the key step.

RX(4) OF 4 M ==> N



(4) →

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

RX(4) RCT M 188730-19-0

STAGE(1)

RGT O 4136-95-2 2,4,6-Cl₃C₆H₂COCl, P 121-44-8 Et₃N
SOL 109-99-9 THF

STAGE(2)


RGT Q 1122-58-3 4-DMAP
SOL 108-88-3 PhMe

PRO N 186692-84-2

NTE key step

REFERENCE COUNT: 27

THERE ARE 27 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT



H
YIELD 75%

RX(2) RCT G 201136-77-8

STAGE(1)

RGT I 4136-95-2 2,4,6-Cl₃CGH₂COCl, J 121-44-8 Et₃N
SOL 109-99-9 THF

STAGE(2)


RGT K 1122-58-3 4-DMAP
SOL 108-88-3 PhMe

PRO H 201136-78-9

REFERENCE COUNT:


20 THERE ARE 20 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L90 ANSWER 27 OF 28 CASREACT COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 127:293040 CASREACT Full-text
TITLE: Total Syntheses of Epothilones A and B
AUTHOR(S): Mang, Dongfang; Bertinato, Peter; Balog, Aaron; Su, Dai-Shi; Kamenecka, Ted; Sorensen, Erik; Danishefsky, Samuel J.
CORPORATE SOURCE: Laboratory for Bioorganic Chemistry, Sloan-Kettering Institute for Cancer Research, New York, NY, 10021, USA
SOURCE: Journal of the American Chemical Society (1997), 119(42), 10073-10092
CODEN: JACSAT; ISSN: 0002-7863
PUBLISHER: American Chemical Society
DOCUMENT TYPE: Journal
LANGUAGE: English
GI



AB Convergent, stereocontrolled total syntheses of the microtubule-stabilizing macrolides epothilones A (I; R = H) and B (I; R = Me) have been achieved. Four distinct ring-forming strategies were pursued. Of these four, three were reduced to practice. In one approach, the action of a base on a substance possessing an acetate ester and a nonenolizable aldehyde brought about a remarkably effective macrocyclization simultaneously creating the C2-C3 bond and the hydroxyl-bearing stereocenter at C-3. Alternatively, the 16-membered macrolide of the epothilones could be fashioned through a C12-C13 ring-closing olefin metathesis and through macrolactonization of the appropriate hydroxy acid. The application of a stereospecific B-alkyl Suzuki coupling strategy permitted the establishment of a cis C12-C13 olefin, thus setting the stage for an eventual site- and diastereoselective epoxidation reaction. The development of a novel cyclopropane solvolysis strategy for incorporating the geminal Me groups of the epothilones, and the use of Lewis acid catalyzed diene-aldehyde cyclocondensation (LACDAC) and asym. allylation methodol. are also noteworthy.


RX(3) OF 59 H ==> I...



H

(3)

RX(1) OF 1 A ==> B



(1) →

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

RX(1) RCT A 201136-77-8

STAGE(1)

RGT C 121-44-8 Et3N, D 4136-95-2 2,4,6-Cl3C6H2COCl

SOL 109-99-9 THE

STAGE(2)

RGT E 1122-58-3 4-DMAP

SOL 108-88-3 PMe

PRO B 209260-71-9

REFERENCE COUNT: 40

THERE ARE 40 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L90 ANSWER 26 OF 28 CASREACT COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 128:101936 CASREACT Full-Text

TITLE: Total synthesis of 26-hydroxyepothilone B and related analogs

AUTHOR(S): Nicolaou, K. C.; Ninkovic, Sacha; Finlay, M. Ray V.;

CORPORATE SOURCE: Sarabia, Francisco; Li, Tianhu
Department of Chemistry and Biochemistry, University of California, California, 92093, USA

SOURCE: Chemical Communications (Cambridge) (1997), (24), 2343-2344


CODEN: CHCOFS; ISSN: 1359-7345

PUBLISHER: Royal Society of Chemistry

DOCUMENT TYPE: Journal


LANGUAGE: English

GI



AB A series of 26-substituted epothilones B, e.g. I, were constructed by total synthesis involving a selective Wittig olefination, an aldol reaction and a macrolactonization as key steps.

RX(2) OF 2 G ==> H




(2) →

methodology for the rapid, highly selective and convergent construction of epothilone Bepothilone B and analogs


AUTHOR(S): Nicolaou, K. C.; Hepworth, David; Finlay, M. Ray V.; Paul King, N.; Werschkun, Barbara; Elgot, Antony
 CORPORATE SOURCE: Department of Chemistry, The Skaggs Inst. Chem. Biol., The Scripps Res. Inst., La Jolla, CA, 92037, USA
 SOURCE: Chemical Communications (Cambridge) (1999), (6), 519-520

CODEN: CHCOFS; ISSN: 1359-7345
 PUBLISHER: Royal Society of Chemistry
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 GI




AB During a synthesis of 16-desmethylepothilone B (I) new methods for the convergent and highly stereoselective synthesis of epothilone B and analogs were developed.

RX(18) OF 37 BK + BL ==> BQ



BK



(18)

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

RX(18) RCT BK 226940-48-3, BL 108-88-3

STAGE(1)
 RGT BR 4136-95-2 2,4,6-CL3C6H2COCl, BC 121-44-8 Et3N
 SOL 109-99-9 THF

STAGE(2)
 CAT 1122-58-3 4-DMAP
 SOL 108-88-3 PhMe

PRO BQ 226940-49-4
 NTE key step

REFERENCE COUNT: 13 THERE ARE 13 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L90 ANSWER 25 OF 28 CASREACT COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 129:189151 CASREACT Full-text
 TITLE: Total synthesis of 26-hydroxy-epothilone B and related analogs via a macrolactonization based strategy

AUTHOR(S): Nicolaou, K. C.; Finlay, M. Ray V.; Ninkovic, Sacha; Sarabia, Francisco

CORPORATE SOURCE: Department of Chemistry and The Skaggs Institute for Chemical Biology, The Scripps Research Institute, La Jolla, CA, 92037, USA


SOURCE: Tetrahedron (1998), 54(25), 7127-7166

CODEN: TETRAH; ISSN: 0040-4020

PUBLISHER: Elsevier Science Ltd.

DOCUMENT TYPE: Journal

LANGUAGE: English
 GI



AB The chemical synthesis of a series of 26-substituted epothilones B was described. Fully protected 26-hydroxydesoxy-epothilone B I (R = SiMe2OMe3, R1 = CPh3), prepared via a macrolactonization strategy, served as a common precursor to the designed epothilones described. The synthesized compounds were members of a large epothilone library of a number of antitumor agents.

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

RX(4) RCT T 241129-39-5

STAGE(1)
RGT V 84033-33-8 Benzoyl chloride, trichloro-, W 121-44-8 Et3N
SOL 109-99-9 THF

STAGE(2)
RGT X 1122-58-3 4-DMAP
SOL 108-88-3 PhMe, 109-99-9 THF


PRO U 241129-40-8
REFERENCE COUNT: 12 THERE ARE 12 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L90 ANSWER 23 OF 28 CASREACT COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 131:31829 CASREACT Full-text
TITLE: A process for the preparation of ring-opened
epothilone intermediates which are useful for the
preparation of epothilone analogs


INVENTOR(S): Kim, Soong-Hoon; Borzilleri, Robert M.
PATENT ASSIGNEE(S): Bristol-Myers Squibb Company, USA
SOURCE: PCT Int. Appl., 20 pp.
CODEN: PIXXDZ

DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9927890	A2	19990610	WO 1998-US25408	19981130
W:	AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW			
RW:	GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LJ, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
US 6365749	B1	20020402	US 1998-170582	19981013
CA 2312098	A1	19990610	CA 1998-2312098	19981130
EP 1035824	A1	20000920	EP 1998-960564	19981130
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI			
AU 739380	B2	20011011	AU 1999-16134	19981130
JP 2003522722	T	20030729	JP 2000-522878	19981130
ZA 9810993	A	20000601	ZA 1998-10993	19981201
PRIORITY APPLN. INFO.:			US 1997-67550P	19971204
OTHER SOURCE(S):			WO 1998-US25408	19981130
GI			MAPAT 131:31829	



AB A process to produce ring opened epothilones (I), (NR1R2 = N3, (un)substituted amine) and their use in the preparation of epothilone analogs (II) is presented. Thus, epothilone B is cleaved with NaN3, azide reduced to amine and macrolactamized with diphenylphosphoryl azide to give II in 40% yield.




* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *


RX(3) RCT F 219990-25-7
RGT K 26386-88-9 (PRO)2P(O)N3
PRO J 219989-84-1
SOL 68-12-2 DMF

L90 ANSWER 24 OF 28 CASREACT COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 131:31819 CASREACT Full-text
TITLE: Synthesis of 16-desmethylepothilone B: improved

PAGE 1-A



PAGE 1-B



D

(2) →

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

RX(2) RCT D 193146-51-9

PRO B 152044-54-7

NTE lit. ref.

REFERENCE COUNT: 82

THERE ARE 82 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L50 ANSWER 22 OF 28 CASREACT COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER:

131:286299 CASREACT Full-text

TITLE: New Chemical Synthesis of the Promising Cancer

Chemotherapeutic Agent 12,13-Desoxyepothilone B:

Discovery of a Surprising Long-Range Effect on the

Diastereoselectivity of an Aldol Condensation

Harris, Christina R.; Kuduk, Scott D.; Balog, Aaron;

Savin, Ken; Glunz, Peter W.; Danishefsky, Samuel J.

Laboratory for Bioorganic Chemistry, The

Sloan-Kettering Institute for Cancer Research, New


York, NY, 10021, USA

155

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB The epothilones are naturally occurring cytotoxic mols. that possess the remarkable ability to arrest cell division through the stabilization of microtubule assemblies. In vivo studies with 12,13-desoxyepothilone B (dEpoB) (1), have established that the desoxy compound is well tolerated and virtually curative against a variety of sensitive and resistant xenograft tumors in animal models. In light of these discoveries, a chemical synthesis of dEpoB would be able to support a serious and substantial discovery research program directed toward the clin. development of this mol. The overall strategy for this endeavor assumed the ability to synthesize dEpoB from three constructs which include an achiral β,δ -diketo ester construct A (II), an (S)-2-methylpentenal moiety B (III), and the thiazoyl-containing vinyl iodide moiety C (IV). It was envisioned that a diastereoselective aldol condensation between an achiral C5-C6 (Z)-metalenoate derived from construct A and an (S)-2-methylalkanal fragment, B, would generate the desired C6-C7 bond. Second, a B-alkyl Suzuki coupling between the vinyl iodide construct C and an alkyl borane would form the C11-C12 bond. Finally, a late-stage reduction of the C3 ketone to the requisite C3 alc. with high asym. induction would permit introduction of the β,δ -diketo ester fragment A, into the synthesis as a readily accessible achiral building block. The governing concepts the new synthesis are described.

RX(4) OF 5 T ==> D



(4) →

156

SOL 109-99-9 THE

STAGE(2)

RGT AF 1122-58-3 4-DMAP

SOL 108-88-3 PhMe

PRO AC 186692-84-2

REFERENCE COUNT: 41

THERE ARE 41 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L90 ANSWER 21 OF 28 CASREACT COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 131.351125 CASREACT Full-text

TITLE: Syntheses of (-)-epothilone B

AUTHOR(S): Schinzer, Dieter; Bauer, Armin; Schieber, Jennifer

CORPORATE SOURCE: Chemisches Institut der Otto-von-Guericke-Universität,

Magdeburg, D-39106, Germany

SOURCE: Chemistry--A European Journal (1999), 5(9), 2492-2500


CODEN: CEUJED; ISSN: 0947-6539

PUBLISHER: Wiley-VCH Verlag GmbH

DOCUMENT TYPE: Journal


LANGUAGE: English

GI



AB The enantioselective total synthesis of epothilone A was achieved via the catalytic coupling of I and II. The key step in the preparation of I was the catalytic cyansilylation of III. II was prepared via a catalytic organic acetalization followed by an aldol reaction.

RX(5) OF 6 ...C ==> AC



C

(3) →


* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

RX(5) RCT C 188730-19-0

STAGE(1)

RGT AD 4136-95-2 2,4,6-Cl3C6H2COCl, AE 121-44-8 Et3N

153



AB Two efficient routes for the total synthesis of (-)-epothilone B (I) are reported. One strategy is based on ring-closing metathesis, and a second synthesis on a macrolactonization. The key fragments are available on large scale to provide sufficient material for biol. tests. Thiazole fragment II (TBDMS = SiMe2CMe3) was obtained by an improved route starting from (S)-malic acid. The first synthesis is based on our preceding paper. The critical tri-substituted double bond C12-13 in our second approach was constructed by a highly efficient Pd-mediated coupling reaction. Ring closure was achieved by macrolactonization.

154

STAGE(2)

STAGE(3)

SOL 141-78-6 AcOEt

PRO AL 219989-84-1

NTE PHOSPHATE BUFFER USED INSECOND STAGE

REFERENCE COUNT: 80 THERE ARE 80 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L90 ANSWER 19 OF 28 CASREACT COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 133:266631 CASREACT Full-text

TITLE: Total Synthesis of Epothilone A

AUTHOR(S): Zhu, Bin; Panek, James S.

CORPORATE SOURCE: Department of Chemistry and the Center for Streamlined Synthesis Metcalf Center for Science and Engineering, Boston University, Boston, MA, 02215, USA

SOURCE: Organic Letters (2000), 2(17), 2575-2578


CODEN: ORLEF7; ISSN: 1523-7060

PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal


LANGUAGE: English

GI




AB Epothilones A (I) and B are potent antitumor natural products with a Taxol-like mechanism of action. A total synthesis of I is reported, which utilized chiral silane-based bond construction methodology to introduce the key C-6 and C-7 stereocenters of fragment (II). The C-15 stereocenter of fragment (III) was established by a lipase-mediated kinetic resolution. The fragments were assembled with a Suzuki coupling reaction and an aldol condensation and cyclized with a Yamaguchi-type macrocyclization reaction.

RX(4) OF 6 V ==> W



V

(4) →



YIELD 73%

RX(4) RCT V 297131-85-2

STAGE(1)

RGT X 4136-95-2 2,4,6-Cl3C6H2COCl, Y 121-44-8 Et3N

SOL 109-99-9 THF

STAGE(2)

RGT Z 1122-56-3 4-DMAP

SOL 108-88-3 PhMe

PRO W 297131-86-3

NTE stereoselective

REFERENCE COUNT: 36

THERE ARE 36 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L90 ANSWER 20 OF 28 CASREACT COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 132:251011 CASREACT Full-text

TITLE: Enantioselective total synthesis of epothilone A using multifunctional asymmetric catalyses

AUTHOR(S): Sawada, Daiisuke; Shibasaki, Masakatsu

CORPORATE SOURCE: Graduate School of Pharmaceutical Sciences, The

University of Tokyo, Tokyo, 113-0033, Japan

SOURCE: Angewandte Chemie, International Edition (2000),

39(1), 209-213


CODEN: ACHIEF5; ISSN: 1433-7851

PUBLISHER: Wiley-VCH Verlag GmbH

DOCUMENT TYPE: Journal

LANGUAGE: English

GI



(8) →

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

RX(8) RCT T 219990-25-7


STAGE(1)
SOL 68-12-2 DMF

STAGE(2)
RGT D 144-55-8 NaHCO₃, 2 26386-88-9 (PhO)2P(O)N3

PRO I 219989-84-1


REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L90 ANSWER 18 OF 28 CASREACT COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 133:321737 CASREACT Full-text
TITLE: A Novel Application of a Pd(0)-Catalyzed Nucleophilic Substitution Reaction to the Regio- and Stereoselective Synthesis of Lactam Analogues of the Epothilone Natural Products
AUTHOR(S): Borzilleri, Robert M.; Zheng, Xiaoping; Schmidt, Robert J.; Johnson, James A.; Kim, Soong-Hoon; DiMarco, John D.; Fairchild, Craig R.; Gougoutas, Jack Z.; Lee, Francis Y. F.; Long, Byron H.; Vite, Gregory D.
CORPORATE SOURCE: Divisions of Discovery Chemistry Oncology Drug Discovery and Analytical Research and Development, Bristol-Myers Squibb Pharmaceutical Research Institute, Princeton, NJ, 08543-4000, USA
SOURCE: Journal of the American Chemical Society (2000), 122(37), 8890-8897
CODEN: JACSAT; ISSN: 0002-7863
PUBLISHER: American Chemical Society
DOCUMENT TYPE: Journal
LANGUAGE: English
GI



AB Several lactam analogs of the epothilones were prepared using a concise semisynthetic approach starting with the unprotected natural products. Highlighted in this strategy is a novel regio- and stereoselective Pd(0)-catalyzed azidation reaction of a macrocyclic lactone. Subsequent reduction and macrolactamization of the resulting azide acid intermediates provided the desired macrolactams in satisfactory overall yields. The entire three-step sequence was streamlined into a "one-pot" process for the epothilone B-lactam, BMS-247550 (I), which is currently undergoing phase I clin. trials. An initial total synthesis route to prepare the lactam analog of epothilone C was completed and compared to the more direct semisynthesis approach. All of the lactam analogs were evaluated in vitro and the results are discussed.

RX(10) OF 115 ...AJ ==> AL...




(10) →

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

RX(10) RCT AJ 219990-25-7

STAGE(1)
RGT AM 26386-88-9 (PhO)2P(O)N3, AB 144-55-8 NaHCO₃
SOL 68-12-2 DMF



CG

(27)

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

RX(27) RCT CG 188730-19-0

STAGE(1)

RGT CS 4136-95-2 2,4,6-Cl₃C₆H₂COCl, BE 121-44-8 Et₃N
SOL 109-99-9 THF

STAGE(2)

RGT BF 1122-58-3 4-DMAP
SOL 108-88-3 PhMe

PRO CR 186692-84-2

NTE STEREOSSELECTIVE

REFERENCE COUNT: 42

THERE ARE 42 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L90 ANSWER 17 OF 28 CASREACT COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 133:362657 CASREACT Full-text
TITLE: A process for the reduction of oxiranyl epothilones to olefinic epothilones

INVENTOR(S): Kim, Soong-Hoon; Johnson, James A.

PATENT ASSIGNEE(S): Bristol-Myers Squibb Co., USA

SOURCE: PCT Int. Appl., 19 pp.

CODEN: PIXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 3

PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2000071521	A1	20001130	WO 2000-US13253	20000515
W:	AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, GR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI,			

SK, SL, TJ, TM, TR, TT, UA, UZ, VN, YJ, ZA, ZW
RW: DK, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, GH, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BE, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG
US 6320045 B1 20011120 US 1999-316796 19990521
CA 2375029 A1 20001130 CA 2000-2375029 20000515
EP 1178968 A1 20020213 EP 2000-930725 20000515
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO
JP 2003500394 T 20030107 JP 2000-619778 20000515
IN 2001-MN1106 A 20070420 IN 2001-MN1106 20010912
MX 2001-PA11053 A 20020722 MX 2001-PA11053 20011030
US 1999-316796 19990521
US 1997-67549P 19971204
US 1998-82563P 19980421
US 1998-170581 19981013
WO 2000-US13253 20000515

OTHER SOURCE(S):

MARPAT 133:362657

GI




AB 12(13)-Olefinic epothilones, such as I and II [R1-6 = H, alkyl, aryl; R1R2 = cycloalkyl; R7 = H, alkyl, aryl, cycloalkyl, heterocyclyl; P1, P2 = H, alkyl, alkanoyl, aryl, silyl, etc.; W = O, NR8; R8 = H, OH, alkyl], were prepared via reduction of the corresponding 12,13-epoxyepothilones using a metal or metal-assisted reagent. The metal or metal-assisted reagent was selected from the group consisting of reactive metalocenes, [N2C(CO2Me)2, cat Rh2(OAc)4], [N2C(CO2Me)2, cat[(n-C7H15CO2)2Rh]2], [Zn-Cu, EtOH], [Mg(Hg), MgBr], Cr, [FeCl3, n-BuLi], [TiCl3, LiAlH4], [TiCl4, Zn], [WCl6, LiAlH4], [NbCl5, NaAlH4], [VCl3, Zn], or [WCl6, n-BuLi]. Thus, epothilone A, a 12,13-epoxyepothilone, was reduced using magnesium turnings and titanocene dichloride in THF to give epothilone C, a 12(13)-(2)-olefin, in 80% yield.

RX(8) OF 18 ...F ==> I...

efficient and was amenable to the production of significant quantities of these lactams. Using our fully synthetically derived lactams, in vitro and in vivo studies were conducted in comparison with advanced clin. candidates, 12,13-desoxyepothilone B and 12,13-desoxyepothilone F, also derived by total synthesis.

RX(2) OF 2 2 J ==> K + L



2 J

(2) ==>

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

RX(2) RCT J 350042-12-5

STAGE(1)

RGT M 7087-68-5 EN(Pr-1)2, N 148893-10-1 1H-1,2,3-Triazolo[4,5-b]pyridinium, 1-bis(dimethylamino)methylene]-, hexafluorophosphate(1-), 3-oxide, O 39968-33-7
3H-1,2,3-Triazolo[4,5-b]pyridine, 3-hydroxy-
SOL 68-12-2 DMF, 75-09-2 CH2Cl2

STAGE(2)

RGT P 64-19-7 AcOH
SOL 7732-18-5 Water, 109-99-9 THF

PRO X 277749-43-6, L 350042-20-5

REFERENCE COUNT: 29 THERE ARE 29 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

LSO ANSWER 16 OF 28 CASREACT COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 134:56502 CASREACT Full-text

TITLE: Enantioselective Total Synthesis of Epothilones A and

B Using Multifunctional Asymmetric Catalysis

AUTHOR(S): Sawada, Daisuke; Kanai, Motomu; Shibasaki, Masakatsu

CORPORATE SOURCE: Graduate School of Pharmaceutical Sciences, The

145

University of Tokyo, Bunkyo-ku Tokyo, 113-0033, Japan
SOURCE: Journal of the American Chemical Society (2000),
122(43), 10521-10532


CODEN: JACSAT; ISSN: 0002-7863

PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal

LANGUAGE: English

GI



AB An enantioselective total synthesis of epothilones A and B using multifunctional asym. catalysis such as a cyanosilylation of an aldehyde, an aldol reaction of an unmodified ketone with an aldehyde, and a protonation in the conjugate addition of a thiol to an α,β -unsatd. thioester has been achieved. Epothilones A and B were divided into fragment A (I), fragment B (II), and fragment C (III). A catalytic asym. synthesis of fragments A and B was accomplished using a catalytic asym. cyanosilylation as a key step. An enantiocontrolled synthesis of fragment C was achieved in two ways. One is the use of a direct catalytic asym. aldol reaction of an unmodified ketone with an aldehyde as a key step, and the other utilizes a catalytic asym. protonation in the conjugate addition of a thiol to an α,β -unsatd. thioester as a key step. Suzuki cross-coupling of fragment A with fragment C followed by Yamaguchi lactonization as key steps led to an enantiocontrolled synthesis of epothilone A. On the other hand, Suzuki cross-coupling of fragment B with fragment C followed by Yamaguchi lactonization accomplished an enantiocontrolled synthesis of epothilone B.

RX(27) OF 319 ...CG ==> CR

146

TITLE:

Methodology based on chiral silanes in the synthesis of polypropionate-derived natural products - total synthesis of epothilone A

Zhu, Bin; Panek, James S.

R. W. Johnson Pharmaceutical Research Institute, Raritan, NJ, 08869, USA

European Journal of Organic Chemistry (2001), (9), 1701-1714

CODEN: EJOCFK; ISSN: 1434-193X

Wiley-VCH Verlag GmbH

Journal

English

GI

AUTHOR(S):


CORPORATE SOURCE:

SOURCE:

PUBLISHER:


DOCUMENT TYPE:

LANGUAGE:




AB Epothilones A and B are natural products with potent antitumor activity. These compounds have a taxol-like mechanism of action against tumor cells. A total synthesis of epothilone A is reported, which is based on the synthesis and union of two advanced fragments: C3-C11 fragment I and C12-C21 fragment II. Bond construction methodol. based on chiral silanes was utilized to introduce the key C6 and C7 stereocenters of fragment I. A lipase-mediated kinetic resolution established the C15 stereocenter of fragment II. The 16-membered lactone was constructed using a three-step sequence: an intermol. B-alkyl Suzuki coupling of I and II, an aldol condensation, and a Yamaguchi-type macrolactonization reaction.

RX(6) OF 7 AD ==> AE



AD

(6) →



RX(6) RCT AD 297131-85-2

RGT AF 121-44-8 Et3N, AG 4136-95-2 2,4,6-Cl3C6H2COCl

PRO AE 297131-86-3

SOL 109-99-9 THF

REFERENCE COUNT: 43

THERE ARE 43 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L90 ANSWER 15 OF 28

CASREACT COPYRIGHT 2007 ACS on STN

135:107175 CASREACT Full-text

On the Interactivity of Complex Synthesis and Tumor

Pharmacology in the Drug Discovery Process: Total

Synthesis and Comparative in Vivo Evaluations of the

15-Aza Epothilones

AUTHOR(S):

Stachel, Shawn J.; Lee, Chul Bum; Spassova, Maria;

Chappell, Mark D.; Bornmann, William G.; Danishefsky,

Samuel J.; Chou, Ting-Chao; Guan, Yongbiao

Laboratories for Bioorganic Chemistry Preclinical

Pharmacology and the Preparative Synthesis Core

Facility, The Sloan-Kettering Institute for Cancer,

Research, New York, NY, 10021, USA

JOURNAL OF ORGANIC CHEMISTRY (2001), 66(12), 4369-4378

CODEN: JOCEAH; ISSN: 0022-3263

AMERICAN CHEMICAL SOCIETY

PUBLISHER:

DOCUMENT TYPE:

LANGUAGE:

English

AB The total syntheses of 12,13,15-desoxy-15(S)-aza-epothilone B (aza-dEpoB);

dEpoB-lactam) and 12,13,15-desoxy-15(R)-aza-epothilone B (15-epi-aza-dEpoB);

15-epi-dEpoB-lactam) have been accomplished via a highly convergent strategy.

We have also successfully oxidized 12,13,15-desoxy-15(S)-aza-epothilone B to

aza-epothilone B (aza-EpoB; EpoB-lactam). Aza-epothilone B has been advanced

to phase I clin. trials by the Bristol-Myers Squibb group. Our synthesis is

PRO K 219989-84-1

NTE alternative prepn. gave lower yields

REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L90 ANSWER 13 OF 28 CASREACT COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER:

135:180640 CASREACT Full-text

TITLE: The 12,13-diol cyclization approach for a truly stereocontrolled total synthesis of epothilone B and the synthesis of a conformationally restrained analog

AUTHOR(S):

Martin, Harry J.; Pofarliev, Peter; Kahlig, Hanspeter; Mulzer, Johann

CORPORATE SOURCE:

Institut für Organische Chemie der Universität Wien, Vienna, 1090, Austria

SOURCE:

Chemistry--A European Journal (2001), 7(10), 2261-2271

PUBLISHER:

Wiley-VCH Verlag GmbH

DOCUMENT TYPE:

Journal

LANGUAGE:


English

GI

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB A highly convergent and stereocontrolled synthesis of epothilone B (I) has been developed. The epoxide moiety in I was generated by regioselective mesylation and base treatment of the 12,13-diol II which was formed by a chelate Cram controlled Grignard addition of (3S)-Br(CH₂)₃CHMeCH:CH₂ and Me ketone III. Both fragments were synthesized from the chiral carbon pool precursors (S)-citronellol and (S)-lactic acid, resp. A highly diastereoselective aldol addition of epoxy-aldehyde IV and the known Southern hemisphere ketone (S)-MeCH₂COCH₂Me₂CH(OSiMe₂OMe)₃CH₂CH:CH₂ delivered the full carbon skeleton, containing all the stereogenic centers of I. Functional group manipulation, macrolactonization and removal of two protecting groups then yielded I. The spatial closeness of the C4-β-Me and C6-Me group in the crystal structure of I inspired the authors to connect them through a methylene bridge to give a cyclohexanone derivative. Thus, the Northern hemisphere aldehyde IV was added to the enolate of a cyclohexanone derivative. Further manipulations and macrolactonization delivered the conformationally restrained epothilone derivative V.

RX(2) OF 2 G ==> H



G

(2) →

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

PAGE 1-B

-Bu-t

PAGE 2-A

FIELD 654

RX(2) RCT G 263761-19-9
RGT I 121-44-8 Et3N, J 4136-95-2 2,4,6-Cl3C6H2COCl, K 1122-58-3

4-DMAP

PRO H 263761-23-5

SOL 108-88-3 PhMe

REFERENCE COUNT: 62

THERE ARE 62 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L90 ANSWER 14 OF 28 CASREACT COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 135:137326 CASREACT Full-text

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

RX(8) RCT X 219990-25-7
 RGT AC 26386-88-9 (PHO)2P(O)N3, J 144-55-8 NaHCO3
 PRO AB 219999-84-1
 SOL 68-12-2 DMF

REFERENCE COUNT: 88
 THERE ARE 88 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L90 ANSWER 12 OF 28 CASREACT COPYRIGHT 2007 ACS on STN
 135:257087 CASREACT Full-text
 TITLE: A process for the preparation of epothilone analogs and intermediates

INVENTOR(S): Li, Wen Sen; Thornton, John E.; Guo, Zhenrong;
 Skamirathan, Shankar; McConlogue, Gary W.

PATENT ASSIGNEE(S): Bristol-Myers Squibb Company, USA
 SOURCE: PCT Int. Appl., 41 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English


FAMILY ACC. NUM. COUNT: 4

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001070716	A1	20010927	WO 2001-US7749	20010312
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, GU, HD, IL, IN, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PK, PL, PT, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
CA 2404212	A1	20010927	CA 2001-2404212	20010312
EP 1265878	A1	20021218	EP 2001-918544	20010312
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR			
HU 200300693	A2	20030828	HU 2003-693	20010312
JP 2003528090	T	20030924	JP 2001-568920	20010312
IN 2002MN01074	A	20050304	IN 2002-MN1074	20020808
MX 2002PA09165	A	20040812	MX 2002-PA9165	20020919
PRIORITY APPLN. INFO.:			US 2000-528526	20000320
OTHER SOURCE(S):			WO 2001-US7749	20010312

MARPAT 135:257087



GI



139

AB The present invention relates to a process for the preparation of epothilone analogs, such as I (R₁ = NH), by initially forming novel ring-opened epothilones and carrying out a macrolactamization reaction thereon. The subject process is amenable to being carried out in a single reaction vessel without isolation of the intermediate compound and provides at least about a three-fold increase in yield over prior processes for preparing the desired epothilone analogs. Thus, ring opening of epothilone B was achieved using NaN₃, PMA3 and Bu₄N⁺-Cl⁻ in THF in the presence of Pd₂(dba)₃.CHCl₃ to form TBA salt I (R = NH₂, R₁ = O⁻Bu₄N⁺) in 93% yield. The ring opened epothilone B TBA salt I underwent intramolecular macrolactamization using K₂CO₃, HOBT, and EDCI in THF and DMF to form lactam I (R₁ = NH) in 92.7% yield.

RX(2) OF 4 ...C ==> K



(2)

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

RX(2) RCT C 361204-09-3

STAGE(1)

RGT L 1310-58-3 KOH

SOL 109-99-9 THF, 68-12-2 DMF

STAGE(2)

RGT M 2592-95-2 1-Benzotriazolol, N 25952-53-8 EDAP

140

SOL 109-99-9 THE

STAGE(2)



RGT AM 1122-56-3 4-DNAP
SOL 108-88-3 PhMe

PRO AU 241129-40-8

L90 ANSWER 11 OF 28 CASREACT COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 135:371566 CASREACT Full-text
TITLE: Process for reduction of oxiranyl epothilones to
olefinic epothilones
INVENTOR(S): Kim, Seong-hoon; Johnson, James A.
PATENT ASSIGNEE(S): Bristol-Myers Squibb Co., USA
SOURCE: U.S., 10 pp., Cont.-in-part of U.S. Ser. No. 170,581.
CODEN: USXXAM


DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 3
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 6320045	B1	20011120	US 1998-316796	19990521
CA 2375029	A1	20001130	CA 2000-2375029	20000515
WO 2000071521	A1	20001130	WO 2000-US13253	20000515
W: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, GR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, LC, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW				
RW: GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GN, GW, ML, MR, NE, NG, SN, TD, TG				
EP 1178968	A1	20020213	EP 2000-930725	20000515
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
HU 2002001467	A2	20021028	HU 2002-1467	20000515
JP 2003500394	T	20030107	JP 2000-619778	20000515
IN 2001MN1106	A	20070420	IN 2001-MN1106	20010912
MX 2001PA11053	A	20020722	MX 2001-PA11053	20011030
PRIORITY APPLN. INFO.:				
US 1997-67549P 19971204				
US 1998-82563P 19980421				
US 1998-170581 19981013				
US 1999-316796 19990521				
WO 2000-US13253 20000515				
OTHER SOURCE(S): MARPAT 135:371566				
GI				



AB This process produced epothilones I (W = O, NR8; R1-R6 = H, (un)substituted alkyl or aryl and R1 and R2 can be cycloalkyl; R7 = H, (un)substituted alkyl, aryl, cycloalkyl or 4-7 membered heterocyclic N-, O-, or S-containing rings; R8 = H, (un)substituted alkyl, OH, (un)substituted O-alkyl; X = CH2 or XY = CH=CH; Z = H or Op1 where P1, P2 = H, (un)substituted alkyl, alkanoyl, acetyl, trialkyl(aryl)silyl) from oxiranyl epothilones via the reaction of the oxiranyl moiety with a metal or metal-assisted reagent selected from the group consisting of reactive metalocenes, or (WCl6, n-Buli). Thus II was prepared in 29% yield in a multistep reaction from epothilone B via the aminoheptadecenoic acid that cyclized to the oxiranyl azepothilone intermediate which was reacted with WCl6 in THF and n-Buli in hexane.


RX(8) OF 16 ...X ==> AB...



(8)

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002012534	A2	200202014	WO 2001-US25112	20010809
WO 2002012534	A3	200209006		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, GR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW				
RW: GE, GM, KE, ME, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
WO 2001092991	A2	20011206	WO 2001-US17352	20010529
WO 2001092991	A3	20020808		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, GR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW				
RW: GE, GM, KE, ME, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
AU 2001075012	A5	20011211	AU 2001-75012	20010529
CA 2417358	A1	20020214	CA 2001-2417358	20010809
AU 2001083275	A5	20020218	AU 2001-83275	20010809
EP 1307579	A2	20030507	EP 2001-962062	20010809
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR				
JP 2004520008	T	20040708	JP 2002-517818	20010809
PRIORITY APPLN. INFO.:				
US 2000-224038P 20000809				
US 2000-237382P 20001004				
US 2000-248387P 20001113				
US 2001-867845 20010529				
US 2000-207331P 20000530				
WO 2001-US17352 20010529				
WO 2001-US25112 20010809				


OTHER SOURCE(S): MARPAT 136:183657
GI



AB The present invention relates to compds., such as I, made by a subset of modules from one or more polyketide synthase ("PKS") genes that are used as starting material in the chemical synthesis of novel mol's., particularly naturally occurring polyketides or derivs. thereof. The biol. derived intermediates ("bio-intermediates") generally represent particularly difficult compds. to synthesize using traditional chemical approaches due to one or more


stereocenters. In one aspect of the invention, an intermediate in the synthesis of epothilone is provided that feeds into the synthetic protocol of Danishefsky and co-workers. In another aspect of the invention, intermediates in the synthesis of discodermolide are provided that feed into the synthetic protocol of Smith and co-workers. By taking advantage of the inherent stereochem. specificity of biol. processes, the syntheses of key intermediates and thus the overall syntheses of compds. like epothilone and discodermolide are greatly simplified.

RX(12) OF 142 ...AQ ----> AU...



AQ

(12) →





AU

RX(12) RCT AQ 241129-39-5

STAGE(1)

RCT AV 4136-95-2 2,4,6-Cl3C6H2COCl, E 121-44-8 Et3N

$$\text{RX}(1) \text{ OF } 1 \quad A + B_{\text{max}} > C$$


* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

PAGE 2-A

RCT	A	289501-50-4	B	75-25-2
RGT	D	1310-73-2		NaOH
PRO	C	402476-95-3		
CAT	56-37-1	PhCH ₂ NEt ₃	Cl	
SOL	7732-18-5	Water	64-17-5	EtOH

L190 ANSWER 10 OF 28 CASREACT COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 136:183657 CASREACT Full-text
TITLE: Process for the biomediated prepa

INVENTOR(S): Santi, Daniel V.; Ashley, Gary; Myles, David C.
PATENT ASSIGNEE(S): Kosan Biosciences, Inc., USA
SOURCE: PCT Int. Appl., 129 pp.

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT:

133

134

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

RX(2) RCT C 361204-09-3

STAGE(1)
RGT K 584-08-7 K2C03
SOL 109-99-9 THF, 68-12-2 DMF



STAGE(2)
RGT L 2592-95-2 1-Benzotriazolol, M 25952-53-8 EDAP

PRO J 219989-84-1
NTE alternative preps. gave lower yields

L90 ANSWER 8 OF 28 CASREACT COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 136:318824 CASREACT Full-text
TITLE: Synthetic and semisynthetic analogs of epothilones:
chemistry and biological activity
AUTHOR(S): Altmann, Karl-Heinz; Blommers, Marcel J. J.;
Caravatti, Giorgio; Florsheimer, Andreas; Nicolaou,
Kyriacos C.; O'Reilly, Terrence; Schmidt, Alfred;
Schinzer, Dieter; Wartmann, Markus
CORPORATE SOURCE: TA Oncology Research, Novartis Pharma AG, Basel,
CH-4002, Switz.
SOURCE: ACS Symposium Series (2001), 796(Anticancer Agents),
112-130
CODEN: ACSMC8; ISSN: 0097-6156
PUBLISHER: American Chemical Society
DOCUMENT TYPE: Journal
LANGUAGE: English

AB Epothilones A and B are naturally occurring microtubule depolymerizing inhibitors, which exhibit potent in-vitro antiproliferative activity. Epothilone B is a 3-30-fold more potent inhibitor of human cancer cell growth than paclitaxel in paclitaxel-sensitive cancer cell lines and in paclitaxel-resistant lines exceeds paclitaxel activity by 102 - 103-fold. In addition, epothilone B exhibits potent in vivo antitumor activity even in multidrug-resistant tumor models. In order to gain a better understanding of the structural requirements for epothilone-mediated cytotoxicity and antitumor activity and to discover analogs with similar potency but perhaps better tolerability in vivo, we have investigated a series of structural modifications involving the epoxide site (C12/C13) and the heterocyclic side-chain of epothilones. In this paper we present the synthesis of these analogs and we discuss the impact of such modifications on tubulin polymerization activity as well as cytotoxicity in vitro.

RX(32) OF 320 ...CP ==> CR...



CR

RX(32) RCT CP 335160-11-7

STAGE(1)
RGT CS 429-41-4 Bu4N.F
SOL 109-99-9 THF

STAGE(2)
RGT CT 4136-95-2 2,4,6-Cl3C6H2OCl, AP-121-44-8 Et3N, CA
1122-58-3 4-DMAP
SOL 109-99-9 THF, 108-88-3 PhMe

STAGE(3)
RGT R 76-05-1 F3CCO2H
SOL 75-09-2 CH2Cl2

PRO CR 188260-10-8

REFERENCE COUNT: 32 THERE ARE 32 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE REFORMAT

L90 ANSWER 9 OF 28 CASREACT COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 136:216592 CASREACT Full-text
TITLE: Procedures for the production of 12,13-cyclopropylepothilone derivatives, as well as for their use in pharmaceutical preparations
PATENT ASSIGNEE(S): Schering AG, Germany
SOURCE: Ger. Offen., 64 pp.
CODEN: GWXXBX
DOCUMENT TYPE: Patent
LANGUAGE: German
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

PAGE 2-A



I
YIELD 33%

RX(30) RCT DH 472962-13-3
RGT AN 4136-95-2 2,4,6-CL3C6H2COCl, AO 121-44-8 Et3N, AP 1122-58-3 4-DMAP
PRO I 472962-14-4
SOL 109-99-9 THF, 108-88-3 PhMe
NTE stereoselective, Yamauchi macrocyclization
REFERENCE COUNT: 31 THERE ARE 31 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L90 ANSWER 7 OF 28 CASREACT COPYRIGHT 2007 ACS ON STN
ACCESSION NUMBER: 137:140388 CASREACT Full-text
TITLE: A process for the preparation of epothilone analogs and intermediates
INVENTOR(S): Li, Wen-Sen; Thornton, John E.; Guo, Zhenrong; Swaminathan, Shankar
PATENT ASSIGNEE(S): Bristol-Myers Squibb Company, USA
SOURCE: PCT Int. Appl., 41 pp.
CODEN: PIXX22

DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 4
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002060904	A2	20020808	WO 2002-US1853	20020122
WO 2002060904	A3	20030227		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, GR, GU, HK, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PK, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, SM, SN, ST, SV, SZ, TD, TG, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW				
RW: GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GN, GW, HT, IL, IN, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PK, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, SM, SN, ST, SV, SZ, TD, TG, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW				
US 6518421 B1 20030211 US 2001-775361 20010201				
US 2003004338 A1 20030102 US 2001-946721 20010905				
AU 2002240014 A1 20020812 AU 2002-240014 20020122				
PRIORITY APPL. INFO.: US 2001-775361 20010201				
US 2000-528526 20000320				
WO 2002-US1853 20020122				
OTHER SOURCE(S): MAREPAT 137:140388				



129

GI



AB The present invention relates to a process for the preparation of epothilone analogs, such as I [R1 = NH (III)], by initially forming novel ring-opened epothilones and carrying out a macrolactamization reaction thereon. The subject process is amenable to being carried out in a single reaction vessel without isolation of the intermediate compound and provides at least about a three-fold increase in yield over prior processes for preparing the desired epothilone analogs. Thus, ring opening of epothilone B was achieved using NaN3, PMe3 and Bu4NCl in THF in the presence of Pd2(dba)3·CHCl3 to form TBA salt I [R = NH2, R1 = O-Bu4N+ (III)] in 93% yield. III underwent intramol. macrolactamization using K2CO3, HOBt, and EDCI in THF and DMF to form II in 92.7% yield.


RX(2) OF 4 ...C ==> J



C: CM 2

(2) →

130



(2) →


* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

RX(2) RCT C 361204-09-3
 RGT K 584-08-7 K2CO3, L 2592-95-2 1-Benzotriazolol, M 25952-53-8
 EDAP
 PRO J 219989-84-1
 SOL 109-99-9 THF, 68-12-2 DMF
 CON SUBSTAGE(1) room temperature → -5 deg C
 SUBSTAGE(2) 5 minutes, -5 deg C
 SUBSTAGE(3) 2 hours, -5 deg C
 SUBSTAGE(4) 8 hours, 0 deg C
 SUBSTAGE(5) 2 hours, 10 deg C
 NTE optimization study

REFERENCE COUNT: 98
 THERE ARE 98 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L90 ANSWER 6 OF 28 CASREACT COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 137:310727 CASREACT Full-text
 TITLE: Chemical synthesis and biological evaluation of novel epothilone B and trans-12,13-cyclopropyl epothilone B analogues
 AUTHOR(S): Nicolaou, K. C.; Ritzel, Andreas; Namoto, Kenji; Buoy, Ruben M.; Diaz, J. Fernando; Andreu, Jose M.; Wartmann, Markus; Altmann, Karl-Heinz; O'Brate, Aurora; Giannakakou, Paraskevi
 CORPORATE SOURCE: Department of Chemistry and Skaggs Institute for Chemical Biology, Scripps Research Institute, La Jolla, CA, 92037, USA
 SOURCE: Tetrahedron (2002), 58(32), 6413-6432
 CODEN: TETRAE; ISSN: 0040-4020
 PUBLISHER: Elsevier Science Ltd.
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 GI

127




AB In addition to the total synthesis of the thiomethyl thiazole side chain analog of epothilone B I, a series of related trans-12,13-cyclopropyl epothilone B analogs, e.g. II, was accomplished. While the synthesis of the epothilone B analog I proceeded through a Stille coupling of a vinyl iodide substrate containing the epothilone macrocycle with the appropriate side chain stannane, that of the cyclopropyl analogs involved a convergent strategy in which a Nozaki-Hiyama-Kishi coupling was used as a means of introducing the side chains prior to Yamaguchi macrolactonization and final elaboration to the target mols. The synthesized analogs were subjected to biol. evaluation involving in vitro tubulin polymerization, affinity for the microtubule Taxol binding site and cell cytotoxicity assays. The results identified the methylthio thiazole side chain as a potency enhancing moiety for the epothilones and shed further light on the structure-activity relationships within this important class of chemotherapeutic agents.

RX(30) OF 782 ...DH ==> I...



(30) →

128



YIELD 45%


RX(3) RCT H 514224-63-6
 PRO A 186692-73-9
 CAT 252877-37-5 Synthase, epothilone
 SOL 7732-18-5 Water, 67-68-5 DMSO
 CON 22 hours, 30 deg C, pH 5
 NTE biotransformation, enzymic, recombinant epothilone thioesterase
 domain used, phosphate-buffered soln., product distribution
 depends on reaction conditions
 REFERENCE COUNT: 29 THERE ARE 29 CITED REFERENCES AVAILABLE FOR THIS
 RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L90 ANSWER 5 OF 28 CASREACT COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 138:153369 CASREACT Full-text
 TITLE: Process for the preparation of epothilone analogs
 INVENTOR(S): Li, Wen Sen; Thornton, John E.; Guo, Zhenrong;
 Swaminathan, Shankar
 PATENT ASSIGNEE(S): Bristol-Myers Squibb Company, USA
 SOURCE: U.S., 17 pp., Cont.-in-part of U.S. Ser. No. 528,526.
 CODEN: USXXAM
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 4
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 6518421	B1	20030211	US 2001-775361	20010201
US 2003004338	A1	20030102	US 2001-946721	20010905
WO 2002060904	A2	20020808	WO 2002-US1853	20020122
WO 2002060904	A3	20030227		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN,				
CO, CR, CU, CY, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,				
GM, GR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,				
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NZ, PA, PH,				
PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG,				
US, UZ, VN, YU, ZA, ZM, ZW				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TG, TZ, UG, ZM, ZW, AT, BE, CH,				
CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR,				
BF, BU, CF, CG, CI, CM, GN, GW, HT, IL, IN, JP, KE, KG, KP, KR, KZ, LC, LK, LR,				

AU 2002240014 A1 20020812 AU 2002-240014 20020122
 US 39356 E1 20061017 US 2005-56606 20050211
 PRIORITY APPLN. INFO.: US 2000-528526 20000320
 US 2001-775361 20010201
 US 2001-946721 20010905
 WO 2002-US1853 20020122

OTHER SOURCE(S): MARPAT 138:153369
 GI




AB The present invention relates to a process for the preparation of epothilone analogs by initially forming novel ring-opened epothilones and carrying out a macrolactamization reaction thereon. The subject process is amenable to being carried out in a single reaction vessel without isolation of the intermediate compound and provides at least about a three-fold increase in yield over prior processes for preparing the desired epothilone analogs. Thus, epothilone B I (X = O) was treated with NaN₃ and Bu₄NCl- in THF followed by addition of water PMe₃ in THF and equilibrated to 25° then addition of Pd₂(dba)₃.CHCl₃. The resulting solution was stirred under an argon atmosphere for 19 h. to form ring opened salt II in 96% yield. Salt II was then dissolved in THF and DMF, cooled to -5°, treated with K₂CO₃ and stirred for 5 min before adding HOBT and EDCI then stirring for 2 h at -5° to form lactam I (X =NH) in 56% yield from epothilone B.

RX(2) OF 4 ...C ==> J

n-Bu
 n-Bu-N⁺-Bu-n
 n-Bu
 C: CH 1

RX(38) OF 219 ...DB ==> E...




DB

(38)

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

PAGE 2-A

E
YIELD 33%

RX(38) RCT DB 611168-66-2

STAGE(1)

RGT DJ 4136-95-2 2,4,6-CL3C6H2OOC1, BD 121-44-8 Et3N
SOL 109-99-9 THE
CON 1 hour, 0 deg C

STAGE(2)

RGT CF 1122-58-3 4-DMAP
SOL 108-88-3 PhMe
CON 3 hours, 75 deg C

PRO E 611168-68-4

NTE Yamaguchi reaction, addnl. stereoisomeric reactant present
54 THERE ARE 54 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

REFERENCE COUNT:

L90 ANSWER 4 OF 28 CASREACT COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER:

136:316732 CASREACT Full-text

Epothilone C Macrolactonization and Hydrolysis Are

Catalyzed by the Isolated Thioesterase Domain of

Epothilone Polyketide Synthase

123

AUTHOR(S):

Boddy, Christopher N.; Schneider, Tanya L.; Hotta, Kinya; Walsh, Christopher T.; Khosla, Chaitan
Departments of Chemical Engineering, Chemistry and
Biochemistry, Stanford University, Stanford, CA,
94305-5025, USA

CORPORATE SOURCE:

SOURCE:

Journal of the American Chemical Society (2003),
125(12), 3428-3429

CODEN: JACSAT; ISSN: 0002-7863

PUBLISHER:

American Chemical Society

DOCUMENT TYPE:


Journal

LANGUAGE: English

AB Epothilone C is produced by the combined action of one nonribosomal peptide synthetase (NRPS) and nine polyketide synthase (PKS) modules in a multienzyme system. The final step in the biosynthesis is the thioesterase (TE)-catalyzed cyclization of epothilone from the EPOF protein. It has been unclear whether isolated PKS TE domains could exhibit macrolactonization activity. Here we demonstrate that the excised epothilone TE domain can catalyze the efficient cyclization of the N-acetylcysteine thioester of seco-epothilone C to generate epothilone C ($k_{cat}/K_M = 0.41 \pm 0.03 \text{ min}^{-1} \text{ mM}^{-1}$). The TE domain also catalyzes the hydrolysis of both the N-acetylcysteine thioester of seco-epothilone C ($k_{cat} = 0.087 \pm 0.005 \text{ min}^{-1}$, $K_M = 291 \pm 53 \mu\text{M}$) and that of the epothilone C ($k_{cat} = 0.67 \pm 0.01 \text{ min}^{-1}$, $K_M = 117 \pm 5 \mu\text{M}$) to form seco-epothilone C.

RX(3) OF 8 ...H ==> A...

PAGE 1-A



PAGE 1-B



(3)

124

CON 1 hour, room temperature

STAGE(5)

SOL 60-29-7 Et2O, 75-09-2 CH2Cl2
CON room temperature

STAGE(6)

SOL 7732-18-5 Water
CON room temperature

STAGE(7)

SOL 109-99-9 THF
CON room temperature -> -5 deg C

STAGE(8)

RGT CA 121-44-8 Et3N
CON -5 deg C

STAGE(9)

RGT CB 4136-95-2 2,4,6-Cl3C6H2OOC1
CON SUBSTAGE(1) -5 deg C
SUBSTAGE(2) -5 deg C -> 0 deg C
SUBSTAGE(3) 1 hour, 0 deg C

STAGE(10)

RGT CC 1122-58-3 4-DMAP
SOL 108-88-3 PhMe
CON 4 hours, room temperature

PRO BW 867376-54-3, BX 867376-57-6

NTE sixth stage quench

REFERENCE COUNT: 8 THERE ARE 8 CITED REFERENCES AVAILABLE FOR THIS
: RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L90 ANSWER 2 OF 28 CASREACT COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 143:211773 CASREACT Full-text

TITLE: Method for synthesis of Epothilone B lactam derivative

INVENTOR(S): Yan, Jialin

PATENT ASSIGNEE(S): Peop. Rep. China

SOURCE: Faming Zhuanli Shengqing Gongkai Shuomingshu, No pp.

CODEN: CNXEV

DOCUMENT TYPE: Patent

LANGUAGE: Chinese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:


PATENT NO. KIND DATE APPLICATION NO. DATE

CN 1554659 A 20041215 CN 2003-10112901 20031225

PRIORITY APPL. INFO.: CN 2003-10112901 20031225

AB Epothilone B lactam derivative was synthesized from Epothilone B via regional and stereo selective nitridation of the Epsilon B macrolide catalyzed by palladium tri-Ph phosphine. Epothilone B was first ring opened via nitridation reaction to obtain nitronic acid, then processed with tri-Ph phosphine to produce imino phosphorane, later hydrolyzed with ammonium hydroxide to form amino acid, and finally the amino acid was cyclized with DPPA and solid sodium bicarbonate to obtain the target product Epothilone B lactam derivative

RX(3) OF 6 ...G ==> J



(3) →

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

RX(3) RCT G 219990-25-7
RGT K 144-55-8 NaHCO3, L 26386-88-9 (PhO)2P(O)N3
PRO J 219989-84-1

SOL 68-12-2 DMF

CON 24 hours, 4 deg C

L90 ANSWER 3 OF 28 CASREACT COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 139:301287 CASREACT Full-text

TITLE: Design, synthesis, and biological properties of highly potent epothilone B analogues

AUTHOR(S): Nicolaou, K. C.; Sasmal, Pradip K.; Rassias, Gerassimos; Reddy, Mali Venkat; Altmann, Karl-Heinz; Wartmann, Markus; O'Brate, Aurora; Giannakakou, Paraskevi

CORPORATE SOURCE: Department of Chemistry, The Skaggs Institute for Chemical Biology The Scripps Research Institute, La Jolla, CA, 92037, USA

SOURCE: Angewandte Chemie, International Edition (2003), 42(30), 3515-3520

CODEN: ACTEF5; ISSN: 1433-7851

PUBLISHER: Wiley-VCH Verlag GmbH & Co. KGaA



DOCUMENT TYPE: Journal

LANGUAGE: English


AB Epothilones have potent cytotoxicity against tumor cells. We directed our attention toward the synthesis and evaluation of a small designed library of epothilone B analogs. From the library, we found that 12,13-cis-cyclopropane methylsulfanyl epothilone B is extremely potent.

(TBDMS = SiMe₂OMe₃) was prepared from camphosultam V via protection with novel reagent, 4-((trimethylsilyl)methyl)benzyl trichloroacetimidate, basic hydrolysis with LiOH in aqueous THF, stereoselective aldol reaction with undecadienal VI in THF containing LDA, silylation with CF₃SO₂SiMe₂OMe₃ in CH₂Cl₂ containing 2,6-lutidine and regioselective desilylation with MgBr₂ in Et₂O/MeNO₂ containing BuSH, and macrolactonization with 2,4,6-Cl₃C₆H₂COCl in THF containing Et₃N followed by DMAP in PhMe.

RX(28) OF 58 ...BQ + BR ==> BW +
EX



PAGE 1-B



PAGE 1-B



(28) →

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

PAGE 2-A

BW
FIELD 17*



* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

PAGE 2-A

BX
FIELD 33*



RX(28)

STAGE(1)
RGT BY 7789-48-2 MgBr₂
SOL 60-29-7 Et₂O
CON room temperature

STAGE(2)
SOL 75-52-5 MeNO₂
CON room temperature

STAGE(3)
RGT BZ 109-79-5 BuSH
CON room temperature

STAGE(4)
RGT BQ 867376-53-2, BR 867376-58-7
SOL 60-29-7 Et₂O

STAGE(2)
RGT 2 1122-58-3 4-DMAP
SOL 108-88-3 PhMe

PRO AH 240816-03-9
NTE key step

REFERENCE COUNT: 60 THERE ARE 60 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L109 ANSWER 7 OF 7 CASREACT COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 127:108793 CASREACT Full-text

TITLE: Stereoselective syntheses and evaluation of compounds in the 8-desmethylleptothione A series: some surprising observations regarding their chemical and biological properties

AUTHOR(S): Balog, Aaron; Betinato, Peter; Su, Dai-Shi; Meng, Dongfang; Sorensen, Erik; Danishefsky, Samuel J.; Zheng, Yu-Huang; Chou, Ting-Chao; He, Lifeng; Horwitz, Susan B.

CORPORATE SOURCE: Lab. Bioorganic Chem., Sloan-Kettering Inst. Cancer

SOURCE: Res., New York, NY, 10021, USA

CODEN: TELEAY; ISSN: 0040-4039

PUBLISHER: Tetrahedron Letters (1997), 38(26), 4529-4532


DOCUMENT TYPE: Elsevier

LANGUAGE: English

AB The title compds. have been synthesized in a convergent way by recourse to a Weller type dianion construction.

RX(3) OF 15

...I ==> L...



(3)

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

117

RX(3) RCT I 122370-80-2
RGT M 538-75-0 DCC, N 1122-58-3 4-DMAP
PRO L 122370-81-3
SOL 67-66-3 CHCl3

REFERENCE COUNT: 18 THERE ARE 18 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L90 ANSWER 1 OF 28 CASREACT COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 143:422202 CASREACT Full-text

TITLE: Novel protecting reagents, protecting groups and methods of forming and using the same

INVENTOR(S): Avery, Mitchell A.; Chittiboyina, Amar Gopal; Chada, Raji Reddy; Kache, Rajashaker; Jung, Jae Chul

PATENT ASSIGNEE(S): The University of Mississippi, USA

SOURCE: PCT Int. Appl., 82 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO. KIND DATE APPLICATION NO. DATE

WO 2005100329 A1 20051027 WO 2005-US9525 20050323

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, GU, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW
RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, BG, KZ, MD, RU, TJ, TN, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, BU, CF, CG, CI, CM, GN, GW, ML, MR, NE, SN, TD, TG

PRIORITY APPLN. INFO.: US 2004-555896P 20040323

OTHER SOURCE(S): MARPAT 143:422202

GI

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB New protecting reagents TIX-Y (Y = OCNHCCl3, Cl, Br, I, NCO, OCOCl, OCH2Cl, OTs, OMg, ONa, OTf) are provided that allow for the selective placement of a new protecting group onto a reactive site of a multifunctional compound. The TIX-Y reagents are 2, 3, and 4-trialkylsilylyl, triarylsilylyl or a combination of alkyl-aryl silylyl reagents (TIX reagents, I (R, R', R'' = alkyl, aryl), II (R, R', R'' = alkyl, aryl) and III (R, R', R'' = alkyl, aryl)), which carry a TIX protecting group for protecting alcs. as ethers, urethanes, carbonates, acetals; amines as carbamates or ureas; and thiols as ethers or esters. The invention also provides methods of forming the 2, 3, and 4-TIX reagents; introducing the TIX protecting groups to mols. bearing hydroxyl groups, amine groups, or thiol groups; methods of removing the TIX protecting groups; and intermediate compds. formed during any one of these methods. The invention further provides methods useful in producing epothilones and analogs and derivs. thereof. Thus, epothilone derivative IV

118

STAGE(1)

RGT X 4136-95-2 2,4,6-Cl₃C₆H₂COCl, O 121-44-8 Et₃N
SOL 109-99-9 THF

STAGE(2)

RGT Y 1122-58-3 4-DWAP
SOL 108-88-3 PhMe

PRO W 226940-49-4
NTE key step

L109 ANSWER 6 OF 7 CASREACT COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 131.195935 CASREACT Full-text

TITLE: Total synthesis of epothilone E and related side-chain modified analogues via a Stille coupling based strategy

AUTHOR(S):

Nicolaou, K. C.; King, N. P.; Finlay, M. R. V.; He, Y.; Roschangar, F.; Vourloumis, D.; Vallberg, H.; Sarabia, F.; Ninkovic, S.; Hepworth, D.

CORPORATE SOURCE: Department of Chemistry and The Skaggs Institute for Chemical Biology, The Scripps Research Institute, La Jolla, CA, 92037, USA

SOURCE: Bioorganic & Medicinal Chemistry (1999), 7(5), 665-697


CODEN: BMECEP; ISSN: 0968-0896

PUBLISHER: Elsevier Science Ltd.

DOCUMENT TYPE: Journal

LANGUAGE: English

GI




AB A Stille coupling strategy has been utilized to complete a total synthesis of epothilone E from vinyl iodide I (R₁ = I; R₂ = H) and thiazolestannane II. The central core fragment I (R₁ = I; R₂ = H) and its trans-isomer III (R₃ = I) were prepared from triene IV (TBS = SiMe₂OMe₃) using ring-closing metathesis (RCM), and were subsequently coupled to a variety of alternative stannanes to

115

provide a library of epothilone analogs I (R₁ = 2-(5-acetoxypentyl)thiazol-4-yl, 2-(methylthio)thiazol-4-yl, 2-piperidinethiazol-4-yl, 2-methoxythiazol-4-yl, 2-ethoxythiazol-4-yl, thiazol-4-yl, thiazol-2-yl, thiazol-5-yl, 2-(hydroxymethyl)thiazol-4-yl, 2-(acetoxymethyl)thiazol-4-yl, 2-(fluoromethyl)thiazol-4-yl, 2-vinylthiazol-4-yl, 2-ethylthiazol-4-yl, 2-furyl, 2-thienyl, Ph, 3-pyridyl, CH₂C(OEt)Me-(Z), R₂ = H) and III (R₃ = 2-(5-acetoxypentyl)thiazol-4-yl, 2-(methylthio)thiazol-4-yl, 2-piperidinethiazol-4-yl, 2-methoxythiazol-4-yl, 2-ethoxythiazol-4-yl, thiazol-4-yl, thiazol-2-yl, thiazol-5-yl, 2-(hydroxymethyl)thiazol-4-yl, 2-(acetoxymethyl)thiazol-4-yl, 2-(fluoromethyl)thiazol-4-yl, 2-vinylthiazol-4-yl, 2-ethylthiazol-4-yl, 2-furyl, 2-thienyl, Ph, 3-pyridyl, CH₂C(OEt)Me-(Z)). The Stille coupling approach was then used to prepare epothilone B analogs from the key macroketone intermediate I (R₁ = I, R₂ = CH₂OH) which was itself synthesized by a macrolactonization based strategy.


RX(19) OF 264

...BT ==> AH...



BT

(19)



AH
YIELD 84%


RX(19) RCT BT 240816-02-8

STAGE(1)

RGT CB 4136-95-2 2,4,6-Cl₃C₆H₂COCl, BW 121-44-8 Et₃N
SOL 109-99-9 THF

116

PAGE 1-B



AY

(12)

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

RX(12) RCT AY 298702-20-2

STAGE(1)

RCT N 121-44-8 EC3N, BB 50-43-1 Benzoic acid, 2,4,6-trichloro-
SOL 109-99-9 THE

STAGE(2)

RCT BC 1122-58-3 4-DMAP
SOL 108-88-3 PMe

PRO BA 298702-21-3

NTE STEREOSELECTIVE

REFERENCE COUNT: 51

THERE ARE 51 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L109 ANSWER 5 OF 7 CASREACT COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 132:49832 CASREACT Full-text

TITLE: Preparation of 16-desmethyllepothilones for the
treatment of proliferative diseases.

INVENTOR(S): Nicolaou, Kyriacos Costa; Hepworth, David; Finlay,
Maurice Raymond Varschoyle; King, Nigel Paul
PATENT ASSIGNEE(S): Novartis A.-G., Switz.; Novartis-Erfindungen
Verwaltungsgesellschaft m.b.H.; Scripps Research
Institute

SOURCE: PCT Int. Appl., 31 pp.

CODEN: PIXAD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 5

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9967253	A2	19991229	WO 1999-EP4299	19990621
WO 9967253	A3	20000420		

W: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ,

113

DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HU, ID, IL, IN, IS,
JP, KE, KG, KR, KZ, LC, LR, LS, LT, LU, LV, MD, MG, MK,
MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, ST,
TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW
RW: GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW, AT, BE, CH, CY, DE, DK,
ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG,
CI, CM, GN, GW, ML, MR, NE, SN, TD, TG

US 6380394 B1 20020430 US 1998-102602 19980622
AU 9947752 A 20000110 AU 1999-47752 19990621
US 1998-102602 19980622
US 1999-123155P 19990306
US 1999-124633P 19990316
US 1996-32864P 19961213
US 1997-856533 19970514
US 1997-923869 19970904
WO 1999-EP4299 19990621

OTHER SOURCE(S):

MARPAT 132:49832


GI

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB The invention relates to compds. I [X = bond, O; Q = OH, I, H], and methods of
synthesis of I, as well as for the synthesis of epothilone B (II) and their
intermediates. Thus, 16-desmethylepothilone analog III was prepared via
Yamaguchi macrolactonization of hydroxy acid IV. The compds. I can be used
e.g. in the treatment of proliferative diseases.

RX(4) OF 46

...J ==> W...



J

(4)

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

RX(4) RCT J 252986-91-7

114

NTE STEREOSELECTIVE

REFERENCE COUNT: 22

THERE ARE 22 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L109 ANSWER 3 OF 7 CASREACT COPYRIGHT 2007 ACS ON STN

ACCESSION NUMBER: 134:4795 CASREACT Full-text

TITLE: Total Syntheses of Epothilones B and D

AUTHOR(S): Mulzer, Johann; Mantoulidis, Andreas; Oehler,

Elisabeth

CORPORATE SOURCE: Institut fuer Organische Chemie, Universitaet Wien,

Vienna, A-1090, Austria

SOURCE: Journal of Organic Chemistry (2000), 65 (22), 7456-7467

CODEN: JOCEAH; ISSN: 0022-3263


PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Total syntheses of the microtubule stabilizing anticancer drugs epothilone B and D are described, starting from optically pure (S)-malic acid and Me (R)-3-hydroxy-2-methylpropanoate. The synthesis is highly convergent by coupling the three fragments C1-C6 (fragment D), C7-C10 (fragment C), and C11-C21 (fragment B). Key steps are two stereoselective Wittig type olefinations to generate the 12,13- and 16,17-double bonds, an enantioselective Mukaiyama aldehyde addition to synthesize fragment D, and a sulfone anion allyl iodide alkylation to connect fragments B and C. Finally fragment D was attached to the B + C fragment via aldol addition

RX (34) OF 711 ...DL ==> H...



DL

(34)

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

RX (34) RCT DL 193146-26-8

STAGE (1)

RGT RN 1122-58-3 4-DMAP, DO 25952-53-8 EDAP, DP 71561-71-2

111

4-Me2NC5H4N.HCl

SOL 67-66-3 CHCl3

STAGE (2)

RGT AK 12125-02-9 NH4Cl

SOL 7732-18-5 Water

PRO H 189453-35-8

REFERENCE COUNT: 41

THERE ARE 41 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE-FORMAT

L109 ANSWER 4 OF 7 CASREACT COPYRIGHT 2007 ACS ON STN

ACCESSION NUMBER: 133:26634 CASREACT Full-text

TITLE: Total Synthesis and Antitumor Activity of

12,13-Desoxyepothilone F: An Unexpected Solvolysis

Problem at C15, Mediated by Remote Substitution at C21

AUTHOR(S): Lee, Chul Bom; Chou, Ting-Chao; Zhang, Xiu-Guo; Wang,

Zhi-Guang; Kuduk, Scott D.; Chappell, Mark D.;

Stachel, Shawn J.; Danishefsky, Samuel J.

CORPORATE SOURCE: Laboratory for Bioorganic Chemistry, The

Sloan-Kettering Institute for Cancer Research, New

York, NY, 10021, USA

SOURCE: Journal of Organic Chemistry (2000), 65 (20), 6325-6333

CODEN: JOCEAH; ISSN: 0022-3263


PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal

LANGUAGE: English

AB A new epothilone analog, 12,13-desoxyepothilone F (dEpof, 21-hydroxy-12,13-desoxyepothilone B, 21-hydroxyepothilone D), was synthesized and evaluated for antitumor potential. A convergent strategy employed for the semi-practical synthesis of 12,13-desoxyepothilone B (dEpof) has been utilized to yield an amount of dEpof sufficient for relevant biol. studies. The results from an in vitro assay reveal that this new analog is highly active against various tumor cell lines with a potency comparable to that of dEpof. In particular, the growth of resistant tumor cells is inhibited by dEpof at concns. where paclitaxel (Taxol) is basically ineffective. A preliminary assessment of its in vivo activity is also promising. The new analog, containing an addnl. hydroxyl group at C21, exhibits advantages over other epothilones in terms of water solubility, and can serve as a readily functionalizable handle to produce other useful compds. for pertinent biol. studies.

RX (12) OF 128 ...AY ==> BA...



112

STAGE(2)

SOL 108-88-3 PhMe

STAGE(3)

RGT AV 1122-58-3 4-DMAP

SOL 108-88-3 PhMe

PRO W 279226-97-0

NTE stereoselective

REFERENCE COUNT: 25

THERE ARE 25 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L109 ANSWER 2 OF 7 CASREACT COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER:

134:29228 CASREACT Full-text

TITLE: A novel highly stereoselective total synthesis of

epothilone B and of its (12R,13R) acetone

Mulzer, J.; Karig, G.; Pöjarliev, P.

Institut für Organische Chemie der Universität Wien,

Vienna, A-1090, Austria

Tetrahedron Letters (2000), 41(40), 7635-7638


CODEN: TELEAY; ISSN: 0040-4039

Elsevier Science Ltd.

Journal

English

GI




II

AB Stereoselective syntheses of epothilone B and its novel derivative I are described. Key steps are the formation of intermediate II via Sharpless AD-reaction and Davis-Evans-hydroxylation.

RX(30) OF 447


...CI ==> CM...

109



CI

(30)



YIELD 50%

RX(30) RCT CI 312492-96-9

STAGE(1)

RGT CN 429-41-4 Bu4N.F

SOL 109-99-9 THF

STAGE(2)

RGT CO 4136-95-2 2,4,6-Cl3C6H2COCl, AF 121-44-8 Et3N

SOL 108-88-3 PhMe

STAGE(3)

RGT CP 1122-58-3 4-DMAP

SOL 108-88-3 PhMe


PRO CM 312492-96-1

110

L90 28 SEA FILE-CASREACT ABB-ON PLU-ON L85 NOT L88
 L106 21 SEA FILE-CASREACT ABB-ON PLU-ON ("126:251010"/AN OR "127:1087
 93"/AN OR "127:293040"/AN OR "128:101936"/AN OR "129:189151"/AN OR
 OR "131:199535"/AN OR "131:286299"/AN OR "131:31819"/AN OR
 "131:31829"/AN OR "131:351125"/AN OR "132:251011"/AN OR
 "132:49832"/AN OR "133:266631"/AN OR "133:266634"/AN OR
 "134:321737"/AN OR "133:362657"/AN OR "134:178371"/AN OR
 "134:292228"/AN OR "134:4795"/AN OR "134:56502"/AN OR "135:37156
 6"/AN OR "1997:206419"/AN OR "1997:430309"/AN OR "1997:665094"/
 AN OR "1997:787450"/AN OR "1998:378435"/AN OR "1999:176999"/AN
 OR "1999:372044"/AN OR "1999:383492"/AN OR "1999:444724"/AN OR
 "1999:606636"/AN OR "1999:819379"/AN OR "2000:514132"/AN OR
 "2000:52387"/AN OR "2000:597944"/AN OR "2000:624043"/AN OR
 "2000:701228"/AN OR "2000:719579"/AN OR "2000:733774"/AN OR
 "2000:842116"/AN OR "2000:853645"/AN OR "2001:843887"/AN)
 L108 21 SEA FILE-CASREACT ABB-ON PLU-ON L106 AND L43
 L109 7 SEA FILE-CASREACT ABB-ON PLU-ON L108 NOT L90


=> d ibib abs fhlt L109 1-7; d ibib abs fhlt L90 1-28

L109 ANSWER 1 OF 7 CASREACT COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 134:178371 CASREACT Full-text
 TITLE: Synthesis and biological evaluation of highly potent
 analogues of epothilones B and D
 AUTHOR(S): Altmann, K.-H.; Bold, G.; Caravatti, G.; Florzheimer, A.;
 Guagnano, V.; Wartmann, M.
 CORPORATE SOURCE: Novartis Pharma AG, TA Oncology Research, Basel, CH-4002, Switz.
 SOURCE: Bioorganic & Medicinal Chemistry Letters (2000), 10(24), 2765-2768
 CODEN: BMCL68; ISSN: 0960-894X
 PUBLISHER: Elsevier Science Ltd.
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 GI




AB A series of new epothilone B and D analogs incorporating fused hetero-aromatic side chains have been prepared. The synthetic strategy is based on olefin I as the common intermediate and allows variation of the side-chain structure in a highly convergent and stereoselective manner. These epothilone analogs, e.g. II, are more potent inhibitors of cancer cell proliferation than the corresponding parent epothilones B or D.

RX(11) OF 370 ...AR ==> W...



(11) →



YIELD 70%

RX(11) RCT AR 279226-96-9

STAGE(1)
 RGT . AT 4136-95-2 2,4,6-Cl3C6H2COCl, AU 121-44-8 Et3N
 SOL 109-99-9 THF

chain nodes :
 12 13 14 15 16 17 18 19 20 21 22
 ring/chain nodes :
 1 2 3 4 5 6 7 8 9 10 11
 chain bonds :
 2-12 2-13 3-14 3-15 4-16 4-17 6-18 8-19 10-20 11-21 11-22
 ring/chain bonds :
 1-2 2-3 3-4 4-5 5-6 6-7 7-8 8-9 9-10 10-11
 exact/norm bonds :
 1-2 2-3 3-4 4-5 5-6 6-7 6-18 7-8 8-9 8-19 9-10 10-11
 exact bonds :
 2-12 2-13 3-14 3-15 4-16 4-17 10-20 11-21 11-22

Match level :
 1:CLASS 2:CLASS 3:CLASS 4:CLASS 5:CLASS 6:CLASS 7:CLASS 8:CLASS 9:CLASS
 10:CLASS 11:CLASS 12:CLASS 13:CLASS 14:CLASS 15:CLASS 16:CLASS 17:CLASS
 18:CLASS 19:CLASS
 20:CLASS 21:CLASS 22:CLASS

L3 560 SEA FILE-REGISTRY SSS FUL L1
 L34 22933 SEA FILE-REGISTRY ABB-ON PLU-ON OC15/ESS
 L35 27330 SEA FILE-REGISTRY ABB-ON PLU-ON C16/ESS
 L36 726 SEA FILE-REGISTRY ABB-ON PLU-ON NC15/ESS
 L37 0 SEA FILE-REGISTRY ABB-ON PLU-ON NSC14/ESS
 L38 50989 SEA FILE-REGISTRY ABB-ON PLU-ON (L34 OR L35 OR L36 OR L37)
 L39 12165 SEA FILE-REGISTRY ABB-ON PLU-ON L38 AND CASREACT/LC
 L40 2534 SEA FILE-CASREACT ABB-ON PLU-ON L39/PRO
 L42 65 SEA FILE-CASREACT ABB-ON PLU-ON L3/RTT
 L43 59 SEA FILE-CASREACT ABB-ON PLU-ON L42 (L) L40
 L85 59 SEA FILE-CASREACT ABB-ON PLU-ON L43 (L) 1/NS
 L88 31 SEA FILE-CASREACT ABB-ON PLU-ON L43 (L) 4/NS
 L90 28 SEA FILE-CASREACT ABB-ON PLU-ON L85 NOT L88


=>

=> d stat que L109

L1 STR

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

Structure attributes must be viewed using STN Express query preparation:
 Uploading L1.str



chain nodes :
 12 13 14 15 16 17 18 19 20 21 22
 ring/chain nodes :
 1 2 3 4 5 6 7 8 9 10 11
 chain bonds :
 2-12 2-13 3-14 3-15 4-16 4-17 6-18 8-19 10-20 11-21 11-22
 ring/chain bonds :
 1-2 2-3 3-4 4-5 5-6 6-7 7-8 8-9 9-10 10-11
 exact/norm bonds :
 1-2 2-3 3-4 4-5 5-6 6-7 6-18 7-8 8-9 8-19 9-10 10-11
 exact bonds :
 2-12 2-13 3-14 3-15 4-16 4-17 10-20 11-21 11-22

Match level :

1:CLASS 2:CLASS 3:CLASS 4:CLASS 5:CLASS 6:CLASS 7:CLASS 8:CLASS 9:CLASS
 10:CLASS 11:CLASS 12:CLASS 13:CLASS 14:CLASS 15:CLASS 16:CLASS 17:CLASS
 18:CLASS 19:CLASS
 20:CLASS 21:CLASS 22:CLASS

L3 560 SEA FILE-REGISTRY SSS FUL L1
 L34 22933 SEA FILE-REGISTRY ABB-ON PLU-ON OC15/ESS
 L35 27330 SEA FILE-REGISTRY ABB-ON PLU-ON C16/ESS
 L36 726 SEA FILE-REGISTRY ABB-ON PLU-ON NC15/ESS
 L37 0 SEA FILE-REGISTRY ABB-ON PLU-ON NSC14/ESS
 L38 50989 SEA FILE-REGISTRY ABB-ON PLU-ON (L34 OR L35 OR L36 OR L37)
 L39 12165 SEA FILE-REGISTRY ABB-ON PLU-ON L38 AND CASREACT/LC
 L40 2534 SEA FILE-CASREACT ABB-ON PLU-ON L39/PRO
 L42 65 SEA FILE-CASREACT ABB-ON PLU-ON L3/RTT
 L43 59 SEA FILE-CASREACT ABB-ON PLU-ON L42 (L) L40
 L85 59 SEA FILE-CASREACT ABB-ON PLU-ON L43 (L) 1/NS
 L88 31 SEA FILE-CASREACT ABB-ON PLU-ON L43 (L) 4/NS

STAGE(1)
 RCT AW 1191-15-7 ALH(BU-i)2
 SOL 108-88-3 PhMe

STAGE(2)
 RCT BK 7647-01-0 HCl
 SOL 7732-18-5 Water, 67-56-1 MeOH

PRO BJ 226940-68-7
 RX(16) RCT BL 187527-25-9

STAGE(1)
 RCT BE 108-18-9 i-Pr2NH, BF 109-72-8 BuLi
 SOL 109-99-9 THF, 110-54-3 Hexane

STAGE(2)
 RCT BJ 226940-68-7
 SOL 109-99-9 THF

STAGE(3)
 RCT BO 64-19-7 AcOH
 SOL 109-99-9 THF

PRO BM 240815-96-7, BN 240815-97-8

REFERENCE COUNT: 60 THERE ARE 60 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> file casreact
 FILE 'CASREACT' ENTERED AT 12:18:14 ON 11 OCT 2007
 USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT
 COPYRIGHT (C) 2007 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications.

FILE CONTENT:1840 - 6 Oct 2007 VOL 147 ISS 16

New CAS Information Use Policies, enter HELP USAGETERMS for details.

 * CASREACT now has more than 13.8 million reactions *


Some CASREACT records are derived from the ZIC/VINITI database (1974-1999) provided by InfoChem, INPI data prior to 1986, and BioTransformations database compiled under the direction of Professor Dr. Klaus Kieselich.


This file contains CAS Registry Numbers for easy and accurate substance identification.

=> d stat que L90
 LI STR



* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

Structure attributes must be viewed using STN Express query preparation:
 Uploading Li.str





2 BL

7
STERSBN
YIELD 39%BN
YIELD 27%

- RX(10) RCT AK 226940-62-1, AR 95561-65-2
PRO AS 240815-89-8
SOL 71-43-2 Benzene
- RX(11) RCT AS 240815-89-8
STAGE(1)
RGT AW 1191-15-7 AlH(Bu-i)2
SOL 109-99-9 THF, 75-09-2 CH2Cl2
STAGE(2)
RCT AU 76-83-5
RGT Z 1122-58-3 4-DMAP
SOL 68-12-2 DME
- PRO AV 240815-91-2
RX(12) RCT AV 240815-91-2
STAGE(1)
RGT AY 280-64-8 9-BBN
SOL 109-99-9 THF
STAGE(2)
RGT AZ 1310-73-2 NaOH, D 7722-84-1 H2O2;
SOL 7732-18-5 Water
STAGE(3)
RGT AL 288-32-4 1H-Imidazole, BA 603-35-0 PPh3
SOL 60-29-7 Et2O, 75-05-8 MeCN
STAGE(4)
RGT BB 7553-56-2 I2
PRO AX 240815-93-4
NTE (95%;97%)
- RX(13) RCT BC 70113-32-5
STAGE(1)
RGT BE 108-18-9 i-Pr2NH, BF 109-72-8 BuLi
SOL 109-99-9 THF, 110-54-3 Hexane
STAGE(2)
RCT AX 240815-93-4
SOL 109-99-9 THF
PRO BD 240815-94-5
NTE stereoselective
- RX(14) RCT BD 240815-94-5
RGT BI 109536-69-8 2-HO2CC6H4CO3H.Mg
PRO BH 240815-95-6
SOL 7732-18-5 Water, 67-56-1 MeOH, 109-99-9 THF
NTE phosphate buffer (7.0)
- RX(15) RCT BH 240815-95-6

CAT 3144-16-9 10-CSA
SOL 67-56-1 MeOH, 75-09-2 CH₂Cl₂

STAGE(2)

RET H 87413-09-0 Martin's reagent
SOL 75-09-2 CH₂Cl₂

PRO AN 193146-27-9
NTE (82%, 97%)

RX(22) RCT V 185148-95-2

STAGE(1)

RCT X 109-72-8 BuLi, Y 108-18-9 i-Pr₂NH
SOL 109-99-9 THF, 110-54-3 Hexane

STAGE(2)

RCT AN 193146-27-9
SOL 109-99-9 THF

PRO BY 210690-87-2, BZ 250284-01-6

NTE stereoselective key step

REFERENCE COUNT: 82 THERE ARE 82 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L33 ANSWER 23 OF 23 CASREACT COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 131:199535 CASREACT Full-text

TITLE: Total synthesis of epothilone E and related side-chain modified analogues via a Stille coupling based strategy

AUTHOR(S): Nicolaou, K. C.; King, N. P.; Finlay, M. R. V.; He, Y.; Roschangar, F.; Vourloumis, D.; Vallberg, H.; Sarabia, F.; Ninkovic, S.; Hepworth, D.

CORPORATE SOURCE: Department of Chemistry and The Skaggs Institute for Chemical Biology, The Scripps Research Institute, La Jolla, CA, 92037, USA


SOURCE: Bioorganic & Medicinal Chemistry (1999), 7(5), 665-697
CODEN: BMCEP; ISSN: 0968-0896

PUBLISHER: Elsevier Science Ltd.

DOCUMENT TYPE: Journal

LANGUAGE: English

GI








AB A Stille coupling strategy has been utilized to complete a total synthesis of epothilone E from vinyl iodide I (R₁ = I; R₂ = H) and thiazolostannane II. The central core fragment I (R₁ = I; R₂ = H) and its trans-isomer III (R₃ = I) were prepared from triene IV (TBS = SiMe₂CMes) using ring-closing metathesis (RCM), and were subsequently coupled to a variety of alternative stannanes to provide a library of epothilone analogs I (R₁ = 2-(5-acetoxypentyl)thiazol-4-yl, 2-(methylthio)thiazol-4-yl, 2-piperidinethiazol-4-yl, 2-methoxythiazol-4-yl, 2-ethoxythiazol-4-yl, thiazol-4-yl, thiazol-2-yl, thiazol-5-yl, 2-(hydroxymethyl)thiazol-4-yl, 2-(acetoxymethyl)thiazol-4-yl, 2-(fluoromethyl)thiazol-4-yl, 2-vinylthiazol-4-yl, 2-ethylthiazol-4-yl, 2-furyl, 2-thienyl, Ph, 3-pyridyl, CH₃C(OMe)Me (2), R₂ = H) and III (R₃ = 2-(5-acetoxypentyl)thiazol-4-yl, 2-(methylthio)thiazol-4-yl, 2-piperidinethiazol-4-yl, 2-methoxythiazol-4-yl, 2-ethoxythiazol-4-yl, thiazol-4-yl, thiazol-2-yl, 2-thiazol-5-yl, 2-(hydroxymethyl)thiazol-4-yl, 2-(acetoxymethyl)thiazol-4-yl, 2-(fluoromethyl)thiazol-4-yl, 2-vinylthiazol-4-yl, 2-ethylthiazol-4-yl, 2-furyl, 2-thienyl, Ph, 3-pyridyl, CH₃C(OMe)Me (2)). The Stille coupling approach was then used to prepare epothilone B analogs from the key macrolactone intermediate I (R₁ = I, R₂ = CH₂OH) which was itself synthesized by a macrolactonization based strategy.

RX(132) OF 264 COMPOSED OF RX(10), RX(11), RX(12), RX(13), RX(14), RX(15),

RX(16)

RX(132) 2 AK + 2 AR + 2 AU + 2 BC + 2 BL ==> EM + BN





RX(12) RCT AT 4736-60-1

STAGE(1)
RGT X 109-72-8 BuLi
SOL 109-99-9 THF, 110-54-3 Hexane

STAGE(2)
RGT AW 7553-56-2 I2
SOL 109-99-9 THF

STAGE(3)
RGT U 1070-89-9 (Me3Si)2N.Na
SOL 109-99-9 THF

STAGE(4)
RCT AU 188730-08-7
SOL 109-99-9 THF

PRO AV 210690-66-7

RX(21) RCT AP 113453-27-3

STAGE(1)
RGT BS 12621-78-2 Zinc alloy, base, Zn,Cu
CAT 106-93-4 BrCH2CH2Br
SOL 71-43-2 Benzene

STAGE(2)
RGT Q 75-77-4 Me3SiCl, BT 127-19-5 AcNMe2

STAGE(3)
RGT BU 79271-56-0 F3CSO3SiEt3, BT 127-19-5 AcNMe2
CAT 14221-01-3 Pd(PPh3)4

STAGE(4)
RCT AV 210690-66-7
SOL 71-43-2 Benzene

PRO AM 210690-85-0
NTE key step

RX(10) RCT AM 210690-85-0

STAGE(1)

STAGE(5)
 SOL 109-66-0 Pentane
 PRO B 298702-07-5
 NTE STEREOSELECTIVE
 RX(1) RCT A 224580-52-3

STAGE(1)
 RGT D 280-64-8 9-BEN
 SOL 109-99-9 THF

STAGE(2)
 RCT B 298702-07-5
 RGT E 534-17-8 Cs2CO3, F 603-32-7 Ph3As
 CAT 72287-26-4 Palladium, [1,1'-bis(diphenylphosphino-
 xP)ferrocene]dichloro-, (SP-4-2)-
 SOL 68-12-2 DMF

STAGE(3)
 SOL 7732-18-5 Water
 PRO C 298702-16-6
 NTE STEREOSELECTIVE
 RX(9) RCT C 298702-16-6

STAGE(1)
 RGT AR 7647-01-0 HCl
 SOL 67-56-1 MeOH

STAGE(2)
 RGT O 144-55-8 NaHCO3
 SOL 7732-18-5 Water

STAGE(3)
 SOL 75-09-2 CH2Cl2
 PRO AQ 298702-17-7
 NTE STEREOSELECTIVE
 RX(10) RCT AQ 298702-17-7, AS 67-56-1

STAGE(1)
 RGT AV 1333-74-0 H2
 CAT 109361-17-3 Ruthenium, bis[(1R)-[1,1'-binaphthalene]-2,2'-
 diylbis(diphenylphosphine-xP)]di- μ -
 chlorodichloro(N,N-diethylethanamine)di-
 SOL 67-56-1 MeOH


STAGE(2)
 RGT O 144-55-8 NaHCO3

STAGE(3)
 SOL 75-09-2 CH2Cl2

PRO AT 298702-18-8, AU 298702-19-9
 THERE ARE 51 CITED REFERENCES AVAILABLE FOR THIS
 RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

95

L33 ANSWER 22 OF 23 CASREACT COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 131:351125 CASREACT Full-text
 TITLE: Syntheses of (-)-epothilone B
 AUTHOR(S): Schinzer, Dieter; Bauer, Armin; Schieber, Jennifer
 CORPORATE SOURCE: Chemisches Institut der Otto-von-Guericke-Universität,
 Magdeburg, D-39106, Germany
 SOURCE: Chemistry--A European Journal (1999), 5(9), 2492-2500
 CODEN: CEUJED; ISSN: 0947-6539
 PUBLISHER: Wiley-VCH Verlag GmbH
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 GI







AB Two efficient routes for the total synthesis of (-)-epothilone B (I) are reported. One strategy is based on ring-closing metathesis, and a second synthesis on a macrolactonization. The key fragments are available on large scale to provide sufficient material for biol. tests. Thiazole fragment II (TEDMS = SiMe2OMe3) was obtained by an improved route starting from (S)-malic acid. The first synthesis is based on our preceding paper. The critical trisubstituted double bond C12-13 in our second approach was constructed by a highly efficient Pd-mediated coupling reaction. Ring closure was achieved by macrolactonization.

RX(79) OF 215 COMPOSED OF RX(12), RX(21), RX(10), RX(22)
 RX(79) 2 AT + 2 AU + 2 AP + 2 V ==> BY + BZ



96

water solubility, and can serve as a readily functionalizable handle to produce other useful compds. for pertinent biol. studies.


RX(33) OF 128 COMPOSED OF RX(7), RX(1), RX(9), RX(10)
RX(33) 2 AG + 2 AH + AS ==> AT + AU



PAGE 1-B



PAGE 1-B



RX(7) RCT AG 82294-38-0

STAGE(1)
RCT AI 109-72-8 BuLi
SOL 109-99-9 THF

STAGE(2)
RCT AJ 7553-56-2 I2
SOL 109-99-9 THF

STAGE(3)
RCT AK 1070-89-9 (Me3Si)2N.Na

STAGE(4)
RCT AH 298702-12-2
SOL 109-99-9 THF

STAGE (4)

RCT EP 107905-52-2
 RGT ES 17455-13-9 18-Crown-6, BJ 40949-94-8 K (N(SiMe₃)₂)
 SOL 109-99-9 THF

STAGE (5)

RGT AK 12125-02-9 NH₄Cl
 SOL 7732-18-5 Water, 60-29-7 Et₂O

PRO EU 218614-04-1
 NTE stereoselective

RCT EU 218614-04-1

RX (61)

STAGE (1)

RGT AN 1191-15-7 AlH(Bu-1)₂
 SOL 109-99-9 THF, 142-82-5 Heptane

STAGE (2)

RGT AO 304-59-6 Rochelle salt
 SOL 67-56-1 MeOH, 60-29-7 Et₂O

PRO EX 218614-16-5

RCT EX 218614-16-5

RX (55)

STAGE (1)

RGT EF 603-35-0 PPh₃
 SOL 75-05-8 MeCN, 60-29-7 Et₂O

STAGE (2)

RGT CB 288-32-4 1H-Imidazole

STAGE (3)

RGT EY 7553-56-2 I₂

STAGE (4)

RCT ED 218613-98-0
 RGT ES 17455-13-9 18-Crown-6, BJ 40949-94-8 K (N(SiMe₃)₂)
 SOL 109-99-9 THF

STAGE (5)

SOL 109-99-9 THF

STAGE (6)

RGT AK 12125-02-9 NH₄Cl
 SOL 7732-18-5 Water

PRO CV 308357-81-5
 NTE diastereomeric mikt.

RCT CV 308357-81-5

RGT CX 7558-79-4 Na₂HPO₄, CY 11110-52-4 Sodium amalgam
 PRO CW 210690-85-0
 SOL 67-56-1 MeOH, 109-99-9 THF

RCT CW 210690-85-0

RX (29)

STAGE (1)

RGT DA 3144-16-9 10-CSA

SOL 67-56-1 MeOH, 75-09-2 CH₂Cl₂

STAGE (2)

RGT D 144-55-8 NaHCO₃
 SOL 7732-18-5 Water

PRO CZ 210690-99-6

RCT CZ 210690-99-6

RGT BT 77-76-9 Me₂C(OMe)₂

PRO FA 193146-27-9

SOL 73-09-2 CH₂Cl₂, 110-86-1 Pyridine

RCT CU 187527-25-9

RX (57)

STAGE (1)

RGT W 4111-54-0 LiN(Pi-1)₂
 SOL 109-99-9 THF

STAGE (2)

RCT FA 193146-27-9
 SOL 109-99-9 THF

STAGE (3)

RGT AK 12125-02-9 NH₄Cl
 SOL 7732-18-5 Water

PRO FB 193146-50-8, DB 193146-49-5

REFERENCE COUNT: 41

THERE ARE 41 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L33 ANSWER 21 OF 23 CASREACT COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 133:266634 CASREACT Full-text

TITLE: Total Synthesis and Antitumor Activity of

12,13-Desoxyepothilone F: An Unexpected Solvolysis

Problem at C13, Mediated by Remote Substitution at C21

Lee, Chul Bom; Chou, Ting-Chao; Zhang, Xiu-Guo; Wang,

Zhi-Guang; Kuduk, Scott D.; Chappell, Mark D.;

Stachel, Shawn J.; Danishefsky, Samuel J.

Laboratory for Bioorganic Chemistry, The

Sloan-Kettering Institute for Cancer Research, New

York, NY, 10021, USA

SOURCE: Journal of Organic Chemistry (2000), 65(20), 6525-6533


CODEN: JOCEAH; ISSN: 0022-3263

PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal


LANGUAGE: English

AB A new epothilone analog, 12,13-desoxyepothilone F (dEpOF, 21-hydroxy-12,13-desoxyepothilone B, 21-hydroxyepothilone D), was synthesized and evaluated for antitumor potential. A convergent strategy employed for the semi-practical synthesis of 12,13-desoxyepothilone B (dEpOB) has been utilized to yield an amount of dEpOF sufficient for relevant biol. studies. The results from an in vitro assay reveal that this new analog is highly active against various tumor cell lines with a potency comparable to that of dEpOB. In particular, the growth of resistant tumor cells is inhibited by dEpOF at concns. where paclitaxel (Taxol) is basically ineffective. A preliminary assessment of its in vivo activity is also promising. The new analog, containing an addnl. hydroxyl group at C21, exhibits advantages over other epothilones in terms of




2 ET

2 EP




2 ED




2 CU

7
STEPS

PAGE 1-A




PAGE 1-B




FB
YIELD 69% (80)

PAGE 1-A



PAGE 1-B





DB
YIELD 69% (80)

RX(52) RCT ET 18899-14-1

STAGE(1)
RGT R 79-37-8 (COCl)₂, S 67-68-5 DMSO
SOL 75-09-2 CH₂Cl₂

STAGE(2)
RGT T 7087-68-5 EtN(Pr-i)₂

STAGE(3)
RGT AK 12125-02-9 NH₄Cl
SOL 60-29-7 Et₂O, 7732-18-5 Water



PAGE 1-B

-OPH.

CJ
YIELD 50%

RX(15) RCT BG 188730-08-7, BH 4736-60-1

STAGE(1)

RGT BJ 109-72-8 BuLi
SOL 109-99-9 THF

STAGE(2)

RGT AZ 7553-56-2 I2

STAGE(3)

RGT BK 1070-89-9 (Me3Si)2N.Na

PRO BI 210690-66-7
NTE STEREOSELECTIVE

RX(16) RCT BI 210690-66-7
RGT BA 7664-39-3 HF, BB 110-86-1 Pyridine
PRO BL 312730-71-5
SOL 109-99-9 THF
NTE STEREOSELECTIVE

RX(17) RCT BL 312730-71-5, BC 108-24-7
RGT BE 121-44-8 Et3N, BF 1122-58-3 4-DMAP
PRO BM 189453-18-7
SOL 75-09-2 CH2Cl2
NTE STEREOSELECTIVE

RX(26) RCT BV 262375-53-1, BM 189453-18-7

STAGE(1)

RGT CM 280-64-8 9-BBN
SOL 109-99-9 THF

STAGE(2)

RGT CN 7778-53-2 K3PO4
CAT 72287-26-4 Palladium, (1,1'-bis(diphenylphosphino)-
kP)ferrocene]dichloro-, (SP-4-2)-
SOL 68-12-2 DMF, 7732-18-5 Water

PRO CJ 312730-85-1

NTE ULTRASOUND IN FIRST STAGE, STEREOSELECTIVE

REFERENCE COUNT: 42 THERE ARE 42 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L33 ANSWER 20 OF 23 CASREACT COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 134:4795 CASREACT Full-text

TITLE: Total Syntheses of Epothilones B and D

AUTHOR(S): Mulzer, Johann; Mantoulidis, Andreas; Oehler,

Elisabeth

CORPORATE SOURCE: Institut fuer Organische Chemie, Universitaet Wien,

Vienna, A-1090, Austria

SOURCE: Journal of Organic Chemistry (2000), 65(22), 7456-7467

CODEN: JOCEAH; ISSN: 0022-3263

PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Total syntheses of the microtubule stabilizing antitumor drugs epothilone B and D are described, starting from optically pure (S)-malic acid and Me (R)-3-hydroxy-2-methylpropanoate. The synthesis is highly convergent by coupling the three fragments C1-C6 (fragment D), C7-C10 (fragment C), and C11-C21 (fragment B). Key steps are two stereoselective Wittig type olefinations to generate the 12,13- and 16,17-double bonds, an enantioselective Mukaiyama aldol addition to synthesize fragment D, and a sulfone anion allyl iodide alkylation to connect fragments B and C. Finally fragment D was attached to the B + C fragment via aldol addition

RX(411) OF 711 COMPOSED OF RX(52), RX(61), RX(55), RX(28), RX(29), RX(56),

RX(57)

RX(411) 2 ET + 2 EP + 2 ED + 2 CU ==> FB + DB

STAGE(1)

RGT AW 4111-54-0 LIn(Pr-1)2
SOL 109-99-9 THF

STAGE(2)

RGT AS 312492-69-6
SOL 109-99-9 THF

STAGE(3)

RGT J 12125-02-9 NH4Cl
SOL 7732-18-5 Water

PRO AV 321522-36-5

RX(13) RCT AV 321522-36-5, AX 17341-93-4

STAGE(1)

RGT T 110-86-1 Pyridine
SOL 75-09-2 CH2Cl2

STAGE(2)

RGT Y 144-55-8 NaHCO3
SOL 7732-18-5 Water

PRO AV 321522-37-6

RX(14)

STAGE(1)

RGT BA 20816-12-0 OsO4, BB 7529-22-8 Me-morpholineoxide
SOL 109-99-9 THF, 75-65-0 t-BuOH, 7732-18-5 Water

STAGE(2)

RGT V 7772-98-7 Na2S2O3
SOL 7732-18-5 Water, 75-09-2 CH2Cl2

STAGE(3)

RGT C 10028-15-6 Ozone
SOL 64-17-5 EtOH, 7732-18-5 Water

STAGE(4)

RGT Y 144-55-8 NaHCO3
SOL 7732-18-5 Water

PRO AZ 321522-38-7

L33 ANSWER 19 OF 23 CASREACT COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER:

134:56302 CASREACT Full-text
Enantioselective Total Synthesis of Epothilones A and B Using Multifunctional Asymmetric Catalysis

AUTHOR(S):

Sawada, Daisuke; Kanai, Motomu; Shibasaki, Masakatsu
Graduate School of Pharmaceutical Sciences, The University of Tokyo, Bunkyo-ku Tokyo, 113-0033, Japan

SOURCE:

Journal of the American Chemical Society (2000), 122(43), 10521-10532

CODEN: JACSAT; ISSN: 0002-7863

PUBLISHER:

American Chemical Society


DOCUMENT TYPE:

85

LANGUAGE:


English

GI







AB An enantioselective total synthesis of epothilones A and B using multifunctional asym. catalysis such as a cyanosilylation of an aldehyde, an aldol reaction of an unmodified ketone with an aldehyde, and a protonation in the conjugate addition of a thiol to an α,β -unsatd. thioester has been achieved. Epothilones A and B were divided into fragment A (I), fragment B (II), and fragment C (III). A catalytic asym. synthesis of fragments A and B was accomplished using a catalytic asym. cyanosilylation as a key step. An enantiocontrolled synthesis of fragment C was achieved in two ways. One is the use of a direct catalytic asym. aldol reaction of an unmodified ketone with an aldehyde as a key step, and the other utilizes a catalytic asym. protonation in the conjugate addition of a thiol to an α,β -unsatd. thioester as a key step. Suzuki cross-coupling of fragment A with fragment C followed by Yamaguchi lactonization as key steps led to an enantiocontrolled synthesis of epothilone A. On the other hand, Suzuki cross-coupling of fragment B with fragment C followed by Yamaguchi lactonization accomplished an enantiocontrolled synthesis of epothilone B.

RX(90) OF 319 COMPOSED OF RX(15), RX(16), RX(17), RX(26)
RX(90) BG + BH + BC + BV ==> CJ



86



RX(8) RCT AF 321522-34-3, G 67-56-1
RGT AH 584-08-7 K2003
PRO AI 263761-11-1

83

SOL 67-56-1 MeOH
NTE stereoselective

RX(19) RCT AI 263761-11-1

STAGE(1)
RGT AT 1191-15-7 ALH(Bu-1)2
SOL 75-09-2 CH2Cl2

STAGE(2)
RGT J 12125-02-9 NH4Cl
SOL 7732-18-5 Water

PRO AK 263761-13-3

RX(9) RCT AJ 263768-73-6

STAGE(1)
RGT AM 109-72-8 BuLi
SOL 60-29-7 Et2O, 109-99-9 THF, 7732-18-5 Water

STAGE(2)
RGT AK 263761-13-3
SOL 109-99-9 THF

STAGE(3)
RGT J 12125-02-9 NH4Cl
SOL 7732-18-5 Water

PRO AL 263761-14-4

RX(10) RCT AL 263761-14-4

STAGE(1)
RGT AQ 38721-52-7 L-Selectride
SOL 109-99-9 THF

STAGE(2)
RGT AN 74-88-4
RGT AR 680-31-9 HMPT

STAGE(3)
RGT J 12125-02-9 NH4Cl
SOL 7732-18-5 Water

PRO AO 263761-15-5, AP 321522-35-4

RX(11) RCT AO 263761-15-5

STAGE(1)
RGT AT 1191-15-7 ALH(Bu-1)2
SOL 75-09-2 CH2Cl2

STAGE(2)
RGT J 12125-02-9 NH4Cl
SOL 67-56-1 MeOH, 7732-18-5 Water, 60-29-7 Et2O

PRO AS 312492-69-6

RX(12) RCT AU 187283-44-9

84

RCT CD 7553-56-2 I2

PRO CB 335160-07-1
 NTE stereoselective
 RX(32) RCT CB 335160-07-1, BM 113453-27-3
 RGT CF 7440-66-6 Zn, CG 7440-50-8 Cu
 PRO CE 335160-08-2
 CAT 14221-01-3 Pd(PPh₃)₄
 SOL 71-43-2 Benzene
 NTE stereoselective

RX(33) RCT CE 335160-08-2
 RGT AF 3144-16-9 10-CSA
 PRO CI 335160-09-3
 SOL 67-56-1 MeOH
 NTE stereoselective

RX(34) RCT CI 335160-09-3
 RGT F 79-37-8 (COCl)₂, Z 67-68-5 DMSO
 PRO CJ 335160-10-6
 SOL 73-09-2 CH₂Cl₂
 NTE stereoselective

RX(35) RCT CJ 335160-10-6, AL 187283-45-0

STAGE(1)

STAGE(2)

RCT AM 69739-34-0
 RGT AP 108-48-5 2,6-Lutidine

STAGE(3)


RGT AQ 584-08-7 K₂CO₃
 SOL 67-56-1 MeOH

PRO CK 335160-11-7

NTE stereoselective

REFERENCE COUNT: 19 THERE ARE 19 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L33 ANSWER 17 OF 23 CASREACT COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 134:178371 CASREACT Full-text
 TITLE: Synthesis and biological evaluation of highly potent analogues of epothilones B and D
 AUTHOR(S): Altman, K.-H.; Bold, G.; Caravatti, G.; Florsheimer, A.; Guagnano, V.; Wartmann, M.
 CORPORATE SOURCE: Novartis Pharma AG, TA Oncology Research, Basel, CH-4002, Switz.
 SOURCE: Bioorganic & Medicinal Chemistry Letters (2000), 10(24), 2765-2768
 CODEN: BMCLE8; ISSN: 0960-894X
 PUBLISHER: Elsevier Science Ltd.
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 GI




II

AB A series of new epothilone B and D analogs incorporating fused hetero-aromatic side chains have been prepared. The synthetic strategy is based on olefin I as the common intermediate and allows variation of the side-chain structure in a highly convergent and stereoselective manner. These epothilone analogs, e.g. II, are more potent inhibitors of cancer cell proliferation than the corresponding parent epothilones B or D.

RX(27) OF 370 COMPOSED OF RX(9), RX(2)
 RX(27) AP + AN + J ==> L




AP



2 STEPS

Wartmann, Markus; Altmann, Karl-Heinz
 TA Oncology Research, Novartis Pharma AG, Basel,
 CH-4002, Switz.
 Proceedings of ECSOC-3, [and] Proceedings of ECSOC-4,
 Sept. 1-30, 1999 and 2000 (2000), Meeting Date
 1999-2000, 1431-1442. Editor(s): Pombo-Villari,
 Esteban. Molecular Diversity Preservation
 International: Basel, Switz.
 CODEN: 69AXZT


DOCUMENT TYPE:
 CONFERENCE; (computer optical disk)
 LANGUAGE:
 English




I

AB The authors have synthesized epothilone analogs, e.g. I, with modifications in the northern hemisphere and the heterocyclic side-chain. In all three cases the key steps for construction of the macrocyclic skeleton involve Yamaguchi macrocyclization, the build-up of the requisite seco-acid through aldol reaction between the C7-Cl5 aldehyde and the dianion of the O-protected C1-C6 β -hydroxy acid fragment, and the assembly of the C7-Cl5 aldehyde through the appropriate type of Pd(0)-catalyzed coupling reaction. The IC50 for growth inhibition of the KB-31 tumor cell line for I was 0.45 nM.


RX(284) OF 370 COMPOSED OF RX(29), RX(30), RX(31), RX(32), RX(33), RX(34),
 RX(35)
 RX(284) X + BX + BM + AL + AM ==> CX




Y



BM



AL



CX
 YIELD 308

RX(29) RCT Y 188730-08-7, BX 558-13-4

RGT B2 603-35-0 PPh3

PRO BY 335160-05-9

SOL 75-09-2 CH2Cl2

NTE stereoselective

RX(30)

RCT BY 335160-05-9

RGT K 109-72-8 BUL1

PRO CA 335160-06-0

SOL 109-99-9 THF

NTE stereoselective


RX(31)

RCT CA 335160-06-0

STAGE(1)
 RGT CC 1291-32-3 ZrCp2Cl2



SOL 109-99-9 THF

STAGE(2)




AB A highly convergent total synthesis of the natural products epothilone B and D is described. The route is highlighted by efficient generation of a C12-C13 trisubstituted olefin I which exploits a sequential Nozaki-Hiyama-Kishi coupling and a stereoselective thionyl chloride rearrangement.

RX(53) OF 175 COMPOSED OF RX(6), RX(7), RX(13), RX(14)
RX(53) O + N ==> AN




4
STEPS
→

PAGE 1-A



PAGE 1-B



RX(6) RCT O 355009-06-2
RGT U 7719-09-7 SOCL2
PRO T 355009-08-4
SOL 60-29-7 Et2O, 109-66-0 Pentane

RX(7) RCT T 355009-08-4
RGT Y 22560-16-3 Superhydride
PRO X 210690-99-6
SOL 109-99-9 THF
NTE other product detected

RX(13) RCT X 210690-99-6
RGT AC 26299-14-9 PCC
PRO AM 193146-27-9

RX(14) RCT N 187527-25-9, AM 193146-27-9
RGT AO 4111-54-0 LIN(Pr-1)2
PRO AN 193146-49-5

REFERENCE COUNT: 37
NTE stereoselective, other product detected, no exptl.
THERE ARE 37 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L33 ANSWER 16 OF 23 CASREACT COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 134:311010 CASREACT Full-text
TITLE: Synthetic epothilone analogs with modifications in the
northern hemisphere and the heterocyclic
side-chain-synthesis and biological evaluation
AUTHOR(S): End, Nicole; Bold, Guido; Caravatti, Giorgio;

AU
YIELD 42%

PAGE 1-B

RX(6) RCT Z 4736-60-1

STAGE(1)
RCT AC 109-72-8 BuLi
SOL 109-99-9 THF

STAGE(2)
RCT AD 7553-56-2 I2
SOL 109-99-9 THF

STAGE(3)
RCT AE 1070-89-9 (Me3Si)2N.Na

STAGE(4)
RCT AA 298702-12-2
SOL 109-99-9 THF

STAGE(5)
RCT D 64-19-7 AcOH

PRO AB 298702-07-5
NTE STEREOSELECTIVE

RX(8) RCT AL 224580-52-3

STAGE(1)
RCT AN 280-64-8 9-BBN
SOL 109-99-9 THF

STAGE(2)
RCT AB 298702-07-5
RCT AO 603-32-7 Ph3As, AP 534-17-8 Cs2CO3
CAT 72287-26-4 Palladium, [1,1'-bis(diphenylphosphino-
kp)ferrocene]dichloro-, (SP-4-2)-
SOL 68-12-2 DMF

STAGE(3)
SOL 7732-18-5 Water

STAGE(4)
SOL 60-29-7 Et2O

PRO AM 298702-16-6

73

NTE STEREOSELECTIVE

RX(9) RCT AM 298702-16-6

STAGE(1)
RCT AS 7647-01-0 HCl
SOL 67-56-1 MeOH

STAGE(2)
RCT N 144-55-8 NaHCO3

STAGE(3)
SOL 75-09-2 CH2Cl2

PRO AR 298702-17-7
NTE STEREOSELECTIVE

RX(10) RCT AR 298702-17-7, AT 67-56-1

STAGE(1)
RCT AS 7647-01-0 HCl
CAT 109361-17-3 Ruthenium, bis[(1R)-[1,1'-binaphthalene]-2,2'-
diylbis(diphenylphosphine-kp)]di-μ-
chlorodichloro(N,N-diethylethanamine)di-
SOL 67-56-1 MeOH

STAGE(2)
RCT AV 1333-74-0 H2

STAGE(3)
RCT N 144-55-8 NaHCO3

STAGE(4)
SOL 75-09-2 CH2Cl2

PRO AU 298702-19-9
NTE STEREOSELECTIVE

L33 ANSWER 15 OF 23 CASREACT COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 135:180642 CASREACT Full-text
TITLE: Total Synthesis of Epothilones B and D
AUTHOR(S): Taylor, Richard E.; Chen, Yue
CORPORATE SOURCE: Department of Chemistry & Biochemistry, University of
Notre Dame, Notre Dame, IN, 46556-5670, USA
SOURCE: Organic Letters (2001), 3(14), 2221-2224
CODEN: ORLEF7; ISSN: 1523-7060
PUBLISHER: American Chemical Society
DOCUMENT TYPE: Journal
LANGUAGE: English
GI




74

ACCESSION NUMBER: 135:287591 CASREACT Full-text
 TITLE: Preparation of epothillone intermediates
 INVENTOR(S): Vite, Gregory D.; Kim, Soong-Hoon; Hoeefle, Gerhard
 PATENT ASSIGNEE(S): Bristol-Myers Squibb Company, USA
 SOURCE: PCT Int. Appl., 28 pp.
 CODEN: PIXAD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001073103	A2	20011004	WO 2001-US9620	20010323
WO 2001073103	A3	20020523		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HK, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MX, MY, NZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, VZ, VN, YU, ZA, ZW, AM, AZ, BY, BG, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, FR, GB, GR, IE, IT, LJ, LU, MC, NL, PT, SE, TR, BF, BE, CF, CG, CI, CN, GM, GW, ML, MR, NE, SN, TD, TG				
US 2002042109	A1	20020411	US 2001-811808	20010319
US 6593115	B2	20030715	US 2003-447082	20030528
US 2004023345	A1	20040205	US 2000-191975P	20000324
PRIORITY APPL. INFO.: US 2001-811808 20010319				

OTHER SOURCE(S): MARPAT 135:287591
 AB The present invention relates to a process for the preparation of intermediates useful in the synthesis of epothillone analogs by initially enzymically degrading certain epothillone compds. to form ring-open structures containing a carboxyl group which is esterified, the hydroxyl groups on the moiety protected and the resulting compound oxidized by, e.g. ozone, to form a first intermediate. The first intermediate can be reacted with a triphenylphosphine adduct to yield a compound containing an ester group at position 1 which is subsequently hydrolyzed to form a second intermediate.


RX(4) OF 31 ...O + P ==> Q...



-OMe
 FIELD 558

RX(4) RCT O 190369-98-3
 STAGE(1)
 RGT R 1070-89-9 (Me3Si)2N.Na
 SOL 109-99-9 THF
 STAGE(2)
 RGT P 364336-83-4
 SOL 109-99-9 THF
 PRO Q 364336-77-6

L33 ANSWER 14 OF 23 CASREACT COPYRIGHT 2007 ACS on STN



EP
YIELD 86%

RX(6) RCT AB 370578-43-1

STAGE(1)
RGT AD 121-44-8 Et3N
SOL 75-09-2 CH2Cl2

STAGE(2)
RGT AE 7719-09-7 SOCl2
SOL 75-09-2 CH2Cl2

STAGE(3)
RGT G 7732-18-5 Water

STAGE(4)
RGT AF 429-41-4 Bu4N.F
SOL 109-99-9 THF

PRO AC 370578-22-6
NTE stereoselective

RX(64) RCT AC 370578-22-6, EM 994-30-9

STAGE(1)
RGT AD 121-44-8 Et3N
CAT 1122-58-3 4-DMAP
SOL 75-09-2 CH2Cl2

STAGE(2)
RGT D 144-35-8 NaHCO3
SOL 7732-18-5 Water

PRO DL 370578-62-4

RX(32) RCT DL 370578-62-4

STAGE(1)

RGT CK 7529-22-8 Me-morpholineoxide
SOL 75-63-0 t-BuOH, 109-99-9 THF, 7732-18-5 Water

STAGE(2)
CAT 20816-12-0 OsO4
SOL 7732-18-5 Water

STAGE(3)
RGT DN 7631-90-5 NaHSO3

STAGE(4)
RGT DO 546-67-8 Pb(OAc)4
SOL 141-78-6 AcOEt

PRO DM 342607-03-8

RX(33) RCT DQ 219990-08-6

STAGE(1)
RGT AU 4111-54-0 L1N(Pr-i)2
SOL 109-99-9 THF

STAGE(2)
RGT DM 342607-03-8
SOL 109-99-9 THF

STAGE(3)
RGT AA 12125-02-9 NH4Cl
SOL 7732-18-5 Water

PRO DR 342607-02-7
NTE in-situ generated reagent, stereoselective

RX(66) RCT DR 342607-02-7, FO 17341-93-4
RGT E 110-86-1 Pyridine
PRO EO 342607-17-4
SOL 75-09-2 CH2Cl2

RX(45) RCT EO 342607-17-4

STAGE(1)
RGT CK 7529-22-8 Me-morpholineoxide
SOL 75-63-0 t-BuOH, 109-99-9 THF, 7732-18-5 Water

STAGE(2)
CAT 20816-12-0 OsO4

STAGE(3)
RGT DN 7631-90-5 NaHSO3

STAGE(4)
RGT DO 546-67-8 Pb(OAc)4
SOL 141-78-6 AcOEt

PRO EP 342607-35-6

REFERENCE COUNT: 99 THERE ARE 99 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L33 ANSWER 13 OF 23 CASREACT COPYRIGHT 2007 ACS on STN

STAGE(2)

RGT Z 109-72-8 BULI
SOL 109-99-9 THF

STAGE(3)

RGT CJ 37342-97-5 Hydrozirconocene Cl
SOL 109-99-9 THF

STAGE(4)

RGT CK 7553-56-2 I2
SOL 109-99-9 THF

PRO CH 335160-07-1

RX(30) RCT AO 113453-27-3, CH 335160-07-1

STAGE(1)

RGT CM 12621-78-2 Zinc alloy, base, Zn,Cu
CAT 14221-01-3 Pd(PPh3)4
SOL 71-43-2 Benzene

STAGE(2)

RGT BY 3144-16-9 10-CSA
SOL 67-56-1 MeOH

STAGE(3)

RGT BL 67-68-5 DMSO, AB 79-37-8 (COCl)2
SOL 75-09-2 CH2Cl2

PRO CL 335160-10-6

RX(31) RCT H 187283-45-0, CL 335160-10-6

STAGE(1)

RGT BT 4111-54-0 LIN(Pr-i)2

STAGE(2)

RCT F 69739-34-0
RCT BW 108-48-5 2,6-Lutidine

STAGE(3)

RGT CQ 584-08-7 K2CO3
SOL 67-56-1 MeOH


PRO CP 335160-11-7

REFERENCE COUNT: 32 THERE ARE 32 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L33 ANSWER 12 OF 23 CASREACT COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 135:331283 CASREACT Full-text
TITLE: Stereoselective Syntheses of Epothilones A and B via Nitrile Oxide Cycloadditions and Related Studies
AUTHOR(S): Bode, Jeffrey W.; Carreira, Erick M.
CORPORATE SOURCE: Laboratorium fuer Organische Chemie, ETH-Zuerich, Zurich, CH-8092, Switz.
SOURCE: Journal of Organic Chemistry (2001), 66(19), 6410-6424
CODEN: JOCEAH; ISSN: 0022-3263
PUBLISHER: American Chemical Society
DOCUMENT TYPE: Journal



LANGUAGE: English

GI



AB The expedient and fully stereocontrolled synthesis of epothilones A (I; R = H) and B (I; R = Me) are described. The routes described make extensive use of nitrile oxide cycloaddns. as surrogates for aldol addition reactions and have led to the realization of a highly convergent synthesis based on the Kanemasa hydroxyl-directed nitrile oxide cycloaddn. As well, our synthetic efforts have led to the development of new reaction methodologies and served as the proving ground for several modern methods for asym. carbon-carbon bond formation.

RX(451) OF 751 COMPOSED OF RX(6), RX(64), RX(32), RX(33), RX(66), RX(45)
RX(451) AB + EM + DQ + FO ==> EP



SOL 109-99-9 THE

PRO BR 186692-68-2
NTE stereoselective

RX(28) RCT AX 461044-35-9, BR 186692-68-2
RGT BW 603-32-7 Ph3As, BX 534-17-8 Cs2CO3, BY 280-64-8 9-BBN
PRO BV 461044-42-8
CAT 72287-26-4 Palladium, [1,1'-bis(diphenylphosphino)-

kP)ferrocene]dichloro-, (SP-4-2)-
NTE stereoselective, regioselective

REFERENCE COUNT: 54
THERE ARE 54 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L33 ANSWER 11 OF 23 CASREACT COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 136:318824 CASREACT Full-text

TITLE: Synthetic and semisynthetic analogs of epothilones:
chemistry and biological activity

AUTHOR(S): Altmann, Karl-Heinz; Blommers, Marcel J. J.;
Caravatti, Giorgio; Florsheimer, Andreas; Nicolaou, Kyriacos C.; O'Reilly, Terrence; Schmidt, Alfred;
Schlitzer, Dieter; Wartmann, Markus
CORPORATE SOURCE: TA Oncology Research, Novartis Pharma AG, Basel, CH-4002, Switz.

SOURCE: ACS Symposium Series (2001), 796(Anticancer Agents), 112-130

CODEN: ACSMCS; ISSN: 0097-6156

American Chemical Society

PUBLISHER:





DOCUMENT TYPE:

LANGUAGE:

English

AB Epothilones A and B are naturally occurring microtubule depolymerizing inhibitors, which exhibit potent in vitro antiproliferative activity. Epothilone B is a 3-30-fold more potent inhibitor of human cancer cell growth than paclitaxel in paclitaxel-sensitive cancer cell lines and in paclitaxel-resistant lines exceeds paclitaxel activity by 102 - 103-fold. In addition, epothilone B exhibits potent in vivo antitumor activity even in multidrug-resistant tumor models. In order to gain a better understanding of the structural requirements for epothilone-mediated cytotoxicity and antitumor activity and to discover analogs with similar potency but perhaps better tolerability in vivo, we have investigated a series of structural modifications involving the epoxide site (C12/C13) and the heterocyclic side-chain of epothilones. In this paper we present the synthesis of these analogs and we discuss the impact of such modifications on tubulin polymerization activity as well as cytotoxicity in vitro.

RX(146) OF 320 COMPOSED OF RX(29), RX(30), RX(31)
RX(146) BK + CG + AC + H + F ==> CP



RX(29) RCT BK 188730-08-7, CG 558-13-4



STAGE(1)

RGT CI 603-35-0 PPh3


SOL 75-09-2 CH2Cl2

of novel pyranose synthons. The utility of this very convergent and effective method is demonstrated by the concise total synthesis of epothilones. The stereoselective I-catalyzed aldol condensation of acetaldehyde with (2S)-3-hydroxy-2-methylpropanal gave 2,4-dideoxy-4-methyl-L-erythro-pentopyranose. Oxidation of the latter gave 2,4-dideoxy-4-methyl-L-threo-pentonic acid δ -lactone. Alkylation of the δ -lactone gave 2,4-dideoxy-4-methyl-2-(2-propenyl)-L-xyliconic acid δ -lactone. This compound was a chiral synthon needed for the total synthesis of epothilone C [(4S,7R,8S,13Z,16S)-4,8-dihydroxy-5,5,7,9-tetramethyl-16-[(1E)-1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]oxacyclohexadec-13-ene-2,6-dione].

RX(172) OF 316 COMPOSED OF RX(23), RX(24), RX(25), RX(26), RX(27), RX(28)
 RX(172) BC + BH + BN + BQ + AX ==> BV



6
 STEPS



PAGE 1-B

—OBu-t
 BV
 YIELD 65%

RX(23) RCT BC 247900-97-6
 RGT BF 109-63-7 BF3-Et2O
 PRO BE 461044-38-2
 SOL 7732-18-5 Water, 75-05-8 MeCN
 RCT BE 461044-38-2, BH 109-80-8
 PRO BI 461044-39-3
 CAT 7550-4S-0 TiCl4


RX(24) RCT BI 461044-39-3
 RGT BL 79-37-8 (COCl)2, BM 67-68-5 DMSO
 PRO BK 461044-40-6

RX(25) RCT BK 461044-40-6, BN 184246-51-3
 RGT AR 109-72-8 BuLi
 PRO BO 461044-41-7
 SOL 109-99-9 THF
 NTE stereoselective

RX(26) RCT BO 461044-41-7

STAGE(1)
 RGT BS 7616-83-3 Hg(ClO4)2, BT 471-34-1 CaCO3
 SOL 109-99-9 THF, 7732-18-5 Water

STAGE(2)
 RCT BQ 29949-93-7
 RGT V 680-31-9 HMPF, BU 1070-89-9 (Me3Si)2N.Na



BO
YIELD 93%

RX(27) RCT T 193146-30-4D

STAGE(1)

RCT BY 1070-89-9 (Me3Si)2N.Na
SOL 109-99-9 THF
CON 15 minutes, room temperature

STAGE(2)

RCT BP 346652-91-3
CON 15 minutes, -78 deg C -> -40 deg C

PRO BQ 563829-96-3

NTE solid-supported reaction

REFERENCE COUNT:

73 THERE ARE 73 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L33 ANSWER 9 OF 23 CASREACT COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER:

138:237914 CASREACT Full-text

TITLE:

The total synthesis and biological assessment of

trans-epothillone A

AUTHOR(S):

Altman, Karl-Heinz; Bold, Guido; Caravatti, Giorgio;
Denni, Donatienne; Florsheimer, Andreas; Schmidt,
Alfred; Rins, Greta; Wartmann, Markus
Corporate Research, Novartis Pharma AG, Switz.
Helvetica Chimica Acta (2002), 85(11), 4086-4110
CODEN: HCACAV; ISSN: 0018-019X
Verlag Helvetica Chimica Acta
Journal
English

CORPORATE SOURCE:

Corporate Research, Novartis Pharma AG, Switz.

SOURCE:

Helvetica Chimica Acta (2002), 85(11), 4086-4110

PUBLISHER:

Verlag Helvetica Chimica Acta


DOCUMENT TYPE:

Journal


LANGUAGE:

English

RX(123) OF 496 COMPOSED OF RX(18), RX(19), RX(3), RX(39)
RX(123) BF + BH + CW ==> BV



57



AB The total synthesis of (12S,13S)-trans-epothillone A (I) was achieved based on two different convergent strategies. In a first-generation approach, construction of the C(11)-C(12) bond by PdO-catalyzed Negishi-type coupling between the C(12)-to-C(15) trans-vinyl iodide II and the C(7)-to-C(11) alkyl iodide III preceded the (nonselective) formation of the C(6)-C(7) bond by aldol reaction between the C(7)-to-C(15) aldehyde and the dianion derived from the C(1)-to-C(6) acid IV. The lack of selectivity in the aldol step was addressed in a second-generation approach, which involved construction of the C(6)-C(7) bond in a highly diastereoselective fashion through reaction between the acetonide-protected C(1)-to-C(6) diol ("Schinzer's ketone") and the C(7)-to-C(11) aldehyde. As part of this strategy, the C(11)-C(12) bond was established subsequent to the critical aldol step and was based on B-alkyl Suzuki coupling between the C(1)-to-C(11) fragment and C(12)-to-C(15) trans-vinyl iodide II. Both approaches converged at the stage of the 3-O, 7-O-bis-TBS-protected seco acid, which was converted to trans-deoxyepothillone A via Yamaguchi macrolactonization and subsequent deprotection. Stereoselective epoxidn. of the trans C(12)-C(13) bond could be achieved by epoxidn. with Oxone in the presence of the catalyst 1,2,4,5-di-O-isopropylidene-L-erythro-2,3-hexediure-2,6-pyranose, which provided a 8:1 mixture of I and its (12R,13R)-epoxide isomer (V) in 27% yield (34% based on recovered starting material). The absolute configuration of I was established by X-ray crystallog. I is at least equipotent with natural epothillone A (VI) in its ability to induce tubulin polymerization and to inhibit the growth of human cancer cell lines in vitro. In contrast, the biol. activity of V is at least two orders of magnitude lower than that of VI or I.

58

RGT CF 280-64-8 9-BBN
SOL 109-99-9 THF
CON SUBSTAGE(1) room temperature
SUBSTAGE(2) 3 hours, room temperature

STAGE(2)

RGT E 7732-18-5 Water
CON room temperature

STAGE(3)


RGT CA 186692-68-2
RGT CG 603-32-7 Ph3As, CH 534-17-8 Cs2CO3
CAT 72287-26-4 Palladium, [1,1'-bis(diphenylphosphino)-
κP]ferrocenedichloro-, (SP-4-2)-
SOL 68-12-2 DMF
CON SUBSTAGE(1) room temperature
SUBSTAGE(2) 2 minutes, room temperature
SUBSTAGE(3) room temperature
SUBSTAGE(4) 8 hours, room temperature

STAGE(4)

RGT BS 12125-02-9 NH4Cl
SOL 7732-18-5 Water
CON room temperature



PRO CE 461044-42-8

L33 ANSWER 8 OF 23 CASREACT COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 139:197285 CASREACT Full-text
TITLE: A total synthesis of epothilones using solid-supported
reagents and scavengers
AUTHOR(S): Storer, R. Ian; Takemoto, Toshiyasu; Jackson, Philip
S.F. Ley, Steven V.
CORPORATE SOURCE: University Chemical Laboratories, University of
Cambridge, Cambridge, CB2 1EW, UK
SOURCE: Angewandte Chemie, International Edition (2003),
42(22), 2521-2525
CODEN: AClEF5; ISSN: 1433-7851
PUBLISHER: Wiley-VCH Verlag GmbH & Co. KGaA
DOCUMENT TYPE: Journal
LANGUAGE: English
GI




AB A total synthesis of epothilone C (I) with concomitant formal synthesis of
epothilone A is described, using immobilized reagents and scavengers to effect
multistep synthetic transformations and purifications.

RX(27) OF 210 ...T + BP ==> BQ...




BP

PAGE 1-A



5

STEPS



PAGE 1-B

—OBA-t
CE
YIELD 65%

RX (9) RCT AG 247900-97-6, AI 109-80-8

STAGE (1)
RCT AK 7550-45-0 TIC14
SOL 75-09-2 CH2Cl2
CON SUBSTAGE (1) room temperature
SUBSTAGE (2) room temperature -> -78 deg C
SUBSTAGE (4) 30 minutes

STAGE (2)
RCT J 144-55-8 NaHCO3
SOL 7732-18-5 Water

PRO AJ 461044-39-3

RX (25)

STAGE (1)

53

RGT BV 79-37-8 (COCl)2
SOL 75-09-2 CH2Cl2, 67-68-5 DMSO
CON SUBSTAGE (2) 30 minutes

STAGE (2)

RCT AJ 461044-39-3
SOL 75-09-2 CH2Cl2
CON SUBSTAGE (2) 3 hours

STAGE (3)

RCT AE 121-44-8 Et3N

PRO BU 461044-40-6

RX (26) RCT BX 184246-51-3

STAGE (1)

RGT BF 109-72-8 BuLi
SOL 109-99-9 THF
CON SUBSTAGE (2) 15 minutes

STAGE (2)

RCT BU 461044-40-6
CON SUBSTAGE (2) room temperature

STAGE (3)

RGT BS 12125-02-9 NH4Cl
SOL 7732-18-5 Water

PRO BY 461044-41-7

RX (27) RCT BY 461044-41-7

STAGE (1)

RGT CB 7616-83-3 Hg(ClO4)2, CC 471-34-1 CaCO3
SOL 7732-18-5 Water
CON SUBSTAGE (1) room temperature
SUBSTAGE (2) 2 hours, room temperature

STAGE (2)

SOL 60-29-7 Et2O
CON SUBSTAGE (1) room temperature
SUBSTAGE (2) 10 minutes, room temperature

STAGE (3)

RCT BZ 3020-28-8
RGT BG 680-31-9 HMPT, CD 1070-89-9 (Me3Si)2N.Na
SOL 109-99-9 THF
CON SUBSTAGE (3) room temperature
SUBSTAGE (4) 1 hour

STAGE (4)

RGT BS 12125-02-9 NH4Cl
SOL 7732-18-5 Water

PRO CA 186692-68-2

RX (28) RCT BT 461044-35-9

STAGE (1)

54

RX(14) RCT AS 863981-49-1

STAGE(1)

RGT AU 64-17-5 EtOH
CAT 24057-28-1 Pyridinium tosylate
SOL 64-17-5 EtOH
CON 65 deg C

STAGE(2)

RGT Q 79-37-8 (COCl)₂, AV 67-68-5 DMSO
SOL 75-09-2 CH₂Cl₂
CON -78 deg C

STAGE(3)

RGT AW 121-44-8 Et₃N
CON -78 deg C -> room temperature

PRO AT 688318-68-5

NTE Swern oxidn. in stage 2

RX(15) RCT AV 185148-95-2

STAGE(1)

RGT BA 4111-54-0 LiN(Pr-1)₂
CON -78 deg C

STAGE(2)

RCT AT 688318-68-5
CON -78 deg C

PRO AZ 688318-69-6

NTE stereoselective

REFERENCE COUNT: 24

THERE ARE 24 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L33 ANSWER 7 OF 23 CASREACT COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 139:277113 CASREACT Full-text

TITLE: Synthesis of atorvastatin and epothilone synthons via 2-deoxyribose-5-phosphate aldolase-catalyzed asymmetric aldol condensation of aldehydes
Wong, Chi-huey; Liu, Junjie; De Santis, Grace; Burk, Mark

INVENTOR(S):

PATENT ASSIGNEE(S): The Scripps Research Institute, USA; Diversa Corporation

SOURCE: PCT Int. Appl., 63 pp.

CODEN: PIXAD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003077868	A2	20030925	WO 2003-US7982	20030314
WO 2003077868	A3	20040401		

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KR, KP, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NZ, NO, NZ, OM, PH,

PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW



RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG
CA 2479247 A1 20030925
AU 2003225810 A1 20030929
US 2003232416 A1 20031218
EP 1485498 A2 20041215
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK
JP 200520510 T 20050714
US 2007015260 A1 20070118
PRIORITY APPLN. INFO.:
US 2002-364641P 20020314
US 2003-390544 20030314
WO 2003-US7982 20030314

OTHER SOURCE(S):

MARPAT 139:277113

AB The present invention is based on the discovery that 2-deoxyribose-5-phosphate aldolase (DERA, EC 4.1.2.4) and variants thereof can be used to catalyze sequential asym. aldol reactions between a wide variety of donor and acceptor aldehydes. The reaction products typically contain at least two new stereogenic centers and can be produced in enantiomerically pure form. As such, DERA catalyzed asym. aldol chemical can be exploited to produce synthons for the synthesis of a variety of bioactive mols., e.g. epothilone A.



RX(147) OF 294 COMPOSED OF RX(9), RX(25), RX(26), RX(27), RX(28)
RX(147) AG + AI + BX + BZ + BT ==> CE




AB An efficient synthesis of the epothilone B derivative, 26-fluoroepothilone B (I) was realized by early introduction of the synthetically demanding fluoromethyl epoxide function. The presence of a fluoro substituent results in a remarkable increase in the stability of the epoxide, which tolerates the wide range of reaction conditions required for the fragment coupling step and end game transformations.

RX(175) OF 310 COMPOSED OF RX(10), RX(24), RX(25), RX(26), RX(27), RX(14),

RX(15)
RX(175) H + AG + BW + AY ==> AF



7
STEPS



AZ
YIELD 77%

RX(10) RCT H 226940-40-5, AG 188730-08-7

PRO AH 226940-54-1

SOL 108-88-3 PhMe

CON 40 deg C

NTE stereoselective

RX(24) RCT AH 226940-54-1

STAGE(1)

RGT BY 76-05-1 F3CO2H

SOL 109-99-9 THF, 7732-18-5 Water

CON 23 deg C

STAGE(2)

RCT BW 110-87-2

CAT 24057-28-1 Pyridinium tosylate

SOL 75-09-2 CH2Cl2

CON 23 deg C

PRO EX 688318-79-8

RX(25)

RCT EX 688318-79-8

RGT AJ 1191-15-7 ALH(Bu-i)2

PRO BZ 688318-80-1

SOL 109-99-9 THF

CON SUBSTAGE(1) -78 deg C

SUBSTAGE(2) -78 deg C -> 0 deg C

NTE regioselective

RX(26)

RCT BZ 688318-80-1

RGT AL 75-91-2 t-BuOOH

PRO CA 688318-81-2

CAT 546-68-9 Ti(OPr-i)4, 87-91-2 Di-Et L-tartrate

SOL 75-09-2 CH2Cl2

CON 2 hours, -30 deg C

NTE stereoselective, Katsuki-Sharpless epoxidn., mol. sieves used

RX(27)

RCT CA 688318-81-2



RGT AR 38078-09-0 DAST ((Et2N).SF3)

PRO AS 863981-49-1

SOL 75-09-2 CH2Cl2

CON 23 deg C


NTE using Yarovenko or Ishikawa reagents gave comparable results




2
STEPS

EH

PAGE 1-A



PAGE 1-B



FI
YIELD 93%

RX(50) RCT N 193146-30-4
RGT O 603-35-0D PPh₃
PRO EB 725738-64-7D
SOL 108-88-3 PhMe
CON SUBSTAGE(1) room temperature
SUBSTAGE(2) 18 hours, 90 deg C
SUBSTAGE(3) 90 deg C -> room temperature
NTE attachment to solid-supported reagent Fluka polymer bound triphenylphosphine

RX(54) RCT EB 725738-64-7D

STAGE(1)
RGT CY 1070-89-9 (Me₃Si)₂N.Na
SOL 109-99-9 THF
CON SUBSTAGE(1) room temperature
SUBSTAGE(2) 10 minutes, room temperature
SUBSTAGE(3) room temperature -> -78 deg C


STAGE(2)
RCT EH 346652-91-3
CON SUBSTAGE(1) 1 minute, -78 deg C
SUBSTAGE(2) 10 minutes, -78 deg C

PRO EI 583829-96-3


NTE solid-supported reactant

REFERENCE COUNT: 122
THERE ARE 122 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE REFORMAT


L33 ANSWER 6 OF 23 CASREACT COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 140:423500 CASREACT Full-text
TITLE: Total synthesis of 26-fluoro-epothilone B
AUTHOR(S): Koch, Guido; Loiseleur, Olivier; Altmann, Karl-Heinz
CORPORATE SOURCE: Novartis Institutes for Biomedical Research, Basel, 4002, Switz.
SOURCE: Synlett (2004), (4), 693-697
CODEN: SYNLES; ISSN: 0936-5214
PUBLISHER: Georg Thieme Verlag
DOCUMENT TYPE: Journal
LANGUAGE: English
GI



I



(6) →



RX (6)

RCT AE 823203-10-7

STAGE (1)

SOL 109-99-9 THF

CON 23 deg C -> 0 deg C

STAGE (2)

RCT AG 1070-89-9 (Me3Si)2N.Na

SOL 109-99-9 THF

CON 0 deg C

STAGE (3)

RCT AB 823203-08-3

SOL 109-99-9 THF

CON SUBSTAGE (1) 0 deg C

SUBSTAGE (2) 0 deg C -> 23 deg C

SUBSTAGE (3) 5 hours, 23 deg C

STAGE (4)

RGT R 12125-02-9 NH4Cl

SOL 7732-18-5 Water

CON 23 deg C

PRO AF 823203-09-4

NTE last stage quench

REFERENCE COUNT: 1

THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L33 ANSWER 5 OF 23 CASREACT COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 141:140221 CASREACT Full-text

TITLE: Multi-step application of immobilized reagents and

AUTHOR(S): Storer, R. Ian; Takemoto, Toshiyasu; Jackson, Philip

CORPORATE SOURCE: S.; Brown, Deag S.; Bakendale, Ian R.; Ley, Steven V.

SOURCE: Department of Chemistry, University of Cambridge,

Cambridge, CB2 1EW, UK


CODEN: CEUJED; ISSN: 0947-6539

PUBLISHER: Wiley-VCH Verlag GmbH & Co. KGaA

DOCUMENT TYPE: Journal

LANGUAGE: English

GI




AB The total synthesis of the cytotoxic antitumor natural product epothilone C has provided a stage for the exploitation and further development of immobilized reagent methods. A stereoselective convergent synthetic strategy was applied, incorporating polymer-supported reagents, catalysts, scavengers and catch-and-release techniques to avoid frequent aqueous work-up and chromatog. purification. The enantioselective preparation of 3 key fragments heptanone I, (S)-2-methyl-6-heptenal, and thiazole II along with their elaboration via diastereoselective coupling into epothilone C is presented.

RX(112) OF 662 COMPOSED OF RX(50), RX(54)


RX(112) N + EH ==> EI

L33 ANSWER 4 OF 23 CASREACT COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 142:113814 CASREACT Full-text
TITLE: Method for producing Cl-C15 fragments of epothilones
and derivatives thereof
INVENTOR(S): Klar, Ulrich; Buchmann, Bernd; Schwede, Wolfgang;
Skuballa, Werner
PATENT ASSIGNEE(S): Schering Aktiengesellschaft, Germany
SOURCE: PCT Int. Appl., 48 pp.
CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: German
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2005003071	A1	20050113	WO 2004-EP6685	20040619
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GE, GH, GM, GR, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, MY, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TR, TT, TZ, UA, UG, US, VZ, VN, ZA, ZM, ZW			
RW:	BW, GH, GM, KE, LS, MW, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, BG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GN, GW, ML, MR, NE, SN, TD, TG			
DE 10331004	A1	20050224	DE 2003-10331004	20030703
AU 2004254200	A1	20050113	AU 2004-254200	20040619
CA 2531078	A1	20050113	CA 2004-2531078	20040619
EP 1641734	A1	20060405	EP 2004-740122	20040619
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK, HR			
CN 1816514	A	20060809	CN 2004-80019005	20040619
BR 2004012179	A	20060822	BR 2004-12179	20040619
IN 2006DN0056	A	20070824	IN 2006-DN56	20060103
MX 2006PA00172	A	20060427	MX 2006-PA172	20060105
NO 2006000554	A	20060403	NO 2006-554	20060202
US 2007142675	A1	20070621	US 2006-563058	20060619
PRIORITY APPLN. INFO.:			DE 2003-10331004	20030703
OTHER SOURCE(S):			WO 2004-EP6685	20040619
GI			MARPAT 142:113814	






I



II

AB The invention relates to a method for preparing Cl-C15 fragments I [R1a, Rib = H, Cl-10-alkyl, aryl, C7-20-alkyl; R1aRib = (CH2)m; m = 2-5; R2a, R2b = H, Cl-10-alkyl, C2-10-alkenyl, C2-10-alkynyl, aryl, C7-20-alkyl; R2aR2b = (CH2)n; n = 2-5; R3 = H, Cl-10-alkyl, aryl, C7-20-alkyl; R4a, R4b = H, Cl-10-alkyl, aryl, C7-20-alkyl; R4aR4b = (CH2)p; p = 2-5; R5 = H, Cl-10-alkyl, aryl, C7-20-alkyl; R6, R7 = H; R6R7 = bond, O; G = X:CR8, bi- or tricyclic aryl; R8 = H, halogen, (un)substituted Cl-20-alkyl, aryl, C7-20-alkyl; X = O, (OR23)2, C2-10-alkylene- α,ω -dioxy, H(OR9), CRIOR11; R23 = Cl-20-alkyl; R9 = H, protecting group; R10, R11 = H, Cl-10-alkyl, aryl; C7-20-alkyl; CRIOR11 = 5- to 7-membered carbocycle; R13 = CH2OR13a, CH2-halo, CHO, CO2R13b, CO-halo; R13a, R14a = H, SO2alkyl, SO2-aryl, SO2-alkyl; R13aR14a = (CH2)o, CRI5aR15b; o = 2-4; R13b, R14b = H, Cl-10-alkyl, aryl, C7-20-alkyl; R15a, R15b = H, Cl-10-alkyl, aryl, C7-20-alkyl; R15aR15b = (CH2)q; q = 3-6; R20 = O-PG, NHR29, N3; Z = O, H(OR12); R12 = H, PG of epothilones and derivatives. The procedure comprises the bonding of a Cl-C6 fragment, R13CH2CHRI4CRIaR1bC(O)CH2R2aR2b, to a C7-Cl12 fragment, R5C(V)(CH2)3CR4aR4bC(W)R3a [V, W = O, (OR23)2, C2-10-alkylene- α,ω -dioxy, H(OR9)], to form a Cl-Cl12 fragment, R5C(V)(CH2)3CR4aR4bCR3a(O-PG14)CR2aR2bC(I)2CRIaR1bCHRI4CH2R13 [PG = H, protecting group], which is then treated with a Cl13-Cl15 fragment, G-CR20'CH2CHR7'R21 [R7' = H; R20' = halogen, N3, NHR29, OH, O-PG, NR29-PG, Cl-20-(perfluoro)alkylsulfonyloxy, (Cl-4-alkyl, NO2, Cl, Br-substituted) benzyl, NR29SO2Me, NR29C(O)Me, CH2C(O)Me; R21 = OH, halo, O-PG, P-Ph3Hal- (Hal = F, Cl, Br, I), P(O)(OO)2 (Q = Cl-10-alkyl, Ph), P(O)Ph2; R29 = H, Cl-6-alkyl], to form the Cl-Cl15 epothilone intermediate product I. Thus, I [R1a = Rib = R5 = Me, R2a = CH2CH:CH2- β , R2b = β , R20 = OSiMe2OMe3- α , G = 2-methylbenzothiazol-5-yl, PG = SiMe2OMe3, Z = O] was prepared from (S)-4-(2-methyl-3-oxohept-6-en-2-yl)-2,2-dimethyl-1,3-dioxane via lithiation and reaction with (2S,6RS)-2-methyl-6-[(tert-butylidimethylsilyl)oxy]heptanal, tetrahydropyranylation, desilylation with BuNF in THF, oxidation in CH2Cl2 containing N-methylmorpholine N-oxide and catalytic tetrapropylammonium perruthenate, Wittig reaction with [(3S)-3-(2-methylbenzothiazol-5-yl)propyl]triphenylphosphonium iodide, desisopropylideneation/detetracyclopentanylation with catalytic 4-MeOC6H4SO3H in EtOH, silylation with CF3SO2SiMe2OMe3, regioselective desilylation with (1)-camphor-10'-sulfonic acid, Swern oxidation with DMSO/(COCl)2 in CH2Cl2 and carbonyl oxidation with NaOCl2 in aqueous THF/Me3COH. The produced Cl-C15 epothilone intermediate products can be converted into the intrinsically active ingredients II [AK = OC(O), OCH2, CH2C(O), NR29C(O), NR29SO2; R29 = H, Cl-6-alkyl] according to known methods. The invention also relates to the corresponding Cl-Cl12 fragments.

RX(6) OF 60 ...AE + AB ==> AF...



CA
YIELD 31%

RX(22) RCT AV 861124-76-7

STAGE(1)
RCT BV 1070-89-9 (Me3Si)2N.Na
SOL 109-99-9 THF
CON 15 minutes, 0 deg C

STAGE(2)
RCT BS 190370-00-4
SOL 109-99-9 THF
CON 12 hours, -20 deg C

41

STAGE(3)
RCT M 12125-02-9 NH4Cl
SOL 7732-18-5 Water

PRO BU 861124-92-7

RX(23) RCT BU 861124-92-7

STAGE(1)
RCT BX 3144-16-9 10-CSA
SOL 67-56-1 MeOH, 75-09-2 CH2Cl2
CON SUBSTAGE(1) 5 minutes, 0 deg C
SUBSTAGE(2) 0.5 hours, 0 deg C
SUBSTAGE(3) 1 hour, 25 deg C

STAGE(2)
RCT BI 121-44-8 Et3N

PRO BW 861124-94-9

RX(24) RCT BW 861124-94-9

STAGE(1)
RCT BZ 26412-87-3 Pyridine-SO3 (1:1), BO 67-68-5 DMSO, BI 121-44-8 Et3N
SOL 75-09-2 CH2Cl2
CON 0.5 hours, 25 deg C

STAGE(2)
RCT M 12125-02-9 NH4Cl
SOL 7732-18-5 Water, 60-29-7 Et2O

PRO BY 861124-97-2

RX(25) RCT S 187283-45-0, BY 861124-97-2

STAGE(1)
RCT BZ 26412-87-3 Pyridine-SO3 (1:1), BO 67-68-5 DMSO, BI 121-44-8 Et3N
SOL 75-09-2 CH2Cl2
CON 0.5 hours, 25 deg C

STAGE(2)
RCT M 12125-02-9 NH4Cl
SOL 7732-18-5 Water, 60-29-7 Et2O

STAGE(3)
RCT J 69739-34-0
RCT L 108-48-5 2,6-Lutidine
SOL 75-09-2 CH2Cl2
CON 2 hours, 0 deg C

STAGE(4)
RCT BE 7647-01-0 HCl
SOL 7732-18-5 Water

STAGE(5)
RCT CB 584-08-7 K2CO3
CON 15 minutes, 25 deg C

42

CON SUBSTAGE(1) 5 deg C
SUBSTAGE(2) 1 minute, 5 deg C

STAGE(2)

RCT T 220775-18-8
SOL 109-99-9 THF
CON 5 hours

PRO BD 924727-13-9

RCT BD 924727-13-9
PRO BF 279226-51-6
CAT 104-15-4 TsOH
SOL 64-17-5 EtOH
CON 2 hours, room temperature

RX(17)

STAGE(1)

RCT AF 67-68-5 DMSO, BH 79-37-8 (COCl)₂
SOL 75-09-2 CH₂Cl₂
CON SUBSTAGE(1) -78 deg C
SUBSTAGE(2) 10 minutes, -78 deg C

STAGE(2)

RCT BF 279225-51-6
SOL 75-09-2 CH₂Cl₂
CON 30 minutes, -78 deg C

STAGE(3)

RGT AG 121-44-8 Et₃N
CON -78 deg C -> -10 deg C

PRO BG 279226-52-7
NTE Wittig reaction

RX(18)

RCT BI 305840-13-5

STAGE(1)

RGT BL 4111-54-0 LIN(Pr-i)₂
SOL 109-99-9 THF
CON SUBSTAGE(1) -70 deg C
SUBSTAGE(2) 10 minutes, -70 deg C
SUBSTAGE(3) 1 hour, -30 deg C
SUBSTAGE(4) -30 deg C -> -70 deg C

STAGE(2)

RGT AL 7646-85-7 ZnCl₂
SOL 109-99-9 THF
CON SUBSTAGE(1) -70 deg C
SUBSTAGE(2) 10 minutes, -70 deg C
SUBSTAGE(3) 15 minutes, -70 deg C

STAGE(3)

RCT BG 279226-52-7
SOL 109-99-9 THF
CON SUBSTAGE(1) 30 minutes, -70 deg C
SUBSTAGE(2) 3.5 hours, -70 deg C

STAGE(4)

RCT BJ 69739-34-0

PRO BK 924727-14-0

REFERENCE COUNT: 39

THERE ARE 39 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L33 ANSWER 3 OF 23 CASREACT COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 143:172686 CASREACT Full-text

TITLE: New method for totally synthesizing natural product-

epothilones

INVENTOR(S): Yan, Jiaqi

Peop. Rep. China

PATENT ASSIGNEE(S): Faming Zhuanli Shengding Gongkai Shuomingshu, 29 pp.

SOURCE: CODEN: CNXKEV

DOCUMENT TYPE: Patent


LANGUAGE: Chinese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
CN 1418881	A	20030521	CN 2002-153675	20021205
PRIORITY APPLN. INFO.:				
CN 2002-153675 20021205				

GI



1

AB The invention discloses a novel multi-step synthetic method for preparing Epothilone A and epothilone B (I; R = H or Me resp.) in a convergent approach starting from 2,2-dimethyl-3-oxopentanal, propionaldehyde SAMP hydrazone, and Et 2-methylthiazolidin-4-ylcarboxylate.

RX(97) OF 191 COMPOSED OF RX(22), RX(23), RX(24), RX(25)
RX(97) AV + BS + S + J -> CA

STAGE (2)

RCT P 865535-39-3
RGT V 534-17-8 Cs2CO3
CAT 603-32-7 Ph3As, 72287-26-4 Palladium, [1,1'-bis(diphenylphosphino)-κP]ferrocene]dichloro-, (SP-4-2)-
SOL 68-12-2 DMF
CON 2 hours, 0 deg C -> room temperature

PRO T 865535-59-7

RX(11) RCT T 865535-59-7
RGT AQ 1310-65-2 LiOH
PRO AP 865535-60-0
SOL 7732-18-5 Water, 67-63-0 Me2CHOH
CON 2.5 hours, 60 deg C

REFERENCE COUNT: 33
THERE ARE 33 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L33 ANSWER 2 OF 23 CASREACT COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 146:229070 CASREACT Full-text

TITLE: Total synthesis and antitumor activity of ZK-EPO: The first fully synthetic epothilone in clinical development


AUTHOR(S): Klar, Ulrich; Buchmann, Bernd; Schwede, Wolfgang; Skuballa, Werner; Hoffmann, Jens; Lichtner, Rosemarie B.

CORPORATE SOURCE: Schering AG, Research Center Europe, Berlin, Germany
SOURCE: Angewandte Chemie, International Edition (2006), 45(47), 7942-7948

CODEN: ACTEF5; ISSN: 1433-7851


WILEY-VCH Verlag GmbH & Co. KGaA

PUBLISHER: Journal
DOCUMENT TYPE: English
LANGUAGE: GI




AB From about 350 active epothilone analogs synthesized by a highly convergent synthesis, ZK-EPO (1) was chosen for clin. development on the basis of its outstanding preclin. data. This compound exhibits higher activity and efficacy than taxanes, such as paclitaxel and second-generation epothilones, a fast and efficient cellular uptake, no recognition by efflux mechanisms, and an improved therapeutic window.


RX(84) OF 314 COMPOSED OF RX(15), RX(16), RX(17), RX(18)
RX(84) BC + T + BI + BJ ==> BK



BC



BI



4
STEPS

BK
YIELD 73%

RX(15) RCT BC 823203-10-7


STAGE(1)

RGT BE 1070-89-9 (Me3Si)2N.Na
SOL 109-99-9 THF

100.0% DONE 3173 VERIFIED 468 HIT RXNS 23 DOCS
SEARCH TIME: 00.00.04



=> d ibib abs fhlt L33 1-23

L33 ANSWER 1 OF 23 CASREACT COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 146:251631 CASREACT Full-text
TITLE: Total synthesis and biological assessment of
benzimidazole-based analogs of epothilone A:
Ambivalent effects on cancer cell growth inhibition
Cachoux, Frederic; Isarno, Thomas; Wartmann, Markus;
Altman, Karl-Heinz
CORPORATE SOURCE: Prestwick Chemical, Illkirch, Fr.
SOURCE: ChemBioChem (2006), 7(1), 54-57
CODEN: CBCHFX; ISSN: 1439-4227
PUBLISHER: Wiley-VCH Verlag GmbH & Co. KGaA
DOCUMENT TYPE: Journal
LANGUAGE: English
GI



AB The title (12R,13S)- and (12S,13S)-epoxy-benzimidazole epothilone derivs. I
(R12 = β -, α -H, resp.), as well as the corresponding (12Z)- and (12E)- Δ 12-
olefin epoxide precursors, were prepared and evaluated for inhibition of
growth of human cancer cell lines, such as KB-31 and KB-8511.

RX(30) OF 64 COMPOSED OF RX(10), RX(3), RX(4), RX(11)
RX(30) AN + AC + S \rightarrow AP



RX(10) RCT AN 3020-28-8

STAGE(1)

RGT AO 1070-89-9 (Me3Si)2N.Na
SOL 109-99-9 THF
CON 15 minutes, room temperature

STAGE(2)

RCT AC 279226-82-3
CON 30 minutes, -78 deg C

PRO O 865535-38-2
NTE stereoselective, Wittig reaction

RX(3)


RCT O 865535-38-2
RGT Q 3144-16-9 10-CSA
PRO P 865535-39-3
SOL 67-56-1 MeOH, 75-09-2 CH2Cl2
CON 17 hours, room temperature

RX(4)

RCT S 279227-12-2

STAGE(1)

RGT U 280-64-8 9-BEN
SOL 109-99-9 THF
CON 90 minutes, room temperature



chain nodes :
 2 3 4 5 9 10 11 12 13 14 15 17 18 19 21 22 23 24 25 26 27 28
 32 33 34 36 48 50 51 58 59 60 61 62 64 65 67 68 71 72 73 74 75
 77 78 80 81
 83 84 85 88 89 91 92 93 95 96 97 101 106 107 108 109 110 111 112
 113 114 115
 116 117 118 121 123
 ring nodes :
 7 8 49 52 53 54
 ring/chain nodes :
 1 6 16 20 35 37 38 39 40 45 98 99 100
 chain bonds :
 1-2 2-3 2-9 2-10 3-4 3-11 3-12 4-14 4-13 4-95 5-93 5-48 5-15 5-95 6-92
 6-93 7-17 8-18 8-19 20-21 20-22 22-23 22-24 22-25 25-26 25-96 25-106
 32-33 32-34
 35-36 49-50
 49-51 58-59 58-60 58-61 58-89 59-62 59-71 64-65 64-88 67-68
 72-73 72-74
 72-75 77-78 80-81 83-84 84-85 91-92 100-101 107-108 107-112 107-114 107-118
 121 108-109 108-110
 108-111 109-113 109-123 115-116 116-117 116-118
 ring/chain bonds :
 1-20 6-16 35-37 38-39 38-40 98-99
 ring bonds :
 7-8 7-53 8-49 49-52 52-54 53-54
 exact/norm bonds :
 1-2 1-20 2-3 4-95 5-93 5-48 5-15 5-95 6-92 6-16 6-93 7-8 7-53 8-49
 20-22 22-23 25-96 25-106 35-37 38-39 38-40 49-52 52-54 53-54 58-89 59-71
 64-88 67-68
 72-75 80-81 83-84 84-85 91-92 98-99 107-114 107-121 109-123 115-116 116-118

exact bonds :
 2-9 2-10 3-11 3-12 4-14 4-13 7-17 8-18 8-19 20-21 22-23 22-24 25-26
 32-33 32-34 35-36 49-50 49-51 58-59 58-60 58-61 59-62 64-65 72-73 72-74
 77-78 100-101
 107-108 107-112 108-109 108-110 108-111 109-113

G1: (*1), (*2)
 G2: (*3), (*4), (*5)
 G3: H, (*6)
 G4: X, OH, O, (*7)
 G5: (*8), (*9), (*10), (*11)
 G6: (*12), (*13), (*14)
 G7: (*15), (*16), (*17)
 G8: O, N, X, (*18)
 G9: O, P, X


Match level :

1:CLASS 2:CLASS 3:CLASS 4:CLASS 5:CLASS 6:CLASS 7:CLASS 8:CLASS 9:CLASS
 10:CLASS 11:CLASS 12:CLASS 13:CLASS 14:CLASS 15:CLASS 16:CLASS 17:CLASS
 18:CLASS 19:CLASS
 20:CLASS 21:CLASS 22:CLASS 23:CLASS 24:CLASS 25:CLASS 26:CLASS 27:CLASS
 28:CLASS 32:CLASS
 33:CLASS 34:CLASS
 35:CLASS 36:CLASS 37:CLASS 38:CLASS 39:CLASS 40:CLASS
 45:CLASS 48:CLASS
 49:Atom 50:CLASS 51:CLASS 52:Atom 53:Atom 54:Atom 58:CLASS 59:CLASS
 60:CLASS 61:CLASS
 62:CLASS 64:CLASS 65:CLASS 67:CLASS 68:CLASS 71:CLASS 72:CLASS 73:CLASS
 74:CLASS 75:CLASS
 77:CLASS 78:CLASS
 89:CLASS 91:CLASS
 92:CLASS 93:CLASS 95:CLASS 96:CLASS 97:Atom 98:CLASS 99:CLASS 100:CLASS
 101:CLASS 106:CLASS
 107:CLASS 108:CLASS 109:CLASS 110:CLASS 111:CLASS 112:CLASS 113:CLASS
 114:CLASS 115:CLASS
 116:CLASS 117:CLASS
 118:CLASS 121:CLASS 123:CLASS
 Generic attributes :
 97:
 Saturation : Unsaturated
 Type of Ring System : Polycyclic

fragments assigned product role:
 containing 1
 fragments assigned reactant/reagent role:
 containing 107
 node mappings:
 22:108

L33 23 SEA FILE=CASREACT SUB=L30 SSS FUL L31 (468 REACTIONS)

Structure attributes must be viewed using STN Express query preparation:
Uploading Lib.str



chain nodes :
2 3 4 5 9 10 11 12 13 14 15 17 18 19 21 22 23 24 25 26 27 28
32 33 34 36 48 50 51 58 59 60 61 62 64 65 67 68 71 72 73 74 75
77 78 80 81
83 84 85 88 89 91 92 93 95 96 97 101 106
ring nodes :
7 8 49 52 53 54
ring/chain nodes :
1 6 16 20 35 37 38 39 40 45 98 99 100
chain bonds :
1-2 2-3 2-9 2-10 3-4 3-11 3-12 4-14 4-13 4-95 5-93 5-48 5-15 5-95 6-
92
6-93 7-17 8-18 8-19 20-21 20-22 22-23 22-24 22-25 25-26 25-96 25-106
32-33 32-34
35-36 49-50 49-51 58-59 58-60 58-61 58-89 59-62 59-71 64-65 64-88 67-68
72-73 72-74
72-75 77-78 80-81 83-84 84-85 91-92 100-101
ring/chain bonds :
1-20 6-16 35-37 38-39 38-40 98-99
ring bonds :
7-8 7-53 8-49 49-52 52-54 53-54
exact/norm bonds :
1-2 1-20 2-3 3-4 4-95 5-93 5-48 5-15 5-95 6-92 6-16 6-93 7-8 7-53 8-
49
20-22 22-25 25-96 25-106 35-37 38-39 38-40 49-52 52-54 53-54 58-89 59-71
64-88 67-68
72-75 80-81 83-84 84-85 91-92 98-99
exact bonds :
2-9 2-10 3-11 3-12 4-14 4-13 7-17 8-18 8-19 20-21 22-23 22-24 25-26
32-33 32-34 35-36 49-50 49-51 58-59 58-60 58-61 59-62 64-65 72-73 72-74

31

77-78 100-101

G1: (*1), (*2)
G2: (*3), (*4), (*5)
G3: H, (*6)
G4: X, OH, O, (*7)
G5: (*8), (*9), (*10), (*11)
G6: (*12), (*13), (*14)
G7: (*15), (*16), (*17)

Match level :

1:CLASS 2:CLASS 3:CLASS 4:CLASS 5:CLASS 6:CLASS 7:CLASS 8:CLASS 9:CLASS
10:CLASS 11:CLASS 12:CLASS 13:CLASS 14:CLASS 15:CLASS 16:CLASS 17:CLASS
18:CLASS 19:CLASS
20:CLASS 21:CLASS 22:CLASS 23:CLASS 24:CLASS 25:CLASS 26:CLASS 27:CLASS
28:CLASS 32:CLASS
33:CLASS 34:CLASS 35:CLASS 36:CLASS 37:CLASS 38:CLASS 39:CLASS 40:CLASS
45:CLASS 48:CLASS
49:Atom 50:CLASS 51:CLASS 52:Atom 53:Atom 54:Atom 58:CLASS 59:CLASS
60:CLASS 61:CLASS
62:CLASS 64:CLASS 65:CLASS 67:CLASS 68:CLASS 71:CLASS 72:CLASS 73:CLASS
74:CLASS 75:CLASS
77:CLASS 78:CLASS 80:CLASS 81:CLASS 83:CLASS 84:CLASS 85:CLASS 88:CLASS
89:CLASS 91:CLASS
92:CLASS 93:CLASS 95:CLASS 96:CLASS 97:Atom 98:CLASS 99:CLASS 100:CLASS
101:CLASS 106:CLASS

Generic attributes :

97:
Saturation : Unsaturated
Type of Ring System : Polycyclic

L3 560 SEA FILE-REGISTRY SSS FUL L1
L30 69 SEA FILE-CASREACT ABB-ON PLU-ON L3
L31 STR

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *


Structure attributes must be viewed using STN Express query preparation:
Uploading L31b.str

32

AT 368036 T 20070815 AT 1998-946309 19980810
 US 2003144523 A1 20030731 US 2000-485292 20000503
 PRIORITY APPLN. INFO.: DE 1997-19735574 A 19970809
 DE 1997-19735575 A 19970809
 DE 1997-19735576 A 19970809
 DE 1997-19748928 A 19971024
 DE 1997-19749717 A 19971031
 DE 1997-19751200 A 19971113
 DE 1998-19813821 A 19980320
 WO 1998-EP5064 W 19980810

OTHER SOURCE(S): CASREACT 130:168162; MARPAT 130:168162

GI



AB Compds. I (R1 = H, OH, OR7; R2 = H, protective group; R3, R4 = H, Cl-10-alkyl, C7-10-alkyl; R3R4 = (CH2)m; R5, R6 = H, Cl-10-alkyl, aryl, C7-20-aralkyl; R7 = Cl-20-alkyl, C7-20-aralkyl; R1 ≠ OH, when R2 = SiMe2OMe3, R3 = R4 = R5 = Me, R6 = H), useful for the preparation of epothilone and epothilone derivs., are prepared from (R)-, (S)- or (i)-pantolactone (II; R8 = H) via tetrahydropyranation with 3,4-dihydro-2H-pyran and pyridinium P-toluenesulfonate and reduction of THP ether II (R8 = THP) with DIBAL-H, Wittig reaction of lactol III with MePh3P+Br-/BuLi, oxidation of pentenol IV (R9 = CH:CH2, R10 = CH2OH) with (COCl)2/DMSO in CH2Cl2, addition of an organometallic compound, R5CH2Y (Y = Li, MgX, X = Cl, Br, I), to aldehyde IV (R9 = CH:CH2, R10 = CHO), hydroboration and oxidation of hexenol IV (R9 = CH:CH2, R10 = CH(OH)CH2R5) with N-methylmorpholine N-oxide/TPAP and oxidation of keto aldehyde IV (R9 = CH2CHO, R10 = COCH2R5) to keto acid IV (R9 = CH2CO2H, R10 = COCH2R5) or hexenol IV (R9 = CH:CH2, R10 = CH(OH)CH2R5) can be oxidized then alkylated with LiN(CHMe2)2 and R6Z (Z = leaving group) and the resulting ketone IV (R9 = CH:CH2, R10 = COCH2R5R6), can hydroborated and oxidized as above, leading to I.

=> file registry
 FILE 'REGISTRY' ENTERED AT 12:16:30 ON 11 OCT 2007
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
 COPYRIGHT (C) 2007 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 10 OCT 2007 HIGHEST RN 950149-06-1
 DICTIONARY FILE UPDATES: 10 OCT 2007 HIGHEST RN 950149-06-1

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH June 29, 2007

Please note that search-term pricing does apply when conducting SmartSELECT searches.

REGISTRY includes numerically searchable data for experimental and predicted properties as well as tags indicating availability of experimental property data in the original document. For information on property searching in REGISTRY, refer to:

<http://www.cas.org/support/stngen/stdoc/properties.html>

=> file casreact

FILE 'CASREACT' ENTERED AT 12:16:34 ON 11 OCT 2007
 USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT
 COPYRIGHT (C) 2007 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications.

FILE CONTENT:1840 - 6 Oct 2007 VOL 147 ISS 16

New CAS Information Use Policies, enter HELP USAGETERMS for details.

 * CASREACT now has more than 13.8 million reactions *
 * *****

Some CASREACT records are derived from the ZIC/VINITI database (1974-1999) provided by InfoChem, INPI data prior to 1986, and Biotransformations database compiled under the direction of Professor Dr. Klaus Kieselich.

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> d stat que L33
 L1 STR

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

protecting group; R11 = H, protecting group] including all the stereoisomers and their mixts. are prepared E.g., title compound (S)-III [R5 = R6 = Me, R9 = R11 = H, R10 = TBDMS] was prepared in 6 steps from D-(-)-pantolactone via reaction with 3,4-dihydro-2H-pyran, hydride reduction, Wittig reaction with methyltriphenylphosphonium bromide, protection of OH with TBDMS-Cl, de-tetrahydropyranyl, and reduction with borane-THF.

LI02 ANSWER 23 OF 24 CAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 1999:116658 CAPLUS Full-text
DOCUMENT NUMBER: 130:168163

TITLE: New (C13-C15)-fragments, method for their preparation and their application for synthesis of epothilone and epothilone derivatives

INVENTOR(S): *Klar, Ulrich; Schwede, Helfgang; Skuballa, Werner; Buchmann, Bernd;*

Schirmer, Michael

Schering A.-G., Germany

Get. Offen., 14 pp.

CODEN: GWXXBX

DOCUMENT TYPE: Patent

LANGUAGE: German


FAMILY ACC. NUM. COUNT: 4

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 19735575	A1	19990211	DE 1997-19735575	19970809
CA 2299608	A1	19990218	CA 1998-2299608	19980810
WO 9907692	A2	19990218	WO 1998-EP5064	19980810
WO 9907692	A3	19990514		
W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DK, EE, ES, FI, GB, GE, GH, GU, HU, ID, IL, IS, JP, KE, KP, KR, KZ, LC, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW				
RW: GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
AU 9893409	A	19990301	AU 1998-93409	19980810
EP 1005465	A2	20000607	EP 1998-946309	19980810
EP 1005465	B1	20070725		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, CY, AL, MK				
JP 2001512723	T	20010828	JP 2000-506196	19980810
AT 368036	T	20070815	AT 1998-946309	19980810
US 2003144523	A1	20030731	US 2000-485292	20000503
PRIORITY APPLN. INFO.:				
DE 1997-19735575	A	19970809		
DE 1997-19735575	A	19970809		
DE 1997-19735575	A	19970809		
DE 1997-19748926	A	19971024		
DE 1997-19749717	A	19971031		
DE 1997-19751200	A	19971113		
DE 1998-19813821	A	19980320		
WO 1998-EP5064	W	19980810		

OTHER SOURCE(S): CASREACT 130:168163; MARPAT 130:168163

GI



AB The title compds. [I; II; R1 = H, alkyl, aryl, aralkyl; R2 = H, protecting group; R3 = OH, halo, OR6; R6 = protecting group; R4 = H, alkyl; R5 = H, alkyl, aryl, aralkyl] are prepared E.g., title compound I [R1 = Me, R2 = TBDMS, R3 = TBDMS] was prepared in 6 steps from L-(-)-malic acid via cyclization, 3-O-protection of 3(S)-hydroxy-2-tetrahydrofuranone, hydride reduction, ring opening and chain lengthening, 1-O-protection of 3(S)-tert-butylphenylsilyloxy-1,4-pentadiol, and oxidation of 3(S)-tert-butylphenylsilyloxy-5-(tert-butylidimethylsilyloxy)-2-pentanol. This was further treated with Et (2-methyl-4-thiazolylmethyl)phosphonate in THF-hexane containing BuLi to give (E,3S)-1-[[dimethyl(1,1-dimethylethyl)silyloxy]-3-[[[(1,1-dimethylethyl)diphenylsilyloxy]-4-methyl-4-(2-methylthiazol-4-yl)pent-4-ene.

LI02 ANSWER 24 OF 24 CAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 1999:116657 CAPLUS Full-text
DOCUMENT NUMBER: 130:168162

TITLE: New method for the preparation of the C(1)-C(6)-segment of epothilone and epothilone derivatives

INVENTOR(S): *Klar, Ulrich; Schwede, Helfgang; Skuballa, Werner; Buchmann, Bernd;*

Schirmer, Michael

Schering A.-G., Germany

Get. Offen., 17 pp.

CODEN: GWXXBX

DOCUMENT TYPE: Patent

LANGUAGE: German

FAMILY ACC. NUM. COUNT: 4


PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 19735574	A1	19990211	DE 1997-19735574	19970809
CA 2299608	A1	19990218	CA 1998-2299608	19980810
WO 9907692	A2	19990218	WO 1998-EP5064	19980810
WO 9907692	A3	19990514		
W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DK, EE, ES, FI, GB, GE, GH, GU, HU, ID, IL, IS, JP, KE, KP, KR, KZ, LC, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW				
RW: GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
AU 9893409	A	19990301	AU 1998-93409	19980810
EP 1005465	A2	20000607	EP 1998-946309	19980810
EP 1005465	B1	20070725		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, CY, AL, MK				
JP 2001512723	T	20010828	JP 2000-506196	19980810

DE 1998-19813821 A 19980320
WO 1998-EP5064 W 19980810

MARPAT 130:196529

OTHER SOURCE(S):
GI



II

AB Epothilone derivs. of formula I (X = O, alkylene-q, q-dioxy, two alkoxy groups, etc.; Y = O, H₂; Z = O, (H, OH), (H, protected OH); R_{1a}, R_{1b} = H, alkyl, aryl, aralkyl, or together = (CH₂)_m where m = 2, 3, 4, 5; R_{2a}, R_{2b} = H, alkyl, aryl, aralkyl, or together = (CH₂)_n where n = 2, 3, 4, 5; when D-E = CH₂CH₂ or when Y = O, R_{2a} or R_{2b} may not be H/Me; R₃ = H, alkyl, aryl, aralkyl; R_{4a}, R_{4b} = H, alkyl, aryl, aralkyl, or together = (CH₂)_p where p = 2, 3, 4, 5; D-E = CH₂CH₂, CH₂CH, C-tripbond, C, 2,3-oxiranedyl, CH(OH)CH(OH), CH(OH)CH₂; R₅ = H, alkyl, aryl, aralkyl; R₆, R₇ = H, together = a saturated bond or O; R₈ = H, alkyl, aryl, aralkyl all of which may be substituted) are prepared. Thus, the title compds. (4S,7R,8S,9S,13E,16S(E))- and (4S,7R,8S,9S,13Z,16S(E))-4,8-dihydroxy-7-ethyl-16-(1-methyl-2-(2-methyl-4-thiazolyl)ethenyl)-1-oxa-5,9,13-tetramethylcyclohexadec-13-en-2,6-dione (II) were prepared in many steps. The new compds. interact with tubulin by stabilizing formed microtubuli. They are capable of influencing cell division in a phase-specific manner and are suitable for the treatment of malignant tumors, such as ovarian, gastric, colon, breast, lung, head and neck carcinoma, adenocarcinoma, malignant melanoma, and acute lymphocytic and myelocytic leukemia. They are also suited for anti-angiogenesis therapy and for the treatment of chronic inflammatory diseases (psoriasis, arthritis). To prevent uncontrolled cell growth on, and into or applied to polymeric materials. The compds. provided for in the invention can be used alone or, to achieve additive or synergistic effects, in combination with other principles and substance categories used in tumor therapy.

LI02 ANSWER 22 OF 24 CAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 1999:116659 CAPLUS Full-text
DOCUMENT NUMBER: 130:168164
TITLE: New (C1-C6)-fragments, method for their preparation and their application for synthesis of epothilone and epothilone derivatives

INVENTOR(S):
Klar, Ulrich; Schwede, Wolfgang;
Schuballa, Werner; Buchmann, Bernd;
Schirner, Michael
Schering A.-G., Germany
Ger. Offen., 18 pp.

PATENT ASSIGNEE(S):
SOURCE:

CODEN: GWXXBX

Patent

German

4



DOCUMENT TYPE:

FAMILY ACC. NUM. COUNT:

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 19735578	A1	19990211	DE 1997-19735578	19970809
CA 2299608	A1	19990218	CA 1998-2299608	19980810
WO 9907692	A2	19990218	WO 1998-EP5064	19980810
WO 9907692	A3	19990514		
W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DK, EE, ES, FI, GB, GE, GH, GM, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW				
RW: GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LJ, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
AU 9893409	A	19990301	AU 1998-93409	19980810
EP 1005465	A2	20000607	EP 1998-946309	19980810
EP 1005465	B1	20070725		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, CY, AL, MK				
JP 2000-306196	T	20010828	JP 2000-306196	19980810
AT 368036	T	20070815	AT 1998-946309	19980810
US 2003144523	A1	20030731	US 2000-485292	20000503
PRIORITY APPLN. INFO.:				
DE 1997-19735574 A 19970809				
DE 1997-19735575 A 19970809				
DE 1997-19735578 A 19970809				
DE 1997-19748928 A 19971024				
DE 1997-19749717 A 19971031				
DE 1997-19751200 A 19971113				
DE 1998-19813821 A 19980320				
WO 1998-EP5064 W 19980810				
CASREACT 130:168164; MARPAT 130:168164				

OTHER SOURCE(S):
GI



III

AB The title compds. [I; II; III; R₁, R₂ = H, alkyl, aryl, aralkyl; R₃ = CH₂OH, CH₂OR; R₄ = OH, OR; R = CR₇R₈; R₇, R₈ = H, alkyl, aryl, or R₇R₈ = (CH₂)_n; n = 2-6; R₅, R₆ = H, alkyl, aralkyl, or R₅R₆ = (CH₂)_m; m = 2-5; R₉, R₁₀ = H,

from which structure-activity- relationships can be deduced. Epothilone-A (R=H) Epothilone B (R=Me) Epothilone C (R=H) Epothilone D (R=Me).

L102 ANSWER 20 OF 24 CAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 2000:13195 CAPLUS Full-text
 DOCUMENT NUMBER: 132:64110
 TITLE: The preparation process, intermediate products and pharmaceutical use of epothilone derivatives
 INVENTOR(S): Buchmann, Bernd; Klar, Ulrich; Schuballa, Werner; Schwede, Wolfgang;
 Schirmer, Michael; Menrad, Andreas
 PATENT ASSIGNEE(S): Schering A.-G., Germany
 SOURCE: PCT Int. Appl., 86 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: German
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 200000485	A1	20000106	WO 1999-EP4915	19990630
W: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DK, EE, ES, FI, GB, GD, GE, GH, GM, GR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ZA, ZW, AM, AZ, BY, BG, CZ, MD, RU, TJ, TN				
RW: GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, BU, CF, CG, CI, CN, GA, AU 20000117				
DE 19830060	A1	20000210	DE 1998-19830060	19980630
DE 19923001	A1	20001116	DE 1999-19923001	19990513
AU 9950369	A	20000117	AU 1999-50369	19990630
PRIORITY APPLN. INFO.: DE 1998-19830060 A 19980630 DE 1999-19923001 A 19990513 WO 1999-EP4915 W 19990630				
OTHER SOURCE(S): CASREACT 132:64110; MARPAT 132:64110				

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB The invention relates to new epothilone derivs. I (R1a, R1b = H, Cl-10-alkyl, aryl, C7-10-alkyl; R1aR1b = (CH2)m, m = 2 - 5; R2a, R2b = H, Cl-10-alkyl, aryl, C7-10-alkyl; R2aR2b = (CH2)n, n = 2 - 5; R3 = H, Cl-10-alkyl, aryl, C7-10-alkyl; R4a, R4b = H, Cl-10-alkyl, aryl, C7-10-alkyl; R4aR4b = (CH2)m, m = 2 - 5; D-E = CH2CH2, CH-CH, C.tpbond.C, oxirane ring, CH(OH)CH(OH), CH(OH)CH2; R5 = Cl-10-alkyl, aryl, C7-10-alkyl; R6, R7 = H; R6R7 = O, bond; R8 = Cl-10-alkyl, aryl, C7-10-alkyl; R25 = H, Cl-10-alkyl, Cl-11-hydroxyalkyl, Cl-10-haloalkyl; X = O, (OR)2, C2-10-alkylene- α,ω -dioxo, CR11R12; CX = CH(OR)10; R9 = Cl-20-alkyl; R10 = H, protecting group; R11, R12 = H, Cl-10-alkyl, aryl, C7-10-alkyl; R11R12 = CH2, C5-7-carbocyclic ring; Y = O, CY = CH2; CZ = CH(OR)13, R13 = H, protecting group which are prepared via cyclization of ketones II (R15 = H, OH halogen, OR15a, OSO2R15b; R15a = H, SO2-alkyl, SO2-aryl, C7-20-alkyl, (CH2)o, CR16aR16b; R15b = H, Cl-20-alkyl, aryl, C7-20-alkyl; R16a, R16b = H, Cl-10-alkyl, aryl, C7-20-alkyl;

R16aR16b = (CH2)q; q = 2 - 4; q = 3 - 6). Thus, epothilone derivative III was prepared via macrolactonization of carboxylic acid IV with 2,4,6-trichlorobenzoyl chloride and Et3N in THF followed by deprotection with aqueous CF3CO2H in CH2Cl2. I cooperate with tubulin by stabilizing formed microtubuli.

REFERENCE COUNT: 10 THERE ARE 10 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L102 ANSWER 21 OF 24 CAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 1999:126888 CAPLUS Full-text
 DOCUMENT NUMBER: 130:196529
 TITLE: Preparation of new epothilone derivatives, as pharmaceutical agents
 INVENTOR(S): Klar, Ulrich; Schwede, Wolfgang; Schuballa, Werner; Buchmann, Bernd;
 Schirmer, Michael
 PATENT ASSIGNEE(S): Schering Aktiengesellschaft, Germany
 SOURCE: PCT Int. Appl., 185 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: German
 FAMILY ACC. NUM. COUNT: 4
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9907692	A2	19990218	WO 1998-EP5064	19980810
WO 9907692	A3	19990514		
W: AE, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DK, EE, ES, FI, GB, GE, GH, GM, GR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW				
RW: GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, BU, CF, CG, CI, CN, GA, AU 19990211				
DE 19735574	A1	19990211	DE 1997-19735574	19970809
DE 19735575	A1	19990211	DE 1997-19735575	19970809
DE 19735578	A1	19990211	DE 1997-19735578	19970809
DE 19748928	A1	19990429	DE 1997-19748928	19971024
DE 19749717	A1	19990506	DE 1997-19749717	19971031
DE 19751200	A1	19990520	DE 1997-19751200	19971113
DE 19813821	A1	19990923	DE 1998-19813821	19980320
CA 2299608	A1	19990218	CA 1998-2299608	19980810
AU 9893409	A	19990301	AU 1998-93409	19980810
EP 1005465	A2	20000607	EP 1998-946309	19980810
EP 1005465	B1	20070725		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, CY, AL, MK				
JP 2001512723	T	20010828	JP 2000-506196	19980810
ZA 9810403	A	20000515	ZA 1998-10403	19981113
IN 190805	A1	20030823	IN 1998-DE3413	19981116
US 2003144523	A1	20030731	US 2000-485292	20000503
IN 20020601305	A	20050311	IN 2002-DE1305	20021227
PRIORITY APPLN. INFO.: DE 1997-19735574 A 19970809 DE 1997-19735575 A 19970809 DE 1997-19735578 A 19970809 DE 1997-19748928 A 19971024 DE 1997-19749717 A 19971031 DE 1997-19751200 A 19971113				

stabilizing the microtubuli which are formed. They are able to influence the cell division phase-specifically and are suitable for treating malignant tumors such as cancers of the ovaries, stomach, colon, glands, breasts, lungs, head and neck, malignant melanoma and acute lymphocytic and myelocytic leukemia. These compounds are also suitable for anti-angiogenesis therapy and for treating chronic inflammatory diseases (psoriasis, arthritis) and can be deposited on or in polymer materials in order to prevent uncontrolled cell proliferation on medical implants and to improve the compatibility. These derivs. can be used alone or in combination with other principles and classes of substances that can be used in the therapy of tumors to achieve additive or synergistic effects.

L102 ANSWER 18 OF 24 CAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 2000:573798 CAPLUS Full-text
 DOCUMENT NUMBER: 133:177064

Preparation of epothilone derivatives useful as

pharmaceuticals

INVENTOR(S): **Klar, Ulrich; Skuballa, Werner; Buchmann, Bernd; Schwede, Wolfgang;**

Schirmer, Michael

Schering A.-G., Germany

PATENT ASSIGNEE(S): PCT Int. Appl., 141 pp.

SOURCE: CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: German

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2000047584	A2	20000817	WO 2000-EP1104	20000211
WO 2000047584	A3	20001228		
W: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, GR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW				
RW: GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
DE 19907480	A1	20000817	DE 1999-19907480	19990211
CA 2360952	A1	20000817	CA 2000-2360952	20000211
EP 1161430	A2	20011212	EP 2000-920433	20000211
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
BR 200008206	A	20020219	BR 2000-8206	20000211
HU 200105330	A2	20020529	HU 2001-5330	20000211
JP 2002536450	T	20021029	JP 2000-598504	20000211
EE 200100422	A	20021216	EE 2001-422	20000211
IN 2001MN00825	A	20070504	IN 2001-MN825	20010713
BG 105803	A	20020329	BG 2001-105803	20010809
NO 2001003900	A	20011011	NO 2001-3900	20010810
MX 2001PA08148	A	20030721	MX 2001-PA8148	20010810
ZA 2001007458	A	20021210	ZA 2001-7458	20010910
US 7001916	B1	20060221	US 2001-913163	20011207
US 2006040990	A1	20060223	US 2005-189787	20050727
PRIORITY APPLN. INFO.:				
			DE 1999-19907480	A 19990211
			DE 1999-19954229	A 19991104
			WO 2000-EP1104	W 20000211

OTHER SOURCE(S): MARPAT 133:177064 US 2001-913163 A3 20011207
 GI

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB Novel epothilone deriva. I (R4 = R5 = H, Cl-C10 alkyl, aryl, C7-C20 aralkyl; R6, R7 are each H, or together an addnl. bond or O; R8 = Me or H; R9a, R10b together = trimethylene; R2 = Ph, CH2Ph; X = 2-pyridyl, 2-methyl-4-thiazolyl, 2-methyl-4-oxazolyl; or R9a, R10b together = trimethylene; R2 = Me, Et, Pr; X = 2-pyridyl, 2-methyl-4-thiazolyl, 2-methyl-4-oxazolyl; or simultaneously R9a = R10b = Me; R2 = Me, Et, Pr; X = 2-pyridyl, 2-methyl-4-thiazolyl or 2-methyl-4-oxazolyl; and the N and/or S atoms in X can be in an oxidized form; and if R2 and R8 = Me, X can only be a 2-pyridyl residue which is optionally oxidized at the nitrogen atom) and all possible stereoisomers and their mixts were prepared. Thus II was prepared in a multistep sequence from the starting materials III and IV. The novel compds. interact with tubulin by stabilizing the formed microtubuli. The compds. are able to influence the cell division in a phase-specific manner and are suited for treating malignant tumors, for example, ovarian cancer, gastric carcinoma, colon cancer, breast cancer, lung cancer, head and neck cancer, malignant melanoma, and acute lymphocytic and myelocytic leukemia. The inventive compds. are suited for use in anti-angiogenic therapy as well as for treating chronic inflammatory diseases (psoriasis, arthritis). In order to prevent uncontrolled cell proliferations and to improve the compatibility of medical implants, the inventive compds. can be applied or incorporated in polymeric materials. The inventive compds. can be used alone or, in order to achieve additive or synergistic effects, in conjunction with addnl. constituents and substance classes which can be used in tumor therapy.

L102 ANSWER 19 OF 24 CAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 2000:332415 CAPLUS Full-text
 TITLE: Epothilones.

AUTHOR(S): **Klar, Ulrich; Skuballa, Werner; Schwede, Wolfgang; Buchmann, Bernd**

CORPORATE SOURCE: Preclinical Drug Research, Schering AG, Germany, Berlin, D-13342, Germany

SOURCE: Book of Abstracts, 219th ACS National Meeting, San Francisco, CA, March 26-30, 2000 (2000), ORGN-288. American Chemical Society: Washington, D. C.

CODEN: 69CLAC

DOCUMENT TYPE: Conference: Meeting Abstract

LANGUAGE: English

AB The new natural product class of epothilones seems to parallel the biol. behavior of paclitaxel regarding its action on the tubulin system although the chemical structures are quite different. Despite its impressive anti-proliferative effects also against multi drug resistant cell lines epothilone B shows severe toxicity at therapeutic relevant doses in vivo. Thus the need for epothilone analogs with improved properties is obvious. In contrast to paclitaxel structural modifications can be achieved more easily by total synthesis. Therefore we have developed a highly convergent total synthesis which allows an efficient preparation of large amts. of strategic important building blocks with high optical purity, high flexibility regarding structural modifications in most parts of the mol., efficient syntheses of analogs yielding sufficient amts. for in vivo characterization. Using this methodol., more than 250 analogs of epothilone B and D have been synthesized

W: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DK, DM, EE, ES, FI, FR, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KH, KZ, LC, LK, LR, LS, LT, LV, MA, MD, ME, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, AM, AZ, BY, BG, BR, BU, BT, TJ, TM, SZ, TZ, ZW, AT, BE, CH, CY, DE, RW: GH, GM, KE, LS, MW, SD, SI, SZ, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, CA, GN, GW, ML, MR, NE, SN, TD, TG

DE 199008763 A1 20000824 DE 1999-19908763 A 19990218
PRIORITY APPL. INFO.: MARPAT 133:193027 19990218
OTHER SOURCE(S): DE 1999-19908763 A 19990218

↑ STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT ↑


Epothilone derivs. I (R1a, R1b = H, Cl-C10-alkyl, aryl, C7-C20-aralkyl; (CH2)_n m = 1-5; CH2OR2, R2b = H, Cl-C10-alkyl, aryl, C7-C20-aralkyl; (CH2)_n n = 2-5; E or B where t = 1-2, w = 1-2; G, Gl = H, halogen, CN, R24, Cl-C20-acyl, Cl-C20-acyloxy, OR24, CO2R24, N3, NO2, NR24aR24b; R24a, R24b = R24, (CH2)_e e = 4-6; R24 = R3a = H, Cl-C10-alkyl, aryl, C7-C20-aralkyl; R14 = H, OR14a, halogen; R3b = OPG14; R3b, R4a = bond; R4a, R4b = H, F, Cl-C10-alkyl, aryl, C7-C20-aralkyl; R5 = H, Cl-C10-alkyl, aryl, C7-C20-aralkyl; (CH2)_s s-A, where s = 1-4, A = OR22, halogen; R22 = H, protecting group; R6, R7 = H, bond, OR8 = H, F, Cl-C10-alkyl, aryl, C7-C20-aralkyl; X = O, two alkoxy groups OR23, C2-C10-alkylene-α,ω-dihydroxy group straight or branched, H/OR9, CR10R11 where R23 = Cl-C20-alkyl; R9 = H, protecting group; R10, R11 = H Cl-C10-alkyl, aryl, C7-C20-aralkyl or together are a 5-7 membered carbocyclic ring; Y = O or 2 H atoms; Z = O, H/OR12 where R12 = O, protecting group) were prepared in addition to all possible stereoisomers and mixts. Thus II was prepared from 1,3-bis(hydroxymethyl)benzene in a multistep synthesis. These epothilone derivs. interact with tubulin by stabilizing the formed microtubule. The compds. are able to influence the cell division in a phase-specific manner and are suited for treating malignant tumors, for example, ovarian cancer, gastric carcinoma, colon cancer, breast cancer, lung cancer, head and neck cancer, malignant melanoma, and acute lymphocytic and myelocytic leukemia. These derivs. are suited for use in anti-angiogenic therapy as well as for treating chronic inflammatory diseases (psoriasis, arthritis). These compds. can be applied or incorporated in polymeric materials to prevent uncontrolled cell proliferations and to improve the compatibility of medical implants. They can be used alone or in conjunction with adnl. constituents and substance classes to achieve additive or synergistic effects in tumor therapy.

1102 ANSWER 17 OF 24 CAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 2000:592719 CAPLUS Full-text
DOCUMENT NUMBER: 133:193025
TITLE: Preparation of new epothilone d

pharmaceutical uses

INVENTOR(S): Kiaz, Ulrich; Schwede, Wolfgang;
Krusalla, Werner; Buchmann, Bernd;
Schirmer, Michael; Menrad, Andreas
PATENT ASSIGNEE(S): Schering A.-G., Germany
PCT Int. Appl., 54 pp.
CODEN: PIXXD2
DOCUMENT TYPE: Patent

19

[illegible]

Epothilone derivs. I (R1a, R1b = H, Cl-C10 alkyl, aryl; C7-C20 alkyl; or together are (CH₂)_m m = 1-5; or CH₂OH; R2a, R2b = H, Cl-C10 alkyl, aryl; C7-C20 alkyl; or together are (CH₂)_n n = 2-5; G1-G-E-E1 = CR₃AR₃b-CR₄C-CH₂; CR₃AR₃b-CD(T)R₄-CHD(T)-CH₂ (2,3-epoxy)-CR₃AR₃b-CR₄OC-CH₂; CR₃AR₃b-COH(H)R₄-CHO(H)-CH₂; CR₃AR₃b-CR₄-CH=CH where R₃a = H, Cl-C10 alkyl, aryl; C7-C20 alkyl; R14 = H, OR14a, halogen, OSO₂OR14b; R3b = OPG14 or R3b, R14a = bond; R₄ = H, Cl-C10 alkyl, aryl; C7-C20 alkyl; R5 = H, Cl-C10 alkyl, aryl; C7-C20 alkyl, (CH₂)₅-A s = 1-4, A = OR22 or halogen; R22 = H or protecting group; R6, R7 = H, O, bond; R8 = H, Cl-C10 alkyl, aryl; C7-C20 alkyl; X = O, OR23, C7-C10-alkylene- α,ω -dihydroxy which can be a straight chain or branched; H/OR9 or the group CR10R11 where R23 = Cl-C20 alkyl; R9 = H or a protecting group; R10, R11 = H, Cl-C20 alkyl, aryl; C7-C20 alkyl or R10, R11 together form a 5-7 membered carbocyclic ring; Y = O or 2 H atoms; Z = O or H/OR12 where R12 = H or a protecting group) were prepared in addition to all possible stereoisomers and mixts. Thus II was prepared from (4)-1-acetoxypentad-4-one in a multistep synthesis. These epothilone derivs. interact with tubulin by

IN 2001M01305 A 20070504 IN 2001-MN1305 20011019
 BG 106053 A 20020531 BG 2001-106053 20011026
 NO 2001005278 A 20011221 NO 2001-5278 20011029
 KX 2001PA11039 A 20030630 MX 2001-PA11039 20011030
 ZA 2001009859 A 20030228 ZA 2001-9859 20011129
 B1 7125993 B1 20061024 US 2002-979939 20020606
 US 2005113429 A1 20050526 US 2004-965802 20041018
 IN 2005MN00837 A 20070608 IN 2005-MN837 20050802
 A1 2006046997 A1 20060302 US 2005-214998 20050831
 JP 2007224038 A 20070906 JP 2007-104224 20070411
 DE 1999-19921086 A 19990430
 DE 1999-19954228 A 19991104
 DE 2000-10013363 A 20000309
 DE 2000-10015836 A 20000327
 JP 2000-615619 A3 20000501
 WO 2000-18657 W 20000501
 IN 2001-MN1305 A3 20011019
 US 2002-979939 A3 20020606

OTHER SOURCE(S):

AB The title compds. were prepared by various combinations of 3 fragments making up the mols. Thus, [4S,7R,8S,9S,13Z,16S(E)]-4,8-dihydroxy-16-[1-methyl-2-(2-pyridyl)ethenyl]-1-oxa-5,9,13-tetramethyl-7-(3-butynyl)-13-cyclohexadecene-2,6-dione was prepared in several steps starting from (4S)-4-(2-methyl-1-oxo-2-propyl)-2,2-dimethyl[1,3]dioxane and 5-(trimethylsilyl)-4-pentynylmagnesium bromide.

LI02 ANSWER 14 OF 24 CAPIUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2000:738730 CAPIUS Full-text

DOCUMENT NUMBER: 133:309795

TITLE: Preparation of new epothilone derivatives and their pharmaceutical uses

INVENTOR(S): **Klar, Ulrich; Schwede, Wolfgang;**

Struballe, Werner; Buchmann, Bernd;

Schirmer, Michael

Schering A.-G., Germany

Ger. Offen., 74 pp.

CODEN: GWXXBX

PATENT ASSIGNEE(S): Patent

DOCUMENT TYPE: German

LANGUAGE: 1

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:


PATENT NO. KIND DATE APPLICATION NO. DATE

DE 19908767 A1 20001019 DE 1999-19908767 19990218

PRIORITY APPLN. INFO.: DE 1999-19908767 19990218

OTHER SOURCE(S): MARPAT 133:309795

GI



AB

New epothilone derivs. I (R1e, R1b = R2a, R2b = same or different H, alkyl, aryl, aralkyl or (CH2)m,n m, n = 2-5; R3 = H, alkyl, aryl, aralkyl; R4a, R4b = same or different H, alkyl, aralkyl or (CH2)p = 2-5, CH2CH2, CH=CH, C.tp.bond.C, epoxy, CH(OH)CH(OH), CH(OH)CH2; D-E = a group; R5 = H, alkyl, aryl, aralkyl; R6, R7 = H, bond, O; R8 = H, alkyl, aryl, aralkyl; X = O, OR23 alkylene- α , ω -diol group straight or branched, OR9 or the C10R11 group where R23 = alkyl, R9 = H or protecting group and R10, R11 = same or different H, alkyl, aryl, aralkyl or R10, R11 = together with methylene are a 5-7 membered carbocyclic ring; Y = O or two H; Z = O or H/OR12 and R12 = H or a protecting group) were prepared. Thus E- and Z-II were prepared via a multistep synthesis. I cooperate with tubulin by stabilizing formed microtubuli. I are able phase specifically to affect the cell division and are suitable for the treatment of malignant ovarian, stomach, colon, adeno, breast, lung, head and neck tumors, malignant melanomas, acute lymphocytic and myelocytic leukemia. Derivs. of I are suitable for use in anti-angiogenic therapy as well as for treating chronic inflammatory diseases (psoriasis, arthritis). In order to prevent uncontrolled cell proliferations and to improve the compatibility of medical implants I can be applied or incorporated into polymeric materials. I can be used alone or to achieve additive or synergistic effects in combination with further principles and substance classes applicable in tumor therapy.

LI02 ANSWER 15 OF 24 CAPIUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2000:592721 CAPIUS Full-text

DOCUMENT NUMBER: 133:193028

TITLE: Preparation of 16-halogen epothilone derivatives and their use as antitumor agents

INVENTOR(S): **Klar, Ulrich; Struballe, Werner;**

Buchmann, Bernd; Schwede, Wolfgang;

Schirmer, Michael

Schering Aktiengesellschaft, Germany

PCT Int. Appl., 105 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: German

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

FAMILY ACC. NUM. COUNT: 3

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2000066589	A1	20001109	WO 2000-1B657	20000501
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GR, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, ME, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW				
RW: GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GN, GW, ML, MR, NE, SN, TD, TG				
DE 19921086	A1	20001102	DE 1999-19921086	19990430
DE 19954228	A1	20010913	DE 1999-19954228	19991104
DE 10015836	A1	20011011	DE 2000-10015836	20000327
CA 2371226	A1	20001109	CA 2000-2371226	20000501
BR 2000010190	A	20020108	BR 2000-10190	20000501
EP 1173441	A1	20020123	EP 2000-922826	20000501
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, MC, PT, IE, SI, LT, LV, FI, RO				
JP 2002543203	T	20021217	JP 2000-615619	20000501
EE 200100568	A	20030217	EE 2001-568	20000501
NZ 514989	A	20040227	NZ 2000-514989	20000501
IN 7072750	B2	20040506	AU 2000-43103	20000501
IN 2001MN01305	A	20070504	IN 2001-MN1305	20011019
BG 106053	A	20020531	BG 2001-106053	20011026
NO 2001005278	A	20011221	NO 2001-5278	20011029
MX 2001PA1039	A	20030630	MX 2001-PA11039	20011030
US 7125953	B1	20061024	US 2002-979939	20020606
IN 2005MN00837	A	20070608	IN 2005-MN837	20050802
US 2006046997	A1	20060302	US 2005-214988	20050831
PRIORITY APPL. INFO.:			DE 1999-19921086	A1 19990430
			DE 1995-19954228	A1 19991104
			DE 2000-10015836	A1 20000327
			DE 2000-10013863	A1 20000309
			WO 2000-1B657	W 20000501
			IN 2001-MN1305	A3 20011019
			US 2002-979939	A3 20020606

OTHER SOURCE(S):

MARPAT 133:362656

GI

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB The antitumor agents, 6-alkenyl-, 6-alkynyl-, 6-alkynyl- and 6-epoxyepothilones I (R_{1a}, R_{1b} are same or different = H, C1-C10 alkyl, C6-C12 aryl, C7-C20 aralkyl each optionally substituted; or together = (CH₂)_m m = 1-5 or -CH₂CH₂-; R_{2a}(R_{2b} replace a with b) = H, substituted alkyl, aryl, aralkyl, (CH₂)_nra-C₁tpbond (or =)C-(CH₂)_npa-R_{2a}, Q, Q1 where n = 0-5; R_{3a} = H, substituted alkyl, aryl 4; pa, pb = the same or different and = 0-3; R_{3b} = H, substituted alkyl, aryl or alkyl; R_{3b} = OH, OPG14; R₄ = H, OR14a, halogen and R_{4a} = H, SO₂-alkyl, SO₂-aryl or SO₂-aralkyl; R₄ = H, substituted alkyl, aryl or alkyl, halogen, OR25, CN; R_{26a}, R_{26b} = same or different = H, substituted alkyl, aryl or aralkyl, C1-C10 acyl or if pa or pb > 0, addnl. a group OR27; R₂₅ = R₂₇ = R₂₂ = H, PG; R₅ = H, substituted alkyl, aryl or alkyl, (CH₂)_s s = 1-4, T = OR22 or halogen; R₆, R₇ = H or together = bond or O; G = X-CR8 or bi- or tricyclic

13

aryl radical and R₈ = H, halogen, CN, or substituted alkyl, aryl or aralkyl; X = O, two OR23 groups, C2-C10-alkylene- α,ω -dioxy straight chain or branched; H/OR9 or CR10R11 group and R₂₃ = alkyl radical, R₉ = H, PG, R10, R11 = same or different = H, substituted alkyl, aryl or alkyl, or together with the methylene are a 5-7 carbocyclic ring; D-E = CH₂CH₂ or OCH₂; A = OC(O), OCH₂, CH₂C(O), NR₂SO₂ and R₂₉ = H, alkyl; Z = O or H/OR12 and R12 = H, PG) were prepared. Thus II was prepared in a multistep synthesis starting from (4S)-4-(2-methyl-1-oxoprop-2-yl)-2,2-dimethyl(1,3)dioxane and 5-trimethylsilylpent-4-in-1-yl magnesium bromide. II had an IC₅₀ value [nM] of 3.0 for the growth inhibition of human MCF-7 breast- and 75 for multidrug resistant NCI/ADR carcinoma cell lines with a selectivity of 2.5. The new epothilone derivs. interact with tubulin by stabilizing microtubuli that are formed. They are able to influence the cell-splitting in a phase-specific manner and are therefore useful in treating diseases or conditions associated with the need for cell growth, division and/or proliferation. Thus the epothilone derivs. are suitable for treating malignant tumors, e.g., ovarian, stomach, colon, adenocarcinoma, breast, lung, head and neck carcinomas, malignant melanoma, acute lymphocytic and myelocytic leukemia; and for anti-angiogenesis therapy as well as for treatment of chronic inflammatory diseases (such as psoriasis, arthritis).

REFERENCE COUNT: 4

THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L102 ANSWER 13 OF 24 CAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 2000:772379 CAPLUS Full-text
DOCUMENT NUMBER: 133:321769

TITLE: 6-Alkenyl and 6-alkynyl derivatives of epothilone
INVENTOR(S): Klier, Ulrich; Schwede, Wolfgang;

Strubelle, Werner; Buchmann, Bernd;

Hoffmann, Jens; Lichtner, Rosemarie

Schering A.-G., Germany

Ger. Offen., 18 pp.

PATENT ASSIGNEE(S): CODENT: GWXBX

DOCUMENT TYPE: Patent

LANGUAGE: German

FAMILY ACC. NUM. COUNT: 3

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 19921086	A1	20001102	DE 1999-19921086	19990430
CA 2371226	A1	20001109	CA 2000-2371226	20000501
WO 2000066589	A1	20001109	WO 2000-1B657	20000501
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GR, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, ME, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW				
RW: GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GN, GW, ML, MR, NE, SN, TD, TG				
BR 2000010190	A	20020108	BR 2000-10190	20000501
EP 1173441	A1	20020123	EP 2000-922826	20000501
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, MC, PT, IE, SI, LT, LV, FI, RO				
HU 200201010	A2	20020828	HU 2002-1010	20000501
JP 2002543203	T	20021217	JP 2000-615619	20000501
EE 200100568	A	20030217	EE 2001-568	20000501
NZ 514989	A	20040227	NZ 2000-514989	20000501
AU 772750	B2	20040506	AU 2000-43103	20000501

14

TITLE: Subcellular distribution of epothilones in human tumor cells

AUTHOR(S): Lichtner, R. B.; Rotgeri, A.; Bunte, T.;

Ruchmann, B.; Hoffmann, J.; **Schwede, W.;** **Skuballa, W.;** **Klar, U.**

CORPORATE SOURCE: Research Laboratories of Schering AG, Berlin, 13342, Germany

SOURCE: Proceedings of the National Academy of Sciences of the United States of America (2001), 98(20), 11743-11748

CODEN: FNASAG; ISSN: 0027-8424

PUBLISHER: National Academy of Sciences

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Epothilones are a new class of natural and potent antineoplastic agents that stabilize microtubules. Although 12,13-epoxide derivs. are potent antiproliferative agents, the activities of the corresponding 12,13-olefin analogs are significantly decreased. These data were confirmed for two new analogs, 6-propyl-EpoB (pEB) and 6-propyl-EpoD (pED), in comparison with the natural compds. EpoB/EpoD, by using human A431, MCF7, and MDRI-overexpressing NCI/Adr cells. By using tritiated pEB/pED, compound uptake, release, and nuclear accumulation were investigated in A431 and NCI/Adr cells. In these cells, epothilones can principally be recognized and exported by verapamil-sensitive efflux pumps, which are not identical to MDRI. The degree of export depends on the structure, olefin vs. epoxide-analog, and also on the intracellular drug concentration. The accumulation of pED used at 3.5 or 70 nM, resp., was increased in the presence of 10 μ M Verapamil in both cell lines 2- to 8-fold. In contrast, the intracellular levels of pEB were affected by Verapamil only at 3.5 nM pEB in NCI/Adr (2-fold) and not in A431 cells. In addition, strong nuclear accumulation was observed for pEB (40-50%) but not paclitaxel or pED (5-15%) in both cell lines. Our study suggests that differences in growth inhibitory efficacy between epoxide and olefin analogs may be based on different mechanisms of drug accumulation and subcellular distribution.

REFERENCE COUNT:

34 THERE ARE 34 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L102 ANSWER 11 OF 24 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2001:676638 CAPLUS Full-text

DOCUMENT NUMBER: 135:236394

TITLE: Synthesis of radioactively labeled epothilone derivatives and their biochemical and pharmaceutical usage

INVENTOR(S):

Klar, Ulrich; Cav, Juergen; **Skuballa, Ruchmann, Bernd;** Bunte, Thomas; Lichtner, Rosemarie

Schering Aktiengesellschaft, Germany

PCT Int. Appl., 31 pp.

CODEN: PIXXD2

Patent

German

PATENT ASSIGNEE(S):

SOURCE:

DOCUMENT TYPE:

LANGUAGE:

FAMILY ACC. NUM. COUNT: 3

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001066154	A2	20010913	WO 2001-EP2699	20010309
W:	AE, AG, AL, AM, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CN, CR, CU, CZ, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU,			

LV, MA, MD, MG, MK, MM, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW

RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TG, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GN, GW, ML, MR, NE, SN, TD, TG

PRIORITY APPIN. INFO.:


DE 2000-10013363

A 20000309

OTHER SOURCE(S):

GI

MARPAT 135:236394



AB The invention relates to novel radioactively labeled pharmacol. effective epothilone derivs. of general formula (I), where R1 represents O-PG and hydroxyl, where PG is a protective group; R2a, R2b are the same or different and represent, independent of one another, hydrogen C1-C10 alkyl, aryl, C7-C20 alkyl or, together, represent a (CH2)m group, where m is equal to 1, 2, 3, 4 or 5; R3 represents a C2-C10 alkyl group, a C2-C10 alkenyl group or a C8-C20 aralkyl each containing 2n tritium atoms, where n equals 1 or 2; R4 represents O-PG and hydroxyl; R5 represents hydrogen C1-C10 alkyl, aryl, C7-C20 aralkyl and halogen; W-2 represents a CH2-CH2, CH2-O or O-CH2 group; R6 represents hydrogen, C1-C10 alkyl, aryl, C7-C20 aralkyl, (CH2)s-V and halogen, where s equals 1, 2, 3 or 4 and V represents O-PG, hydroxyl or halogen; R7, R8 each represent a hydrogen atom and, together, represent an addnl. bond or an oxygen atom; A represents aryl, C7-C20 aralkyl, and a group R10-CH-C9-, where R9 represents hydrogen, halogen, CN, C1-C20 alkyl, aryl, and C7-C20 aralkyl, and R10 represents hydrogen, C1-C20 alkyl-, aryl-, C7-C20 aralkyl, and; X-Y represents an O-C(=O), an O-CH2, a CH2-C(=O), an NR11-C(=O) and an NR11-SO2 group, wherein R11 represents hydrogen and C1-C10 alkyl. The novel compds. of formula I are valuable pharmaceuticals and valuable diagnostic probes for elucidating, for example, active mechanisms and biochem., pharmacokinetic and/or pharmacodynamic processes.

L102 ANSWER 12 OF 24 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2000:790507 CAPLUS Full-text

DOCUMENT NUMBER: 133:362656

TITLE: Preparation of 6-alkenyl-, 6-alkynyl- and 6-epoxyepothilone derivatives and their antitumor activity

INVENTOR(S):

Klar, Ulrich; **Schwede, Wolfgang;**

Skuballa, Werner; **Ruchmann, Bernd;**

Hoffmann, Jens; Lichtner, Rosemarie

Schering Aktiengesellschaft, Germany

PCT Int. Appl., 298 pp.

CODEN: PIXXD2

Patent


English

PATENT ASSIGNEE(S):

SOURCE:

DOCUMENT TYPE:

LANGUAGE:



II

Oxa-epothilones, such as I [R3 = heteroaryl, heteroaryalkenyl, heteroarylhaloalkenyl, etc.; R8, R8a = H, alkyl, arylalkyl; R8R8a = alkylene, heteroalkylene; R10 = H, alkyl, alkenyl, alkynyl; R1R1a = bond, O; R16 = H, CN, alkyl, halogen; X = O, NH], were prepared for a variety of therapeutic uses, such as treatment of malignant tumors, proliferative diseases, leukemia, and chronic inflammatory diseases, as well as for anti-angiogenic SE therapy. Thus, oxa-epothilone II was prepared via a multistep synthetic sequence starting from (R)-1,2-propanediol, and [(3S,4Z)-3-[(1*E*,1*E*-dimethylethynyl)dimethylsilyloxy]-4-fluoro-5-(2-methyl-4-thiazolyl)-4-pentenyl]triphenylphosphonium iodide. Pharmaceutical formulations of the prepared oxa-epothilones were discussed, but specific biol. activity data was not presented.

LL102 ANSWER 9 OF 24 CAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 2001:780370 CAPLUS Full-text

ACCESSION NUMBER:

DOCUMENT NUMBER:

DOCUMENT NUMBER: 133:331294
TITLE: Preparation of epothilone derivatives for

INVENTOR(S):
Buchmann, Bernd; Klar, Ulrich;

Hoffmann. Jens: Lichtner Rosemarie, Skuballa, Wernier, Schwesig, Möllgang.

PATENT ASSIGNEE(S): Schering A.-G., Ger

SOURCE: Ger. Offen., 4

CODEN:

DOCUMENT TYPE: Patent

LANGUAGE:	NTM COUNT:
FAMILY ACC	

PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 10020517	A1	200111025	DE 2000-10020517	200000419
WO 2001081342	A2	20011101	WO 2001-EP4552	200100419

9

SN 10/563058 Page 10 of 172 STIC STN SEARCH RESULTS

WO 2001081342	A3	20020510
---------------	----	----------

W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CH, CY, CZ, DK, DR, DZ, EE, ES, FI, GB, GD, GE, GH, GR, GU, HR, HU, ID, IL, IN, IS, JP, KE, KP, KR, LC, LC, LK, LR, LT, LU, LV, MA, MD, MG, MK, MM, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VU, YU, ZW, ZA, ZW, AM, AZ, BY, GB, KZ, MD, RU, TJ, TM	
RW:	GH, GA, KE, LS, MA, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, CZ, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GU, ML, MR, NE, SN, TD, TG	
EP 126740	A2	EP 20030122
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR	20010419
JP 2003531207	T	JP 2001-578432
NO 2002005029	A	NO 2002-5029
US 2004058969	A1	US 2002-257925
PRIORITY APPLN. INFO.:		DE 2000-10020517
		WO 2001-EP4552
		W 20010419




II

Epothilones, such as I [R3 = heteroaryl, heteroarylalkenyl, heteroarylhaloalkyl, etc.]; R8, R9a = H, alkyl, arylalkyl; R8R9a = alkylene, heteroalkylene; R10 = H, alkyl, alkynyl, alkynyl; R1R16a = bond, O; R16 = H, CN, alkyl, halogen; X = O, NH; X1 = O, CH2], were prepared for a variety of therapeutic uses, such as treatment of malignant tumors, proliferative diseases, leukemia, and chronic inflammatory diseases. Thus, epothilone II was prepared via a multistep synthetic sequence starting from (3S)-dihydro-3-hydroxy-4,4-dimethyl-2(3H)-furanone, L-(-)-malic acid, and [(2-methyl-4-thiazolylmethyl)phosphonic acid di-Et ester. Pharmaceutical formulations of the prepared oxo-epothilones were discussed, but specific biol. activity data was not presented.

LL102 ANSWER 10 OF 24 CAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 2001:729040 CAPLUS Full-text
DOCUMENT NUMBER: 136:95676

Bernsd; Schwede, Wolfgang;
Skuballa, Werner
 Schering Aktiengesellschaft, Germany
 PCT Int. Appl., 30 pp.
 CODEN: PIXXD2
 Patent
 German
 1
 DOCUMENT TYPE:
 LANGUAGE:
 FAMILY ACC. NUM. COUNT:
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003053949	A1	20030703	WO 2002-EP14758	20021223
W: AE, AG, AL, AM, AT, AU, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DK, DM, DZ, EC, EE, ES, FI, GE, GD, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
DE 10164592	A1	20030703	DE 2001-10164592	20011221
AU 2002356783	A1	20030709	AU 2002-356783	20021223
US 2003176710	A1	20030918	US 2002-326263	20021223
PRIORITY APPL. INFO.: DE 2001-10164592 A 20011221 WO 2002-EP14758 W 20021223				
OTHER SOURCE(S): CASREACT 139:85166; MARPAT 139:85166				



AB The invention relates to C1-C6 fragments I [R1a, R1b = H, C1-10-alkyl, aryl, C7-20-alkyl, (CH2)m; m = 2 - 5; R2a, R2b = H, C1-10-alkyl, C1-10-alkenyl, C1-10-alkynyl, C7-20-alkyl, (CH2)n; n = 2 - 5; R15a, R15b = H, C1-10-alkyl, aryl, C7-20-alkyl, (CH2)q; q = 3 - 6] of epothilones and to an efficient method for producing such fragments and the deriva. thereof. Thus, (4S)-4-(2-methyl-3-oxohept-6-en-2-yl)-2,2-dimethyl-1,3-dioxane [I; R1a = Me, R2a = CH2CH:CH2, R2b = H, R15a = R15b = Me] was prepared from (3S)-1-hydroxy-2,2-dimethyl-3-(tetrahydropyranyloxy)-4-pentene, (S)-HOCH2CMe2CH(OTHP)CH:CH2, via O-benzoylation with PhCH2Br, hydroboration with BH3-THF complex, dehydratetetrahydropyranylation- isopropylidenation with Me2C(OMe)2 in MeCOMe containing catalytic tosyl acid, hydrogenolytic debenzoylation, Swern oxidation, Crignard reaction with MeMgBr, oxidn, with TPAT in CH2Cl2 contg, N-methylmorpholine N-oxide and alkylation with allyl bromide.
 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS
 REFERENCE COUNT: 3
 RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L102 ANSWER 7 OF 24 CAPIUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 20021132142 CAPIUS Full-text
 DOCUMENT NUMBER: 136:309773
 TITLE:
 AUTHOR(S):
 Synthesis and biological activity of epothilones
Klar, Ulrich; Skuballa, Werner;
Buchmann, Bernd; Schwede, Wolfgang;
 Bunte, Thomas; Hoffmann, Jens; Lichtner, Rosemarie B.
 Research Laboratories of Schering AG, Berlin, D-13342,
 Germany
 SOURCE: ACS Symposium Series (2001), 796(Anticancer Agents),
 131-147
 CODEN: ACSMC8; ISSN: 0097-6156
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal; General Review
 LANGUAGE: English
 AB A review. The total synthesis and biol. activity of epothilone analogs are described. Selected SAR data indicate the possibility to improve activity and selectivity by structural modifications. The new comds. may help to elucidate the therapeutic potential of this class of anticancer drugs.
 REFERENCE COUNT: 16
 THERE ARE 16 CITED REFERENCES AVAILABLE FOR THIS
 RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L102 ANSWER 8 OF 24 CAPIUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 2001780372 CAPIUS Full-text
 DOCUMENT NUMBER: 135:331295
 TITLE:
 INVENTOR(S):
 Preparation of oxa-epothilone derivatives for pharmaceutical use in the treatment of cancer
Schwede, Wolfgang; Klar, Ulrich;
Skuballa, Werner; Buchmann, Bernd;
 Hoffmann, Jens; Lichtner, Rosemarie
 Schering A.-G., Germany
 Ger. Offen., 46 pp.
 CODEN: GWXXBX
 Patent
 German
 1
 DOCUMENT TYPE:
 LANGUAGE:
 FAMILY ACC. NUM. COUNT:
 PATENT INFORMATION:

PATENT ASSIGNEE(S):
 SOURCE:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 10020899	A1	20011025	DE 2000-10020899	20000420
WO 2001081341	A2	20011101	WO 2001-EP4351	20010419
WO 2001081341	A3	20020425		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DK, DM, DZ, EE, ES, FI, GE, GD, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AE				

(DOWNLOADING INTERRUPTED)
 THE REST OF THIS BIB INFORMATION WAS NOT AVAILABLE AT THE TIME OF PRINTING

active ingredients II (AK = OC(O), OCH₂, CH₂CI(O), NR29C(O), NR29S(O), R29 = H, Cl-6-alkyl) according to known methods. The invention also relates to the corresponding Cl-C12 fragments.

REFERENCE COUNT:

1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L102 ANSWER 4 OF 24 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2003:719306 CAPLUS Full-text

DOCUMENT NUMBER: 139:240340

TITLE: Use of epothilones in the treatment of brain diseases

associated with proliferative processes

Lichtner, Rosemarie; Rotgeri, Andrea; **Klar**,

Ulrich; Hoffmann, Jens; **Buchmann**, **Bernd**

; **Schwede**, **Hilfgang**; **Stuballa**, **Haznar**

Schering A.-G., Germany

PCT Int. Appl., 53 pp.

CODEN: PIXD2

Patent

English

2

FAMILY ACC. NUM. COUNT:

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003074053	A1	20030912	WO 2003-EP2085	20030228
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GE, GR, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MX, MY, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, BG, CH, CY, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR, BF, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GN, GW, ML, MR, NE, SN, TD, TG				
EP 1340498	A1	20030903	EP 2002-4745	20030228
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR				
CA 2477403	A1	20030912	CA 2003-2477403	20030228
EP 1480643	A1	20030916	EP 2003-743360	20030228
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
BR 2003008154	A	20050104	BR 2003-8154	20030228
JP 2005525360	T	20050825	JP 2003-572570	20030228
MX 2004PA08450	A	20050713	MX 2004-PAB450	20040901
NO 2004004175	A	20041201	NO 2004-4175	20040930
ZA 2004007905	A	20060426	ZA 2004-4175	20040930
EP 2002-4745	US	2002-361062P	EP 2002-4745	20020301
WO 2003-EP2085	WO	2003-EP2085	WO 2003-EP2085	20030228

OTHER SOURCE(S):

AB The invention provides the use of an Epothilone, which shows an average distribution coefficient between plasma and brain of 0.3 to 1.5 in the mouse i.v. bolus injection assay, for the preparation of a medicament for the treatment of a brain disease associated with proliferative processes.

REFERENCE COUNT:

10 THERE ARE 10 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L102 ANSWER 5 OF 24 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2003:693140 CAPLUS Full-text

DOCUMENT NUMBER: 139:191465

TITLE: Use of epothilones in the treatment of brain diseases

associated with proliferative processes

Lichtner, Rosemarie; Rotgeri, Andrea; **Buchmann**,

Bernd; Hoffmann, Karin; **Klar**, **Ulrich**;

Schwede, **Hilfgang**; **Stuballa**, **Haznar**

Schering Aktiengesellschaft, Germany

Eur. Pat. Appl., 27 pp.

CODEN: EPXXDW

Patent

English

2

FAMILY ACC. NUM. COUNT:

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 1340498	A1	20030903	EP 2002-4745	20030301
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR				
CA 2477403	A1	20030912	CA 2003-2477403	20030228
WO 2003074053	A1	20030912	WO 2003-EP2085	20030228
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GE, GR, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, BG, CH, CY, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR, BF, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GN, GW, ML, MR, NE, SN, TD, TG				
AU 2003215618	A1	20030916	AU 2003-215618	20030228
US 2004019088	A1	20040129	US 2003-375043	20030228
EP 1480643	A1	20041201	EP 2003-743360	20030228
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
BR 2003008154	A	20050104	BR 2003-8154	20030228
JP 2005525360	T	20050825	JP 2003-572570	20030228
MX 2004PA08450	A	20050713	MX 2004-PAB450	20040901
NO 2004004175	A	20041201	NO 2004-4175	20040930
ZA 2004007905	A	20060426	ZA 2004-4175	20040930
EP 2002-4745	US	2002-361062P	EP 2002-4745	20020301
WO 2003-EP2085	WO	2003-EP2085	WO 2003-EP2085	20030228

OTHER SOURCE(S):

AB The invention provides the use of an epothilone, which shows an average distribution coefficient between plasma and brain of 0.3-1.5 in the mouse i.v. bolus injection assay, for the preparation of a medicament for the treatment of a brain disease associated with proliferative processes.

REFERENCE COUNT:

10 THERE ARE 10 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L102 ANSWER 6 OF 24 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2003:511314 CAPLUS Full-text


DOCUMENT NUMBER: 139:85166

TITLE: Method for producing Cl-C6 fragments of epothilones

and the derivatives thereof

Klar, **Ulrich**; Berger, Markus; **Buchmann**,

DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 142:355075
 GI



I

AB An efficient chiral pool synthesis of the C1-C6 fragment of epothilones, e.g. I, starting from readily available (-)-pantolactone is described.
 REFERENCE COUNT: 27
 THERE ARE 27 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L102 ANSWER 3 OF 24 CAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 2005:29293 CAPLUS Full-text
 DOCUMENT NUMBER: 142:113814

TITLE: Method for producing C1-C15 fragments of epothilones and derivatives thereof


INVENTOR(S): Klat, Ulrich; Buchmann, Bernd; Schwede, Wolfgang; Skraballa, Werner
 Schering Aktiengesellschaft, Germany
 PCT Int. Appl., 48 pp.
 CODEN: PIXX22

PATENT ASSIGNEE(S): Patent
 DOCUMENT TYPE: German
 LANGUAGE: 1
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2005003071	A1	20050113	WO 2004-EP6685	20040619
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, GU, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SV, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CG, CI, CM, GN, GW, GW, ML, MR, NE, SN, TD, TG				
DE 10331004	A1	20050224	DE 2003-10331004	20030703
AU 2004254200	A1	20050113	AU 2004-254200	20040619
CA 2531078	A1	20050113	CA 2004-2531078	20040619
EP 1641734	A1	20060405	EP 2004-740122	20040619
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK, HR				
QN 1816514	A	20060809	QN 2004-80019005	20040619
BR 2004012179	A	20060822	BR 2004-12179	20040619
IN 2006DN00056	A	20070824	IN 2006-DN56	20060103
MX 2006PA00172	A	20060427	MX 2006-PA172	20060105

NO 2006000554 A 20060403 NO 2006-554 20060202
 US 2007142675 A1 20070621 US 2006-563058 20060619
 PRIORITY APPLN. INFO.: DE 2003-10331004 A 20030703
 WO 2004-EP6685 W 20040619
 OTHER SOURCE(S): CASREACT 142:113814; MARPAT 142:113814
 GI

for 8 h



I

II

AB The invention relates to a method for preparing C1-C15 fragments I [R1a, R1b = H, C1-10-alkyl, aryl, C7-20-alkenyl, R1aR1b = (CH2)m; m = 2 - 5; R2a, R2b = H, C1-10-alkyl, C2-10-alkenyl, C2-10-alkynyl, aryl, C7-20-alkenyl, R2aR2b = (CH2)n; n = 2 - 5; R3 = H, C1-10-alkyl, aryl, C7-20-alkenyl, R3a, R3b = H, C1-10-alkyl, aryl, C7-20-alkenyl, R4aR4b = (CH2)p; p = 2 - 5; R5 = H, C1-10-alkyl, aryl, C7-20-alkenyl, R6, R7 = H; R6R7 = bond, O; G = X:CR8, bi- or tricyclic aryl; R8 = H, halogen, (un)substituted C1-20-alkyl, aryl, C7-20-alkenyl; X = O, (OR23)2, C2-10-alkylene-α,ω-dioxy, H(OR9), CRIOR11; R23 = C1-20-alkyl; R9 = H, protecting group; R10, R11 = H, C1-10-alkyl, aryl, C7-20-alkenyl, CRIOR11 = 5 - to 7-membered carbocycle; R13 = CH2OR13a, CH2-halo, CHO, CO2R13b, CO-halo; R13a, R14a = H, SO2alkyl, SO2-aryl, SO2-alkyl, R13aR14a = (CH2)q; q = 3 - 6; R20 = O-PG, R6R7 = bond, R13 = CO2H, R14 = OSiMe2CMe3, Z = O] C7-20-alkenyl; R15a, R15b = H, C1-10-alkyl, aryl, C7-20-alkenyl; R15aR15b = (CH2)q; q = 3 - 6; R20 = O-PG, NHR29, N3; Z = O, H(OR12); R12 = H, PG] of epothilones and derivs. The procedure comprises the bonding of a C1-C6 fragment, R13CH2CHRI4CRIA1R1BC(:O)CHR2aR2b, to a C7-C12 fragment, R5C(:V)(CH2)3CR4aR4bC(:W)R3a [V, W = O, (OR23)2, C2-10-alkylene-α,ω-dioxy, H(OR9)], to form a C1-C12 fragment, R5C(:V)(CH2)3CR4aR4bCR3a(O-PG)4)CR2aR2bC(:Z)CRIA1R1BCRIA4CH2R13 [PG = H, protecting group], which is then treated with a C13-C15 fragment, G-CR20-CH2CHR7-R21 [R7 = H; R20 = halogen, N3, NHR29, OH, O-PG, NR29-PG, C1-20-(perfluoro)alkylsulfonyloxy, (C1-4-alkyl, NO2, Cl, Br-substituted) benzylloxy, NR29SO2Me, NR29C(:O)Me, CH2C(:O)Me; R21 = OH, halo, O-PG, P-Ph3Hal- (Hal = F, Cl, Br, I), P(O)(OQ)2 (Q = C1-10-alkyl, Ph), P(O)Ph2; R29 = H, C1-6-alkyl], to form the C1-C15 epothilone intermediate product I. Thus, I [R1a = R1b = R5 = Me, R2a = CH2CH:CH2-β, R2b = R4b = H-α, R3 = H-β, R4a = Me-β, R6R7 = bond, R13 = CO2H, R14 = OSiMe2CMe3, Z = O] was prepared from (S)-4-(2-methyl-3-oxohept-6-en-2-yl)-2,2-dimethyl-1,3-dioxane via lithiation and reaction with (ZS,6RS)-2-methyl-6-((tert-butylidimethylsilyl)oxy)heptanal, tetrahydropyranlation, desilylation with Bu4NF in THF, oxidation in CH2Cl2 containing N-methylmorpholine N-oxide and catalytic tetrapropylammonium perchlorate, Wittig reaction with [(3S)-3-(2-methylbenzothiazol-5-yl)propyl]triphenylphosphonium iodide, Geisopropylidenation/tetrahydropyranlation with catalytic 4-MeOC6H4SO3H in EtOH, silylation with CF3SO2SiMe2CMe3, regioselective desilylation with (i)-camphor-10-sulfonic acid, Swern oxidation with DMSO/(COCl)2 in CH2Cl2 and carbonyl oxidation with NaOCl2 in aqueous THF/Me3COH. The produced C1-C15 epothilone intermediate products can be converted into the intrinsically

=> file registry
 FILE 'REGISTRY' ENTERED AT 12:11:27 ON 11 OCT 2007
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
 COPYRIGHT (C) 2007 AMERICAN CHEMICAL SOCIETY (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 10 OCT 2007 HIGHEST RN 950149-06-1
 DICTIONARY FILE UPDATES: 10 OCT 2007 HIGHEST RN 950149-06-1

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH JUNE 29, 2007

Please note that search-term pricing does apply when conducting SmartSELECT searches.

REGISTRY includes numerically searchable data for experimental and predicted properties as well as tags indicating availability of experimental property data in the original document. For information on property searching in REGISTRY, refer to:

<http://www.cas.org/support/stngen/stdoc/properties.html>

=> file caplus
 FILE 'CAPLUS' ENTERED AT 12:11:29 ON 11 OCT 2007
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
 COPYRIGHT (C) 2007 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 11 OCT 2007 VOL 147 ISS 16
 FILE LAST UPDATED: 10 OCT 2007 (20071010/ED)

Effective October 17, 2005, revised CAS Information Use Policies apply. They are available for your review at:

<http://www.cas.org/infopolicy.html>
 "OBI" IS DEFAULT SEARCH FIELD FOR 'CAPLUS' FILE


=> d stat que L100
 L91 80 SEA FILE=CAPLUS ABB=ON PUJ=ON KJAR U7/AU
 L92 116 SEA FILE=CAPLUS ABB=ON PUJ=ON BUCHMANN B7/AU
 L93 60 SEA FILE=CAPLUS ABB=ON PUJ=ON SCHWEDE W7/AU
 L94 186 SEA FILE=CAPLUS ABB=ON PUJ=ON SKUBALLA W7/AU
 L102 24 SEA FILE=CAPLUS ABB=ON PUJ=ON L91 AND L92 AND L93 AND L94

=> d ibib abs L102 tot

L102 ANSWER 1 OF 24 CAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 2005:132732 CAPLUS Full-Text
 DOCUMENT NUMBER: 142:355075
 TITLE: Total synthesis and antitumor activity of 2K-EPO: The first fully synthetic epothilone in clinical development

AUTHOR(S): Kjar, Ulrich; Buchmann, Bernd; Schwede, Wolfgang; Skuballa, Werner; Hoffmann, Jens; Lichtner, Rosemarie B. Schering AG, Research Center Europe, Berlin, Germany
 CORPORATE SOURCE: Angewandte Chemie, International Edition (2006), 45(47), 7942-7948

CODEN: ACIEF5 ISSN: 1433-7851
 PUBLISHER: Wiley-VCH Verlag GmbH & Co. KGaA
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 146:229070
 GI



AB From about 350 active epothilone analogs synthesized by a highly convergent synthesis, 2K-EPO (1) was chosen for clin. development on the basis of its outstanding preclin. data. This compound exhibits higher activity and efficacy than taxanes, such as paclitaxel and second-generation epothilones, a fast and efficient cellular uptake, no recognition by efflux mechanisms, and an improved therapeutic window.

REFERENCE COUNT: 39 THERE ARE 39 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L102 ANSWER 2 OF 24 CAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 2005:132732 CAPLUS Full-Text
 DOCUMENT NUMBER: 142:355075
 TITLE: Efficient chiral pool synthesis of the C1-C6 fragment of epothilones

AUTHOR(S): Kjar, Ulrich; Roehr, Bodo; Kuczynski, Frank; Schwede, Wolfgang; Berger, Markus; Skuballa, Werner; Buchmann, Bernd

CORPORATE SOURCE: Research Laboratories of Schering AG, Berlin, 13342, Germany
 SOURCE: Synthesis (2005), (2), 301-305
 CODEN: SYNTBF; ISSN: 0039-7881
 PUBLISHER: Georg Thieme Verlag

D STAT QUE L19

FILE 'BABS' ENTERED AT 15:47:02 ON 11 OCT 2007

D STAT QUE L14

L25 FILE 'ZCAPLUS, BEILSTEIN, BABS' ENTERED AT 15:47:21 ON 11 OCT 2007
25 DUP REM L6 L19 L14 (7 DUPLICATES REMOVED)
ANSWERS '1-18' FROM FILE ZCAPLUS
ANSWERS '19-25' FROM FILE BEILSTEIN
D IBIB ABS HITSTR L25 1-18
D IDE ALLREF L25 19-25

FILE HOME

FILE STNGUIDE :

FILE CONTAINS CURRENT INFORMATION.

LAST RELOADED: Oct 5, 2007 (20071005/UP).

FILE REGISTRY

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 10 OCT 2007 HIGHEST RN 950149-06-1

DICTIONARY FILE UPDATES: 10 OCT 2007 HIGHEST RN 950149-06-1

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH June 29, 2007

Please note that search-term pricing does apply when conducting SmartSELECT searches.

REGISTRY includes numerically searchable data for experimental and predicted properties as well as tags indicating availability of experimental property data in the original document. For information on property searching in REGISTRY, refer to:

<http://www.cas.org/support/stngen/stndoc/properties.html>

FILE ZCAPLUS

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS is strictly prohibited.

FILE COVERS 1907 - 11 Oct 2007 VOL 147 ISS 16

FILE LAST UPDATED: 10 Oct 2007 (20071010/ED)

New CAS Information Use Policies, enter HELP USAGETERMS for details.

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> d his full

(FILE 'HOME' ENTERED AT 15:09:07 ON 11 OCT 2007)

FILE 'STNGUIDE' ENTERED AT 15:22:34 ON 11 OCT 2007

FILE 'REGISTRY' ENTERED AT 15:31:01 ON 11 OCT 2007

L1 STRUCTURE UPLOADED
L2 0 SEA SSS SAM L1
L3 SCREEN 1008
L4 0 SEA SSS SAM L1 AND L3
L5 68 SEA SSS FUL L1 AND L3
SAVE TEMP LAO058STR1CL/A L5

FILE 'ZCAPLUS' ENTERED AT 15:37:30 ON 11 OCT 2007

L6 18 SEA ABB=ON PLU=ON L5
E US2006-563058/APPS
L7 1 SEA ABB=ON PLU=ON US2006-563058/AP
D SCA
L8 1 SEA ABB=ON PLU=ON L6 AND L7
D SCA

FILE 'BEILSTEIN' ENTERED AT 15:40:57 ON 11 OCT 2007

L9 0 SEA SSS SAM L1
L10 0 SEA SSS SAM L1 AND L3
L11 38 SEA SSS FUL L1 AND L3
L12 26 SEA ABB=ON PLU=ON L11/COM
L13 5 SEA ABB=ON PLU=ON L12 AND BABSAN/FA
SEL BABSAN

FILE 'BABS' ENTERED AT 15:42:39 ON 11 OCT 2007

L14 7 SEA ABB=ON PLU=ON (6300090/BABSAN OR 6630563/BABSAN OR
6085475/BABSAN OR 6376421/BABSAN OR 6410256/BABSAN OR 6473119/B
ABSAN OR 6597156/BABSAN)

FILE 'BEILSTEIN' ENTERED AT 15:43:07 ON 11 OCT 2007

L15 21 SEA ABB=ON PLU=ON L12 NOT L13
L16 14 SEA ABB=ON PLU=ON L15 AND RN/FA

FILE 'REGISTRY' ENTERED AT 15:43:45 ON 11 OCT 2007

L17 14 SEA ABB=ON PLU=ON L5 AND BEILSTEIN/LC NOT CAPLUS
L18 0 SEA ABB=ON PLU=ON L5 AND BEILSTEIN/LC NOT CAPLUS/LC

FILE 'BEILSTEIN' ENTERED AT 15:44:11 ON 11 OCT 2007

L19 7 SEA ABB=ON PLU=ON L15 NOT L16
L20 252 SEA ABB=ON PLU=ON KLAR U?/AU
L21 351 SEA ABB=ON PLU=ON BUCHMANN B?/AU
L22 436 SEA ABB=ON PLU=ON SCHWEDE W?/AU
L23 369 SEA ABB=ON PLU=ON SKUBALLA W?/AU
L24 0 SEA ABB=ON PLU=ON L19 AND (L20 OR L21 OR L22 OR L23)
D COST

FILE 'REGISTRY' ENTERED AT 15:46:28 ON 11 OCT 2007

FILE 'ZCAPLUS' ENTERED AT 15:46:31 ON 11 OCT 2007
D STAT QUE L6

FILE 'BEILSTEIN' ENTERED AT 15:46:51 ON 11 OCT 2007

Field Availability:

Code	Name	Occurrence
BRN	Beilstein Records	1
MF	Molecular Formula	1
FW	Formular Weight	1
LN	Lawson Number	3
FS	File Segment	1
CTYPE	Compound Type	1
CONSID	Constitution ID	1
TAUTID	Tautomer ID	1
BSO	Beilstein Citation	1
DED	Entry Date	1
DUPD	Update Date	1
NMR	Nuclear Magnetic Resonance	1

This substance also occurs in Reaction Documents:

Code	Name	Occurrence
RX	Reaction Documents	1
RXPRO	Substance is Reaction Product	1

All References:

ALLREF

1. Aberhart et al., J.Chem.Soc.Perkin Trans.1, CODEN: JCPRB4, <1974>, 816,823

BRN	Beilstein Records	1
CN	Chemical Name	1
AUN	Autonomname	1
MF	Molecular Formula	1
FW	Formular Weight	1
LN	Lawson Number	6
FS	File Segment	1
CTYPE	Compound Type	1
CONSID	Constitution ID	1
TAUTID	Tautomer ID	1
BSO	Beilstein Citation	1
DED	Entry Date	1
DUPD	Update Date	1
CDER	Chemical Derivative	1

This substance also occurs in Reaction Documents:


Code	Name	Occurrence
RX	Reaction Documents	1
RXPRO	Substance is Reaction Product	1

All References:
ALLREF

1. Fehr, T. et al., J.Chem.Soc.Perkin Trans.1, CODEN: JCPRB4, <1974>, 836-847

L25 ANSWER 25 OF 25 BEILSTEIN COPYRIGHT 2007 BEILSTEIN MDL on STN

Beilstein Records (BRN):	1677640
Molec. Formula (MF):	C40 H60 D2 O21
Molecular Weight (MW):	880.93
Lawson Number (LN):	17586, 1155, 680
File Segment (FS):	Stereo compound
Compound Type (CTYPE):	heterocyclic
Constitution ID (CONSID):	1560573
Tautomer ID (TAUTID):	1638549
Beilstein Citation (BSO):	5-17
Entry Date (DED):	1988/11/30
Update Date (DUPD):	1990/02/07




RX	Reaction Documents	1
RXPRO	Substance is Reaction Product	1

All References:
ALLREF

1. Aberhart et al., J.Chem.Soc.Perkin Trans.1, CODEN: JCPRB4, <1974>, 816,823

L25 ANSWER 24 OF 25 BEILSTEIN COPYRIGHT 2007 BEILSTEIN MDL on STN

Beilstein Records (BRN):	1677845
Chemical Name (CN):	2-butyl-18-(3,4-dimethoxy-5-methoxymethyl-tetrahydro-furan-2-yloxy)-32-hydroxy-3,7,11,15,19,21,23,25,27-nonamethoxy-4,16-dimethyl-tritriacont-16-enoic acid 1-(2-hydroxy-1-methyl-ethyl)-3-methoxy-2-methyl-6-(N,N',N'-trimethyl-guanidino)-hexyl ester
Autonom Name (AUN):	2-butyl-18-(3,4-dimethoxy-5-methoxymethyl-tetrahydro-furan-2-yloxy)-32-hydroxy-3,7,11,15,19,21,23,25,27-nonamethoxy-4,16-dimethyl-tritriacont-16-enoic acid 1-(2-hydroxy-1-methyl-ethyl)-3-methoxy-2-methyl-6-(N,N',N'-trimethyl-guanidino)-hexyl ester
Molec. Formula (MF):	C71 H139 N3 O19
Molecular Weight (MW):	1338.89
Lawson Number (LN):	17586, 3238, 2817, 2294, 1762, 289
File Segment (FS):	Stereo compound
Compound Type (CTYPE):	heterocyclic
Constitution ID (CONSID):	1561576
Tautomer ID (TAUTID):	1636624
Beilstein Citation (BSO):	5-17
Entry Date (DED):	1988/11/30
Update Date (DUPD):	1991/03/25




Field Availability:

Code	Name	Occurrence
------	------	------------

1. Chakraborty, T. K.; Dutta, S., Tetrahedron Lett., CODEN: TELEAY, 39(1-2), <1998>, 101-104; BABS-6085475

L25 ANSWER 23 OF 25 BEILSTEIN COPYRIGHT 2007 BEILSTEIN MDL on STN

Beilstein Records (BRN): 1965671
 Chemical Name (CN): heptadecane-1,3,5,7,9,11,16-heptaol
 Autonom Name (AUN): heptadecane-1,3,5,7,9,11,16-heptaol
 Molec. Formula (MF): C17H36O7
 Molecular Weight (MW): 352.47
 Lawson Number (LN): 687
 Compound Type (CTYPE): acyclic
 Constitution ID (CONSID): 1838397
 Tautomer ID (TAUTID): 1903620
 Beilstein Citation (BSO): 5-01
 Entry Date (DED): 1989/06/29
 Update Date (DUPD): 1997/12/03



Field Availability:

Code	Name	Occurrence
BRN	Beilstein Records	1
BPR	Beilstein Preferred RN	1
RN	CAS Registry Number	1
CN	Chemical Name	1
AUN	Autonomname	1
MF	Molecular Formula	1
FW	Formular Weight	1
LN	Lawson Number	1
CTYPE	Compound Type	1
CONSID	Constitution ID	1
TAUTID	Tautomer ID	1
BSO	Beilstein Citation	1
DED	Entry Date	1
DUPD	Update Date	1
IR	Infrared Spectrum	1
MP	Melting Point	1
MS	Mass Spectrum	1

This substance also occurs in Reaction Documents:

Code	Name	Occurrence
=====	=====	=====

Chemical Name (CN): 12-benzyloxy-7-(tert-butyl-dimethyl-silanyloxy)-3-hydroxy-4,4,6,8-tetramethyl-5-oxo-dodecanoic acid tert-butyl ester

Autonom Name (AUN): 12-benzyloxy-7-(tert-butyl-dimethyl-silanyloxy)-3-hydroxy-4,4,6,8-tetramethyl-5-oxo-dodecanoic acid tert-butyl ester

Molec. Formula (MF): C33 H58 O6 Si

Molecular Weight (MW): 578.90

Lawson Number (LN): 5228, 3798, 3777, 2672, 318

File Segment (FS): Stereo compound

Compound Type (CTYPE): isocyclic


Constitution ID (CONSID): 6834281

Tautomer ID (TAUTID): 7583399

Beilstein Citation (BSO): 6-06

Entry Date (DED): 1998/11/09

Update Date (DUPD): 1998/11/09




Field Availability:

Code	Name	Occurrence
BRN	Beilstein Records	1
CN	Chemical Name	1
AUN	Autonomname	1
MF	Molecular Formula	1
FW	Formular Weight	1
LN	Lawson Number	5
FS	File Segment	1
CTYPE	Compound Type	1
CONSID	Constitution ID	1
TAUTID	Tautomer ID	1
BSO	Beilstein Citation	1
DED	Entry Date	1
DUPD	Update Date	1

This substance also occurs in Reaction Documents:

Code	Name	Occurrence
RX	Reaction Documents	2
RXREA	Substance is Reaction Reactant	1
RXPRO	Substance is Reaction Product	1

All References:
ALLREF



Field Availability:

Code	Name	Occurrence
BRN	Beilstein Records	1
CN	Chemical Name	1
AUN	Autonomname	1
MF	Molecular Formula	1
FW	Formular Weight	1
LN	Lawson Number	3
FS	File Segment	1
CTYPE	Compound Type	1
CONSID	Constitution ID	1
TAUTID	Tautomer ID	1
DED	Entry Date	1
DUPD	Update Date	1

This substance also occurs in Reaction Documents:

Code	Name	Occurrence
RX	Reaction Documents	2
RXREA	Substance is Reaction Reactant	1
RXPRO	Substance is Reaction Product	1

All References:

ALLREF

1. Dong, Steven D.; Sundermann, Kurt; Smith, Karen M. J.; Petryka, Joseph; Liu, Fenghua; Myles, David C., Tetrahedron Lett., CODEN: TELEAY, 45(9), <2004>, 1945 - 1948; BABS-6591064
2. Dong, Steven D.; Sundermann, Kurt; Smith, Karen M. J.; Petryka, Joseph; Liu, Fenghua; Myles, David C., Tetrahedron Lett., CODEN: TELEAY, 45(9), <2004>, 1945 - 1948; BABS-6441824

L25 ANSWER 22 OF 25 BEILSTEIN COPYRIGHT 2007 BEILSTEIN MDL on STN

Beilstein Records (BRN): 7955235

Field Availability:

Code	Name	Occurrence
BRN	Beilstein Records	1
CN	Chemical Name	1
AUN	Autonomname	1
MF	Molecular Formula	1
FW	Formular Weight	1
LN	Lawson Number	4
FS	File Segment	1
CTYPE	Compound Type	1
CONSID	Constitution ID	1
TAUTID	Tautomer ID	1
DED	Entry Date	1
DUPD	Update Date	1

This substance also occurs in Reaction Documents:

Code	Name	Occurrence
RX	Reaction Documents	2
RXREA	Substance is Reaction Reactant	1
RXPRO	Substance is Reaction Product	1

All References:

ALLREF

1. Dong, Steven D.; Sundermann, Kurt; Smith, Karen M. J.; Petryka, Joseph; Liu, Fenghua; Myles, David C., Tetrahedron Lett., CODEN: TELEAY, 45(9), <2004>, 1945 - 1948; BABS-6591064
2. Dong, Steven D.; Sundermann, Kurt; Smith, Karen M. J.; Petryka, Joseph; Liu, Fenghua; Myles, David C., Tetrahedron Lett., CODEN: TELEAY, 45(9), <2004>, 1945 - 1948; BABS-6441824

L25 ANSWER 21 OF 25 BEILSTEIN COPYRIGHT 2007 BEILSTEIN MDL on STN

Beilstein Records (BRN): 9738862

Chemical Name (CN): 3,7-bis-(tert-butyl-dimethyl-silanyloxy)-4,4,6,8-tetramethyl-5,12-dioxo-tridecanoic acid

Autonom Name (AUN): 3,7-bis-(tert-butyl-dimethyl-silanyloxy)-4,4,6,8-tetramethyl-5,12-dioxo-tridecanoic acid

Molec. Formula (MF): C29 H58 O6 Si2

Molecular Weight (MW): 558.94

Lawson Number (LN): 3798, 3777, 2674

File Segment (FS): Stereo compound

Compound Type (CTYPE): acyclic

Constitution ID (CONSID): 8203590

Tautomer ID (TAUTID): 9128603

Entry Date (DED): 2004/10/23

Update Date (DUPD): 2007/02/05

CTYPE	Compound Type	1
CONSID	Constitution ID	1
TAUTID	Tautomer ID	1
DED	Entry Date	1
DUPD	Update Date	1

This substance also occurs in Reaction Documents:

Code	Name	Occurrence
RX	Reaction Documents	2
RXREA	Substance is Reaction Reactant	1
RXPRO	Substance is Reaction Product	1


All References:

ALLREF

1. Dong, Steven D.; Sundermann, Kurt; Smith, Karen M. J.; Petryka, Joseph; Liu, Fenghua; Myles, David C., Tetrahedron Lett., CODEN: TELEAY, 45(9), <2004>, 1945 - 1948; BABS-6591064
2. Dong, Steven D.; Sundermann, Kurt; Smith, Karen M. J.; Petryka, Joseph; Liu, Fenghua; Myles, David C., Tetrahedron Lett., CODEN: TELEAY, 45(9), <2004>, 1945 - 1948; BABS-6441824


L25 ANSWER 20 OF 25 BEILSTEIN COPYRIGHT 2007 BEILSTEIN MDL on STN

Beilstein Records (BRN):	9738952
Chemical Name (CN):	3,7-bis-(tert-butyl-dimethyl-silanyloxy)- 4,4,6,8-tetramethyl-5,12-dioxo-tridecanoic acid methyl ester
Autonom Name (AUN):	3,7-bis-(tert-butyl-dimethyl-silanyloxy)- 4,4,6,8-tetramethyl-5,12-dioxo-tridecanoic acid methyl ester
Molec. Formula (MF):	C30 H60 O6 Si2
Molecular Weight (MW):	572.97
Lawson Number (LN):	3798, 3777, 2674, 289
File Segment (FS):	Stereo compound
Compound Type (CTYPE):	acyclic
Constitution ID (CONSID):	8203696
Tautomer ID (TAUTID):	9128457
Entry Date (DED):	2004/10/23
Update Date (DUPD):	2007/02/05



L25 ANSWER 19 OF 25 BEILSTEIN COPYRIGHT 2007 BEILSTEIN MDL on STN


Beilstein Records (BRN): 9747518
 Chemical Name (CN): 3,7-bis-(tert-butyl-dimethyl-silanyloxy)-
 4,4,6,8-tetramethyl-5,12-dioxo-tridecanoic
 acid 2-methyl-3-(2-methyl-thiazol-4-yl)-1-
 (2-oxo-ethyl)-allyl ester
 Autonom Name (AUN): 3,7-bis-(tert-butyl-dimethyl-silanyloxy)-
 4,4,6,8-tetramethyl-5,12-dioxo-tridecanoic
 acid 2-methyl-3-(2-methyl-thiazol-4-yl)-1-
 (2-oxo-ethyl)-allyl ester
 Molec. Formula (MF): C39 H69 N O7 S Si2
 Molecular Weight (MW): 752.21
 Lawson Number (LN): 31322, 3798, 3777, 2674
 File Segment (FS): Stereo compound
 Compound Type (CTYPE): heterocyclic
 Constitution ID (CONSID): 8211765
 Tautomer ID (TAUTID): 9136618
 Entry Date (DED): 2004/10/23
 Update Date (DUPD): 2007/02/05



Field Availability:


Code	Name	Occurrence
BRN	Beilstein Records	1
CN	Chemical Name	1
AUN	Autonomname	1
MF	Molecular Formula	1
FW	Formular Weight	1
LN	Lawson Number	4
FS	File Segment	1

Absolute stereochemistry.




RN 53294-58-9 ZCAPLUS

CN 1,2,3,5,7,9,11,16-Heptadecaneoctol (9CI) (CA INDEX NAME)




RN 53294-63-6 ZCAPLUS

CN 1,2,3,5,7,9,11,16-Heptadecaneoctol, octaacetate (9CI) (CA INDEX NAME)




RN 53294-64-7 ZCAPLUS

CN 1,3,5,7,9,14-Pentadecanehexol (9CI) (CA INDEX NAME)




RN 53294-65-8 ZCAPLUS

CN 1,3,5,7,9,14-Pentadecanehexol, hexaacetate (9CI) (CA INDEX NAME)



RN 53294-66-9 ZCAPLUS

CN 1,3,5,7,9,11,16-Heptadecaneheptol, heptaacetate (9CI) (CA INDEX NAME)



AB The structures of secopyrimycins A [MeO2CCHBuCH(OH)CHMe(CH2)2[CH(OH)(CH2)3]2[CH(OH)]2Me], B (I), and C, [AcNH(CH2)3[CH(OH)CHMe]2CH2OH] were detd. from their high resolution mass spectra and/or those of their simpler derivs. and chemical degradation products.


IT 54799-27-8

RL: PRP (Properties)

(mol. structure of, mass spectrum in relation to)

RN 54799-27-8 ZCAPLUS

CN 1,3,5,7,9,11,16-Heptadecaneheptol, 2-(α -D-arabinofuranosyloxy)-(9CI) (CA INDEX NAME)



PAGE 1-B

—CH2—OH

—OH

L25 ANSWER 18 OF 25 ZCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1974:437769 ZCAPLUS Full-text

DOCUMENT NUMBER: 81:37769

TITLE: Constitution of primycin. I. Characterization, functional groups, and degradation to the secopyrimycins

AUTHOR(S): Aberhart, John; Jain, Rup C.; Fehr, Theo; De Mayo, Paul; Szilagyi, Imre

CORPORATE SOURCE: Dep. Chem., Univ. West. Ont., London, ON, Can.

SOURCE: Journal of the Chemical Society, Perkin Transactions 1: Organic and Bio-Organic Chemistry (1972-1999) (1974), (7), 816-26
CODEN: JCPRB4; ISSN: 0300-922X

DOCUMENT TYPE: Journal

LANGUAGE: English

GI For diagram(s), see printed CA Issue.

AB The proposed structure of primycin (I) was determined mainly from the chemical and spectral data of its degradation products.

IT 53294-56-7P 53294-58-9P 53294-63-6P

53294-64-7P 53294-65-8P 53294-66-9P


RL: SPN (Synthetic preparation); PREP (Preparation) (preparation of)

RN 53294-56-7 ZCAPLUS

CN 1,3,5,7,9,11,16-Heptadecaneheptol, 2-[(2,3,5-tri-O-acetyl- α -D-arabinofuranosyl)oxy]-, heptaacetate (9CI) (CA INDEX NAME)


arabinofuranosyl)oxy]-, 6-[[[(dimethylamino)iminomethyl]methylamino]-1-(2-hydroxy-1-methylethyl)-3-methoxy-2-methylhexyl ester, monohydrochloride (9CI) (CA INDEX NAME)

PAGE 1-A



● HCl

PAGE 1-B



L25 ANSWER 17 OF 25 ZCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1974:436929 ZCAPLUS Full-text

DOCUMENT NUMBER: 81:36929

TITLE: Constitution of primycin. II. Mass spectra of the secoprimycins

AUTHOR(S): Gracey, D. E. Fergus; Baczynskyj, Lubomir; Martin, Trevor I.; MacLean, David B.

CORPORATE SOURCE: Dep. Chem., McMaster Univ., Hamilton, ON, Can.

SOURCE: Journal of the Chemical Society, Perkin Transactions 1: Organic and Bio-Organic Chemistry (1972-1999) (1974), (7), 827-36

CODEN: JCPRB4; ISSN: 0300-922X

DOCUMENT TYPE: Journal


LANGUAGE: English

GI For diagram(s), see printed CA Issue.


— OMe

CN 16-Tritriacontenoic acid, 2-butyl-3,7,11,15,19,21,23,25,27-nonamethoxy-4,16-dimethyl-32-oxo-18-[(2,3,5-tri-O-methyl- α -D-arabinofuranosyl)oxy]-, methyl ester (9CI) (CA INDEX NAME)

PAGE 1-A




PAGE 1-B



CN α -D-Arabinofuranoside, 2,4,6,8,10-pentamethoxy-15-(methoxy-d3)-1-(methoxy-d3-methyl)hexadecyl 2,3,5-tri-O-methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



CN 16-Tritriacontenoic acid, 2-butyl-32-hydroxy-3,7,11,15,19,21,23,25,27-nonamethoxy-4,16-dimethyl-18-[(2,3,5-tri-O-methyl- α -D-

CN 16-Tritriacontenoic acid, 32-(acetyloxy)-2-butyl-3,7,11,15,19,21,23,25,27-nonamethoxy-4,16-dimethyl-18-[(2,3,5-tri-O-methyl- α -D-arabinofuranosyl)oxy]-, methyl ester (9CI) (CA INDEX NAME).

PAGE 1-A




PAGE 1-B



CN 16-Tritriacontenoic acid, 2-butyl-32-hydroxy-3,7,11,15,19,21,23,25,27-nonamethoxy-4,16-dimethyl-18-[(2,3,5-tri-O-methyl- α -D-arabinofuranosyl)oxy]-, methyl ester (9CI) (CA INDEX NAME)

PAGE 1-A



DOCUMENT TYPE: Journal

LANGUAGE: English

GI For diagram(s), see printed CA Issue.

AB Methylation of primycin (I) gave, after chromatog., the trimethylated urea and guanidine derivs. (II and III). The structure of I was determined by a spectral study of the ozonolysis products of II and III and their degradation products.

IT 53293-97-3P 53293-98-4P 53293-99-5P

53294-00-1P 53294-03-4P 53294-04-5P


53294-05-6P 53294-53-4P 53503-23-4P

RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation of)

RN 53293-97-3 ZCAPLUS

CN α -D-Arabinofuranoside, 15-hydroxy-1-(hydroxymethyl)-2,4,6,8,10-pentamethoxyhexadecyl 2,3,5-tri-O-methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 53293-98-4 ZCAPLUS


CN α -D-Arabinofuranoside, 15-(acetyloxy)-1-[(acetyloxy)methyl]-2,4,6,8,10-pentamethoxyhexadecyl 2,3,5-tri-O-methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 53293-99-5 ZCAPLUS

CN 1,2,16-Heptadecanetriol, 3,5,7,9,11-pentamethoxy- (9CI) (CA INDEX NAME)



RN 53294-00-1 ZCAPLUS


CN Hexadecanal, 15-hydroxy-2,4,6,8,10-pentamethoxy- (9CI) (CA INDEX NAME)



RN 197634-37-0 ZCAPLUS

CN 4,15-Dioxa-3,16-disilaoctadecan-7-one, 5-(2-hydroxyethyl)-
2,2,3,3,6,6,8,10,17,17-decamethyl-16,16-diphenyl-9-(phenylmethoxy)-,
[5S-(5R*,8S*,9R*,10R*)]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 197634-39-2 ZCAPLUS

CN Dodecanoic acid, 3-[[[(1,1-dimethylethyl)dimethylsilyl]oxy]-12-[[[(1,1-
dimethylethyl)diphenylsilyl]oxy]-4,4,6,8-tetramethyl-5-oxo-7-
(phenylmethoxy)-, [3S-(3R*,6S*,7R*,8R*)]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L25 ANSWER 16 OF 25 ZCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1974:463919 ZCAPLUS Full-text


DOCUMENT NUMBER: 81:63919

TITLE: Constitution of primycin. III. Degradation of
methylated primycin, and the structure of primycin
AUTHOR(S): Fehr, Theo; Jain, Rup C.; De Mayo, Paul; Motl, O.;
Szilagyi, Imre; Baczynskyj, Lubomir; Gracey, D. E.
Fergus; Holland, Herbert L.; MacLean, David B.

CORPORATE SOURCE: Dep. Chem., Univ. West. Ont., London, ON, Can.

SOURCE: Journal of the Chemical Society, Perkin Transactions
1: Organic and Bio-Organic Chemistry (1972-1999)
(1974), (7), 836-47


CODEN: JCPRB4; ISSN: 0300-922X



RN 197634-31-4 ZCAPLUS

CN 3-Decanone, 2-(2,2-dimethyl-1,3-dioxan-4-yl)-10-[[[(1,1-dimethylethyl)diphenylsilyl]oxy]-2,4,6-trimethyl-5-(phenylmethoxy)-, [4S-[4R*(4S*,5R*,6R*)]]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 197634-33-6 ZCAPLUS

CN 5-Dodecanone, 12-[[[(1,1-dimethylethyl)diphenylsilyl]oxy]-1,3-dihydroxy-4,4,6,8-tetramethyl-7-(phenylmethoxy)-, [3S-(3R*,6S*,7R*,8R*)]]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 197634-35-8 ZCAPLUS

CN 4,17-Dioxa-3,18-disilaicosan-9-one, 7-[[[(1,1-dimethylethyl)diphenylsilyl]oxy]-2,2,3,3,8,8,10,12,19,19-decamethyl-18,18-diphenyl-11-(phenylmethoxy)-, [7S-(7R*,10S*,11R*,12R*)]]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



IT 197634-28-9P 197634-29-0P 197634-30-3P

197634-31-4P 197634-33-6P 197634-35-8P

197634-37-0P 197634-39-2P


RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(intermediates in the total synthesis of epothilones A and B)

RN 197634-28-9 ZCAPLUS

CN 3-Decanone, 2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-10-[[[(1,1-dimethylethyl)dimethylsilyl]oxy]-5-hydroxy-2,4,6-trimethyl-, (4R,5S,6S)-(9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 197634-29-0 ZCAPLUS

CN 3-Decanone, 2-(2,2-dimethyl-1,3-dioxan-4-yl)-10-[[[(1,1-dimethylethyl)dimethylsilyl]oxy]-2,4,6-trimethyl-5-(phenylmethoxy)-, [4S-[4R*(4S*,5R*,6R*)]]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 197634-30-3 ZCAPLUS

CN 3-Decanone, 2-(2,2-dimethyl-1,3-dioxan-4-yl)-10-hydroxy-2,4,6-trimethyl-5-(phenylmethoxy)-, [4S-[4R*(4S*,5R*,6R*)]]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML,
MR, NE, SN, TD, TG


AU 9721493	A	19980319	AU 1997-21493	19970115
AU 716610	B2	20000302		
EP 923583	A1	19990623	EP 1997-914077	19970115
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI				
NZ 334821	A	20001222	NZ 1997-334821	19970115
JP 2001500851	T	20010123	JP 1998-511141	19970115
US 5969145	A	19991019	US 1997-921512	19970902
US 6043372	A	20000328	US 1999-344713	19990625
US 6156905	A	20001205	US 2000-478466	20000106

PRIORITY APPLN. INFO.:


DE 1996-19636343	A	19960830
US 1996-27480P	P	19960926
DE 1996-19645361	A	19961028
DE 1996-19645362	A	19961028
WO 1997-DE111	W	19970115
US 1997-921512	A3	19970902
US 1999-344713	A3	19990625

OTHER SOURCE(S): CASREACT 127:346234

GI



I



II

AB Intermediates, e.g. 2-(2,2-dimethyl-1,3-dioxan-4-yl)-2-methyl-3-pentanone, 6-[(tert-butyldimethylsilyloxy)-2-methylhexanal, (S,4E)-3-benzyloxy-1-(tert-butyldimethylsilyloxy)-4-methyl-5-(2-ethyl-thiazol-4-yl)-4-pentene (I), (4S,6S)-10(tert-butyldimethylsilyloxy)-2-(2,2-dimethyl-1,3-dioxan-4-yl)-5-hydroxy-2,4,6-trimethyl-3-decanone (II) and (3S,6R,7S,8S)-7-benzyloxy-3-(tert-butyldimethylsilyloxy)-12-(tert-butyldiphenylsilyloxy)-4,4,6,8-tetramethyl-5-oxododecanoic acid, in the total synthesis of epothilones A and B are described. Epothilones A and B are natural products,.

IT 197634-08-5P

RL: PNU (Preparation, unclassified); PREP (Preparation)
(intermediates in the total synthesis of epothilones A and B)

RN 197634-08-5 ZCAPLUS

CN Dodecanoic acid, 12-[[[(1,1-dimethylethyl)dimethylsilyl]oxy]-3,7-dihydroxy-4,4,6,8-tetramethyl-5-oxo-, methyl ester, [3S-(3R*,6S*,7R*,8R*)]]- (9CI)
(CA INDEX NAME)

Absolute stereochemistry.

AB The title compds. [I; II; III; R1, R2 = H, alkyl, aryl, aralkyl; R3 = CH2OH, CH2OR; R4 = OH, OR; R = CR7R8; R7, R8 = H, alkyl, aryl, or R7R8 = (CH2)n; n = 2-6; R5, R6 = H, alkyl, aralkyl, or R5R6 = (CH2)m; m = 2-5; R9, R10 = H, protecting group; R11 = H, protecting group] including all the stereoisomers and their mixts. are prepared E.g., title compound (S)-III [R5 = R6 = Me, R9 = R11 = H, R10 = TBDPS] was prepared in 6 steps from D-(-)-pantolactone via reaction with 3,4-dihydro-2H-pyran, hydride reduction, Wittig reaction with methyltriphenylphosphonium bromide, protection of OH with TBDPS-Cl, de-tetrahydropyranyl, and reduction with borane-THF.

IT 197634-28-9P


RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(preparation of new C1-C6-fragments and application for synthesis of epothilone and epothilone derivs.)

RN 197634-28-9 ZCAPLUS

CN 3-Decanone, 2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-10-[[[(1,1-dimethylethyl)dimethylsilyl]oxy]-5-hydroxy-2,4,6-trimethyl-, (4R,5S,6S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.



L25 ANSWER 15 OF 25

ACCESSION NUMBER:

ZCAPLUS COPYRIGHT 2007 ACS on STN

1997:708545 ZCAPLUS Full-text

DOCUMENT NUMBER:

127:346234

TITLE:

Intermediate products within the total synthesis of Epothilones A and B

INVENTOR(S):

Schinzer, Dieter; Limberg, Anja; Boehm, Oliver M.

PATENT ASSIGNEE(S):

Schering A.-G., Germany

SOURCE:

Ger., 14 pp.

CODEN: GWXXAW

DOCUMENT TYPE:

Patent

LANGUAGE:

German

FAMILY ACC. NUM. COUNT: 2

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 19636343	C1	19971023	DE 1996-19636343	19960830
DE 19645361	A1	19980430	DE 1996-19645361	19961028
WO 9808849	A1	19980305	WO 1997-DE111	19970115
W: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR,				

TITLE: New (C1-C6)-fragments, method for their preparation and their application for synthesis of epothilone and epothilone derivatives

INVENTOR(S): Klar, Ulrich; Schwede, Wolfgang; Skuballa, Werner; Buchmann, Bernd; Schirner, Michael

PATENT ASSIGNEE(S): Schering A.-G., Germany

SOURCE: Ger. Offen., 18 pp.
CODEN: GWXXBX

DOCUMENT TYPE: Patent

LANGUAGE: German




FAMILY ACC. NUM. COUNT: 4

PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 19735578	A1	19990211	DE 1997-19735578	19970809
CA 2299608	A1	19990218	CA 1998-2299608	19980810
WO 9907692	A2	19990218	WO 1998-EP5064	19980810
WO 9907692	A3	19990514		
W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DK, EE, ES, FI, GB, GE, GH, GM, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW				
RW: GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
AU 9893409	A	19990301	AU 1998-93409	19980810
EP 1005465	A2	20000607	EP 1998-946309	19980810
EP 1005465	B1	20070725		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, CY, AL, MK				
JP 2001512723	T	20010828	JP 2000-506196	19980810
AT 368036	T	20070815	AT 1998-946309	19980810
US 2003144523	A1	20030731	US 2000-485292	20000503
PRIORITY APPLN. INFO.:			DE 1997-19735574	A 19970809
			DE 1997-19735575	A 19970809
			DE 1997-19735578	A 19970809
			DE 1997-19748928	A 19971024
			DE 1997-19749717	A 19971031
			DE 1997-19751200	A 19971113
			DE 1998-19813821	A 19980320
			WO 1998-EP5064	W 19980810

OTHER SOURCE(S): CASREACT 130:168164; MARPAT 130:168164

GI




Absolute stereochemistry.



RN 220774-78-7 ZCAPLUS

CN 3-Undecanone, 2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-10-hydroxy-2,6-dimethyl-4-(phenylmethyl)-5-[(tetrahydro-2H-pyran-2-yl)oxy]-, (4S,5R,6S)-(9CI) (CA INDEX NAME)


Absolute stereochemistry.




RN 220774-80-1 ZCAPLUS

CN 2,9-Undecanedione, 10-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-6,10-dimethyl-8-(phenylmethyl)-7-[(tetrahydro-2H-pyran-2-yl)oxy]-, (6S,7R,8S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.




L25 ANSWER 14 OF 25 ZCAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 1999:116659 ZCAPLUS Full-text
 DOCUMENT NUMBER: 130:168164



RN 220774-61-8 ZCAPLUS

CN 3-Undecanone, 2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-10-hydroxy-2,6-dimethyl-4-(phenylmethyl)-5-[(tetrahydro-2H-pyran-2-yl)oxy]-, (4R,5S,6S)-(9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 220774-62-9 ZCAPLUS

CN 2,9-Undecanedione, 10-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-6,10-dimethyl-8-(phenylmethyl)-7-[(tetrahydro-2H-pyran-2-yl)oxy]-, (6S,7S,8R)-(9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 220774-76-5 ZCAPLUS

CN 3-Undecanone, 2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-10-[[[1,1-dimethylethyl)diphenylsilyl]oxy]-2,6-dimethyl-4-(phenylmethyl)-5-[(tetrahydro-2H-pyran-2-yl)oxy]-, (4S,5R,6S)-(9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 220774-58-3 ZCAPLUS

CN 3-Undecanone, 2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-10-[[(1,1-dimethylethyl)diphenylsilyl]oxy]-5-hydroxy-2,6-dimethyl-4-(phenylmethyl)-, (4R,5S,6S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 220774-59-4 ZCAPLUS

CN 3-Undecanone, 2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-10-[[(1,1-dimethylethyl)diphenylsilyl]oxy]-5-hydroxy-2,6-dimethyl-4-(phenylmethyl)-, (4S,5R,6S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 220774-60-7 ZCAPLUS

CN 3-Undecanone, 2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-10-[[(1,1-dimethylethyl)diphenylsilyl]oxy]-2,6-dimethyl-4-(phenylmethyl)-5-[(tetrahydro-2H-pyran-2-yl)oxy]-, (4R,5S,6S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


Absolute stereochemistry.



RN 220774-21-0 ZCAPLUS

CN 3-Undecanone, 2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-10-[[[(1,1-dimethylethyl)diphenylsilyl]oxy]-4-ethyl-2,6-dimethyl-5-[(tetrahydro-2H-pyran-2-yl)oxy]-, (4R,5S,6S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 220774-22-1 ZCAPLUS

CN 3-Undecanone, 2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-4-ethyl-10-hydroxy-2,6-dimethyl-5-[(tetrahydro-2H-pyran-2-yl)oxy]-, (4R,5S,6S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 220774-23-2 ZCAPLUS

CN 2,9-Undecanedione, 10-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-8-ethyl-6,10-dimethyl-7-[(tetrahydro-2H-pyran-2-yl)oxy]-, (6S,7S,8R)- (9CI) (CA INDEX NAME)

melanoma, and acute lymphocytic and myelocytic leukemia. They are also suited for anti-angiogenesis therapy and for the treatment of chronic inflammatory diseases (psoriasis, arthritis). To prevent uncontrolled cell growth on, and for better tolerability of, medical implants, the derivs. can be introduced into or applied to polymeric materials. The compds. provided for in the invention can be used alone or, to achieve additive or synergistic effects, in combination with other principles and substance categories used in tumor therapy.

IT 220775-76-8


RL: RCT (Reactant); RACT (Reactant or reagent)

(preparation of epothilone derivs. as antitumor agents)

RN 220775-76-8 ZCAPLUS

CN 2,9-Undecanedione, 10-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-6,8,10-trimethyl-7-[(tetrahydro-2H-pyran-2-yl)oxy]-, (6S,7S,8R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



IT 220774-19-6P 220774-20-9P 220774-21-0P
220774-22-1P 220774-23-2P 220774-58-3P
220774-59-4P 220774-60-7P 220774-61-8P
220774-62-9P 220774-76-5P 220774-78-7P
220774-80-1P


RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(preparation of epothilone derivs. as antitumor agents)

RN 220774-19-6 ZCAPLUS

CN 3-Undecanone, 2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-10-[[[(1,1-dimethylethyl)diphenylsilyl]oxy]-4-ethyl-5-hydroxy-2,6-dimethyl-, (4R,5S,6S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 220774-20-9 ZCAPLUS


CN 3-Undecanone, 2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-10-[[[(1,1-dimethylethyl)diphenylsilyl]oxy]-4-ethyl-5-hydroxy-2,6-dimethyl-, (4S,5R,6S)- (9CI) (CA INDEX NAME)

DE 19735575	A1	19990211	DE 1997-19735575	19970809
DE 19735578	A1	19990211	DE 1997-19735578	19970809
DE 19748928	A1	19990429	DE 1997-19748928	19971024
DE 19749717	A1	19990506	DE 1997-19749717	19971031
DE 19751200	A1	19990520	DE 1997-19751200	19971113
DE 19813821	A1	19990923	DE 1998-19813821	19980320
CA 2299608	A1	19990218	CA 1998-2299608	19980810
AU 9893409	A	19990301	AU 1998-93409	19980810
EP 1005465	A2	20000607	EP 1998-946309	19980810
EP 1005465	B1	20070725		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, CY, AL, MK				
JP 2001512723	T	20010828	JP 2000-506196	19980810
ZA 9810403	A	20000515	ZA 1998-10403	19981113
IN 190805	A1	20030823	IN 1998-DE3413	19981116
US 2003144523	A1	20030731	US 2000-485292	20000503
IN 2002DE01305	A	20050311	IN 2002-DE1305	20021227
PRIORITY APPLN. INFO.:				
			DE 1997-19735574	A 19970809
			DE 1997-19735575	A 19970809
			DE 1997-19735578	A 19970809
			DE 1997-19748928	A 19971024
			DE 1997-19749717	A 19971031
			DE 1997-19751200	A 19971113
			DE 1998-19813821	A 19980320
			WO 1998-EP5064	W 19980810

OTHER SOURCE(S): MARPAT 130:196529
GI




AB Epothilone derivs. of formula I [X = O, alkylene- α,ω -dioxy, two alkoxy groups, etc.; Y = O, H₂; Z = O, (H, OH), (H, protected OH); R1a, R1b = H, alkyl, aryl, aralkyl, or together = (CH₂)_m where m = 2, 3, 4, 5; R2a, R2b = H, alkyl, aryl, aralkyl, or together = (CH₂)_n where n = 2, 3, 4, 5; when D-E = CH₂CH₂ or when Y = O, R2a or R2b may not be H/Me; R3 = H, alkyl, aryl, aralkyl; R4a, R4b = H, alkyl, aryl, aralkyl, or together = (CH₂)_p where p = 2, 3, 4, 5; D-E = CH₂CH₂, CH:CH, C.tplbond.C, 2,3-oxiranediy, CH(OH)CH(OH), CH(OH)CH₂; R5 = H, alkyl, aryl, aralkyl; R6, R7 = H, together = a saturated bond or O; R8 = H, alkyl, aryl, aralkyl all of which may be substituted] are prepared. Thus, the title compds. (4S,7R,8S,9S,13E,16S(E))- and (4S,7R,8S,9S,13Z,16S(E))-4,8-dihydroxy-7-ethyl-16-(1-methyl-2-(2-methyl-4-thiazolyl)ethenyl)-1-oxa-5,5,9,13-tetramethylcyclohexadec-13-en-2,6-dione (II) were prepared in many steps. The new compds. interact with tubulin by stabilizing formed microtubuli. They are capable of influencing cell division in a phase-specific manner and are suitable for the treatment of malignant tumors, such as ovarian, gastric, colon, breast, lung, head and neck carcinoma, adenocarcinoma, malignant



RN 220774-23-2 ZCAPLUS

CN 2,9-Undecanedione, 10-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-8-ethyl-6,10-dimethyl-7-[(tetrahydro-2H-pyran-2-yl)oxy]-, (6S,7S,8R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L25 ANSWER 13 OF 25 ZCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1999:126888 ZCAPLUS Full-text

DOCUMENT NUMBER: 130:196529

TITLE: Preparation of new epothilone derivatives as pharmaceutical agents

INVENTOR(S): Klar, Ulrich; Schwede, Wolfgang; Skuballa, Werner; Buchmann, Bernd; Schirner, Michael

PATENT ASSIGNEE(S): Schering Aktiengesellschaft, Germany

SOURCE: PCT Int. Appl., 185 pp. CODEN: PIXXD2


DOCUMENT TYPE: Patent

LANGUAGE: German

FAMILY ACC. NUM. COUNT: 4

PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9907692	A2	19990218	WO 1998-EP5064	19980810
WO 9907692	A3	19990514		
W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DK, EE, ES, FI, GB, GE, GH, GM, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW				
RW: GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
DE 19735574	A1	19990211	DE 1997-19735574	19970809



RN 220774-20-9 ZCAPLUS

CN 3-Undecanone, 2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-10-[[[(1,1-dimethylethyl)diphenylsilyl]oxy]-4-ethyl-5-hydroxy-2,6-dimethyl-, (4S,5R,6S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 220774-21-0 ZCAPLUS

CN 3-Undecanone, 2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-10-[[[(1,1-dimethylethyl)diphenylsilyl]oxy]-4-ethyl-2,6-dimethyl-5-[(tetrahydro-2H-pyran-2-yl)oxy]-, (4R,5S,6S)- (9CI) (CA INDEX NAME)



Absolute stereochemistry.



RN 220774-22-1 ZCAPLUS

CN 3-Undecanone, 2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-4-ethyl-10-hydroxy-2,6-dimethyl-5-[(tetrahydro-2H-pyran-2-yl)oxy]-, (4R,5S,6S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



AB New epothilone derivs. I (R1a,R1b = R2a,R2b = same or different H, alkyl, aryl, aralkyl or (CH₂)_{m,n} m, n = 2-5; R3 = H, alkyl, aryl, aralkyl; R4a,R4b = same or different H, alkyl, aryl, aralkyl or (CH₂)_p = 2-5, CH₂CH₂, CH=CH, C.tplbond.C, epoxy, CH(OH)CH(OH), CH(OH)CH₂; D-E = a group; R5 = H, alkyl, aryl, aralkyl; R6,R7 = H, bond, O; R8 = H, alkyl, aryl, aralkyl; X = O, OR₂₃ alkylene- α ,- ω -dioxy group straight or branched, OR₉ or the CR₁₀R₁₁ group where R₂₃ = alkyl, R₉ = H or protecting group and R₁₀,R₁₁ = same or different H, alkyl, aryl, aralkyl or R₁₀,R₁₁ = together with methylene are a 5-7 membered carbocyclic ring; Y = O or two H; Z = O or H/OR₁₂ and R₁₂ = H or a protecting group) were prepared. Thus E- and Z-II were prepared via a multistep synthesis. I cooperate with tubulin by stabilizing formed microtubuli. I are able phase specifically to affect the cell division and are suitable for the treatment of malignant ovarian, stomach, colon, adeno, breast, lung, head and neck tumors, malignant melanomas, acute lymphocytic and myelocytic leukemia. Derivs. of I are suitable for use in anti-angiogenic therapy as well as for treating chronic inflammatory diseases (psoriasis, arthritis). In order to prevent uncontrolled cell proliferations and to improve the compatibility of medical implants I can be applied or incorporated into polymeric materials. I can be used alone or to achieve additive or synergistic effects in combination with further principles and substance classes applicable in tumor therapy.

IT 220774-19-6P 220774-20-9P 220774-21-0P

220774-22-1P 220774-23-2P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)


(preparation of new epothilone derivs. and their pharmaceutical uses)

RN 220774-19-6 ZCAPLUS

CN 3-Undecanone, 2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-10-[[[(1,1-dimethylethyl)diphenylsilyl]oxy]-4-ethyl-5-hydroxy-2,6-dimethyl-, (4R,5S,6S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Absolute stereochemistry.




IT 303154-57-6P

RL: SPN (Synthetic preparation); PREP (Preparation)
(6-alkenyl and 6-alkynyl derivs. of epothilone)

RN 303154-57-6 ZCAPLUS

CN 3-Undecanone, 2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-10-[[[(1,1-dimethylethyl)diphenylsilyl]oxy]-5-hydroxy-2,6-dimethyl-4-[4-(trimethylsilyl)-3-butyryl]-, (4S,5R,6S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L25 ANSWER 12 OF 25

ACCESSION NUMBER:

ZCAPLUS COPYRIGHT 2007 ACS on STN

DOCUMENT NUMBER:

2000:738730 ZCAPLUS Full-text

TITLE:

133:309795

Preparation of new epothilone derivatives and their pharmaceutical uses

INVENTOR(S):

Klar, Ulrich; Schwede, Wolfgang; Skuballa, Werner; Buchmann, Bernd; Schirner, Michael

PATENT ASSIGNEE(S):

Schering A.-G., Germany

SOURCE:

Ger. Offen., 74 pp.

CODEN: GWXXBX

DOCUMENT TYPE:

Patent

LANGUAGE:


German

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 19908767	A1	20001019	DE 1999-19908767	19990218
PRIORITY APPLN. INFO.:			DE 1999-19908767	19990218
OTHER SOURCE(S):	MARPAT	133:309795		


GI



RN 303154-58-7 ZCAPLUS

CN 3-Undecanone, 2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-10-[[[(1,1-dimethylethyl)diphenylsilyl]oxy]-2,6-dimethyl-5-[(tetrahydro-2H-pyran-2-yl)oxy]-4-[4-(trimethylsilyl)-3-butynyl]-, (4R,5S,6S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 303154-59-8 ZCAPLUS

CN 3-Undecanone, 4-(3-butynyl)-2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-10-hydroxy-2,6-dimethyl-5-[(tetrahydro-2H-pyran-2-yl)oxy]-, (4R,5S,6S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 303154-60-1 ZCAPLUS

CN 2,9-Undecanedione, 8-(3-butynyl)-10-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-6,10-dimethyl-7-[(tetrahydro-2H-pyran-2-yl)oxy]-, (6S,7S,8R)- (9CI) (CA INDEX NAME)

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW
 RW: GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG

BR 2000010190	A	20020108	BR 2000-10190	20000501
EP 1173441	A1	20020123	EP 2000-922826	20000501
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, MC, PT, IE, SI, LT, LV, FI, RO				
HU 200201010	A2	20020828	HU 2002-1010	20000501
JP 2002543203	T	20021217	JP 2000-615619	20000501
EE 200100568	A	20030217	EE 2001-568	20000501
NZ 514989	A	20040227	NZ 2000-514989	20000501
AU 772750	B2	20040506	AU 2000-43103	20000501
IN 2001MN01305	A	20070504	IN 2001-MN1305	20011019
BG 106053	A	20020531	BG 2001-106053	20011026
NO 2001005278	A	20011221	NO 2001-5278	20011029
MX 2001PA11039	A	20030630	MX 2001-PA11039	20011030
ZA 2001009859	A	20030228	ZA 2001-9859	20011129
US 7125893	B1	20061024	US 2002-979939	20020606
US 2005113429	A1	20050526	US 2004-965802	20041018
IN 2005MN00837	A	20070608	IN 2005-MN837	20050802
US 2006046997	A1	20060302	US 2005-214988	20050831
JP 2007224038	A	20070906	JP 2007-104224	20070411

PRIORITY APPLN. INFO.:

DE 1999-19921086	A	19990430
DE 1999-19954228	A	19991104
DE 2000-10013363	A	20000309
DE 2000-10015836	A	20000327
JP 2000-615619	A3	20000501
WO 2000-IB657	W	20000501
IN 2001-MN1305	A3	20011019
US 2002-979939	A3	20020606

OTHER SOURCE(S): MARPAT 133:321769

AB The title compds. were prepared by various combinations of 3 fragments making up the mols. Thus, [4S,7R,8S,9S,13Z,16S(E)]-4,8-dihydroxy-16-[1-methyl-2-(2-pyridyl)ethenyl]-1-oxa-5,5,9,13-tetramethyl-7-(3-butynyl)-13-cyclohexadecene-2,6-dione was prepared in several steps starting from (4S)-4-(2-methyl-1-oxo-2-propyl)-2,2-dimethyl[1,3]dioxane and 5-(trimethylsilyl)-4-pentynylmagnesium bromide.

IT **303154-56-5P 303154-58-7P 303154-59-8P**
303154-60-1P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)


(6-alkenyl and 6-alkynyl derivs. of epothilone)

RN 303154-56-5 ZCAPLUS

CN 3-Undecanone, 2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-10-[[[(1,1-dimethylethyl)diphenylsilyl]oxy]-5-hydroxy-2,6-dimethyl-4-[4-(trimethylsilyl)-3-butynyl]-, (4R,5S,6S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.

Absolute stereochemistry.



IT 303154-57-6P


RL: SPN (Synthetic preparation); PREP (Preparation)

(preparation of 6-alkenyl-, 6-alkynyl- and 6-epoxyepothilone derivs. and their use in pharmaceutical preps.)

RN 303154-57-6 ZCAPLUS

CN 3-Undecanone, 2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-10-[[[(1,1-dimethylethyl)diphenylsilyl]oxy]-5-hydroxy-2,6-dimethyl-4-[4-(trimethylsilyl)-3-butyryl]-, (4S,5R,6S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT:

4

THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L25 ANSWER 11 OF 25 ZCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2000:772379 ZCAPLUS Full-text

DOCUMENT NUMBER: 133:321769

TITLE: 6-Alkenyl and 6-alkynyl derivatives of epothilone

INVENTOR(S): Klar, Ulrich; Schwede, Wolfgang; Skuballa, Werner; Buchmann, Bernd; Hoffmann, Jens; Lichtner, Rosemarie

PATENT ASSIGNEE(S): Schering A.-G., Germany

SOURCE: Ger. Offen., 18 pp.

CODEN: GWXXBX

DOCUMENT TYPE:

Patent


LANGUAGE:

German

FAMILY ACC. NUM. COUNT: 3

PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 19921086	A1	20001102	DE 1999-19921086	19990430
CA 2371226	A1	20001109	CA 2000-2371226	20000501
WO 2000066589	A1	20001109	WO 2000-IB657	20000501



RN 303154-58-7 ZCAPLUS

CN 3-Undecanone, 2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-10-[[[(1,1-dimethylethyl)diphenylsilyl]oxy]-2,6-dimethyl-5-[(tetrahydro-2H-pyran-2-yl)oxy]-4-[4-(trimethylsilyl)-3-butynyl]-, (4R,5S,6S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 303154-59-8 ZCAPLUS

CN 3-Undecanone, 4-(3-butynyl)-2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-10-hydroxy-2,6-dimethyl-5-[(tetrahydro-2H-pyran-2-yl)oxy]-, (4R,5S,6S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 303154-60-1 ZCAPLUS

CN 2,9-Undecanedione, 8-(3-butynyl)-10-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-6,10-dimethyl-7-[(tetrahydro-2H-pyran-2-yl)oxy]-, (6S,7S,8R)- (9CI) (CA INDEX NAME)

DE 1999-19954228	A1 19991104
DE 2000-10015836	A1 20000327
DE 2000-10013363	A 20000309
WO 2000-IB657	W 20000501
IN 2001-MN1305	A3 20011019
US 2002-979939	A3 20020606

OTHER SOURCE(S): MARPAT 133:362656

GI

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB The antitumor agents, 6-alkenyl-, 6-alkynyl- and 6-epoxyepothilones I (R1a, R1b are same or different = H, C1-C10 alkyl, C6-C12 aryl, C7-C20 aralkyl each optionally substituted; or together = (CH2)_m m = 1-5 or -CH2OCH2-; R2a(R2b replace a with b) = H, substituted alkyl, aryl, aralkyl, (CH2)_{ra}-C.tplbond.(or =)C-(CH2)_{pa}-R26a, Q, Q1 where n = 0-5; ra, rb = the same or different and = 0-4; pa, pb = the same or different and = 0-3; R3a = H, substituted alkyl, aryl or aralkyl; R3b = OH, OPG14; R14 = H, OR14a, halogen and R14a = H, SO2-alkyl, SO2-aryl or SO2-aralkyl; R4 = H, substituted alkyl, aryl or aralkyl; halogen, OR25, CN; R26a, R26b = same or different = H, substituted alkyl, aryl or aralkyl, C1-C10 acyl or if pa or pb > 0, addnl. a group OR27; R25 = R27 = R22 = H, PG; R5 = H, substituted alkyl, aryl or aralkyl, (CH2)_s s = 1-4, T = OR22 or halogen; R6, R7 = H or together = bond or O; G = X=CR8 or bi- or tricyclic aryl radical and R8 = H, halogen, CN, or substituted alkyl, aryl or aralkyl; X = O, two OR23 groups, C2-C10-alkylene- α,ω -dioxy straight chain or branched; H/OR9 or CR10R11 group and R23 = alkyl radical, R9 = H, PG, R10, R11 = same or different = H, substituted alkyl, aryl or aralkyl, or together with the methylene are a 5-7 carbocyclic ring; D-E = CH2CH2 or OCH2; A = OC(O), OCH2, CH2C(O), NR29C(O), NR29SO2 and R29 = H, alkyl; Z = O or H/OR12 and R12 = H, PG) were prepared. Thus II was prepared in a multistep synthesis starting from (4S)-4-(2-methyl-1-oxoprop-2-yl)-2,2-dimethyl[1,3]dioxane and 5-trimethylsilylpent-4-in-1-yl magnesium bromide. II had an IC50 value [nM] of 3.0 for the growth inhibition of human MCF-7 breast- and 75 for multidrug resistant NCI/ADR carcinoma cell lines with a selectivity of 2.5. The new epothilone derivs. interact with tubulin by stabilizing microtubuli that are formed. They are able to influence the cell-splitting in a phase-specific manner and are therefore useful in treating diseases or conditions associated with the need for cell growth, division and/or proliferation. Thus the epothilone derivs. are suitable for treating malignant tumors, e.g., ovarian, stomach, colon, adeno-, breast, lung, head and neck carcinomas, malignant melanoma, acute lymphocytic and myelocytic leukemia; and for anti-angiogenesis therapy as well as for treatment of chronic inflammatory diseases (such as psoriasis, arthritis).

IT 303154-56-5P 303154-58-7P 303154-59-8P
303154-60-1P


RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(preparation of 6-alkenyl-, 6-alkynyl- and 6-epoxyepothilone derivs. and their use in pharmaceutical preps.)

RN 303154-56-5 ZCAPLUS

CN 3-Undecanone, 2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-10-[[{(1,1-dimethylethyl)diphenylsilyl]oxy]-5-hydroxy-2,6-dimethyl-4-[4-(trimethylsilyl)-3-butyryl]-, (4R,5S,6S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L25 ANSWER 10 OF 25 ZCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2000:790507 ZCAPLUS Full-text

DOCUMENT NUMBER: 133:362656

TITLE: Preparation of 6-alkenyl-, 6-alkynyl- and 6-epoxyepothilone derivatives and their antitumor activity

INVENTOR(S): Klar, Ulrich; Schwede, Wolfgang; Skuballa, Werner; Buchmann, Bernd; Hoffmann, Jens; Lichtner, Rosemarie

PATENT ASSIGNEE(S): Schering Aktiengesellschaft, Germany

SOURCE: PCT Int. Appl., 298 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 3

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2000066589	A1	20001109	WO 2000-IB657	20000501
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW				
RW: GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
DE 19921086	A1	20001102	DE 1999-19921086	19990430
DE 19954228	A1	20010913	DE 1999-19954228	19991104
DE 10015836	A1	20011011	DE 2000-10015836	20000327
CA 2371226	A1	20001109	CA 2000-2371226	20000501
BR 2000010190	A	20020108	BR 2000-10190	20000501
EP 1173441	A1	20020123	EP 2000-922826	20000501
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, MC, PT, IE, SI, LT, LV, FI, RO				
JP 2002543203	T	20021217	JP 2000-615619	20000501
EE 200100568	A	20030217	EE 2001-568	20000501
NZ 514989	A	20040227	NZ 2000-514989	20000501
AU 772750	B2	20040506	AU 2000-43103	20000501
IN 2001MN01305	A	20070504	IN 2001-MN1305	20011019
BG 106053	A	20020531	BG 2001-106053	20011026
NO 2001005278	A	20011221	NO 2001-5278	20011029
MX 2001PA11039	A	20030630	MX 2001-PA11039	20011030
US 7125893	B1	20061024	US 2002-979939	20020606
IN 2005MN00837	A	20070608	IN 2005-MN837	20050802
US 2006046997	A1	20060302	US 2005-214988	20050831
PRIORITY APPLN. INFO.:			DE 1999-19921086	A1 19990430

PATENT ASSIGNEE(S): Bristol-Myers Squibb Company, USA
 SOURCE: PCT Int. Appl., 28 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001073103	A2	20011004	WO 2001-US9620	20010323
WO 2001073103	A3	20020523		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
US 2002042109	A1	20020411	US 2001-811808	20010319
US 6593115	B2	20030715		
US 2004023345	A1	20040205	US 2003-447082	20030528
PRIORITY APPLN. INFO.:			US 2000-191975P	P 20000324
			US 2001-811808	A3 20010319

OTHER SOURCE(S): CASREACT 135:287591; MARPAT 135:287591

AB The present invention relates to a process for the preparation of intermediates useful in the synthesis of epothilone analogs by initially enzymically degrading certain epothilone compds. to form ring-open structures containing a carboxyl group which is esterified, the hydroxyl groups on the moiety protected and the resulting compound oxidized by, e.g. ozone, to form a first intermediate. The first intermediate can be reacted with a triphenylphosphine adduct to yield a compound containing an ester group at position 1 which is subsequently hydrolyzed to form a second intermediate.


IT **364336-79-8P 364336-83-4P**

RL: PUR (Purification or recovery); RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 (preparation of epothilone intermediates)

RN 364336-79-8 ZCAPLUS

CN Dodecanoic acid, 3,7-dihydroxy-4,4,6,8-tetramethyl-5,12-dioxo-, methyl ester, (3S,6R,7S,8S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 364336-83-4 ZCAPLUS

CN Dodecanoic acid, 3,7-bis[[[1,1-dimethylethyl]dimethylsilyl]oxy]-4,4,6,8-tetramethyl-5,12-dioxo-, methyl ester, (3S,6R,7S,8S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 303154-59-8 ZCAPLUS

CN 3-Undecanone, 4-(3-butynyl)-2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-10-hydroxy-2,6-dimethyl-5-[(tetrahydro-2H-pyran-2-yl)oxy]-, (4R,5S,6S)-(9CI)
(CA INDEX NAME)


Absolute stereochemistry.



RN 303154-60-1 ZCAPLUS

CN 2,9-Undecanedione, 8-(3-butynyl)-10-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-6,10-dimethyl-7-[(tetrahydro-2H-pyran-2-yl)oxy]-, (6S,7S,8R)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.



L25 ANSWER 9 OF 25 ZCAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 2001:731069 ZCAPLUS Full-text
DOCUMENT NUMBER: 135:287591

TITLE: Preparation of epothilone intermediates
INVENTOR(S): Vite, Gregory D.; Kim, Soong-Hoon; Hoefle, Gerhard

303154-59-8P 303154-60-1P


RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(preparation of 12,13-cyclopropylepothilone derivs. and their use in pharmaceutical compns.)

RN 303154-56-5 ZCAPLUS

CN 3-Undecanone, 2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-10-[[[(1,1-dimethylethyl)diphenylsilyl]oxy]-5-hydroxy-2,6-dimethyl-4-[4-(trimethylsilyl)-3-butynyl]-, (4R,5S,6S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 303154-57-6 ZCAPLUS

CN 3-Undecanone, 2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-10-[[[(1,1-dimethylethyl)diphenylsilyl]oxy]-5-hydroxy-2,6-dimethyl-4-[4-(trimethylsilyl)-3-butynyl]-, (4S,5R,6S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 303154-58-7 ZCAPLUS

CN 3-Undecanone, 2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-10-[[[(1,1-dimethylethyl)diphenylsilyl]oxy]-2,6-dimethyl-5-[(tetrahydro-2H-pyran-2-yl)oxy]-4-[4-(trimethylsilyl)-3-butynyl]-, (4R,5S,6S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

FILE BEILSTEIN

FILE LAST UPDATED ON September 26, 2007

FILE COVERS 1771 TO 2007.

FILE CONTAINS 10.119,480 SUBSTANCES

>>>PLEASE NOTE: Reaction Data and substance data are stored in separate documents and can not be searched together in one query. Reaction data for BEILSTEIN compounds may be displayed immediately with the display codes PRE (preparations) and REA (reactions). A substance answer set retrieved after the search for a chemical name, a compounds with available reaction information by combining with PRE/FA, REA/FA or more generally with RX/FA. The BEILSTEIN Registry Number (BRN) is the link between a BEILSTEIN compound and belonging reactions. For more detailed reaction searches BRNs can be searched as reaction partner BRNs Reactant BRN (RX.RBRN) or Product BRN (RX.PBRN).<<<

>>> FOR SEARCHING PREPARATIONS SEE HELP PRE <<<

 * PLEASE NOTE THAT THERE ARE NO FORMATS FREE OF COST. *
 * SET NOTICE FEATURE: THE COST ESTIMATES CALCULATED FOR SET NOTICE *
 * ARE BASED ON THE HIGHEST PRICE CATEGORY. THEREFORE; THESE *
 * ESTIMATES MAY NOT REFLECT THE ACTUAL COSTS. *
 * FOR PRICE INFORMATION SEE HELP COST *

NEW

* **PATENT NUMBERS (PN) AND BABS ACCESSION NUMBERS (BABSAN) CAN NOW BE SEARCHED, SELECTED AND TRANSFERRED.**
 * **NEW DISPLAY FORMATS ALLREF, ALLP AND BABSAN SHOW ALL REFERENCES, ALL PATENT REFERENCES, OR ALL BABS ACCESSION NUMBERS FOR A COMPOUND AT A GLANCE.**


FILE BABS

FILE LAST UPDATED: 25 JUN 2007 <20070625/UP>



FILE COVERS 1980 TO DATE.

=>

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 10041470	A1	20020228	DE 2000-10041470	20000818
PRIORITY APPLN. INFO.:			DE 2000-10041470	20000818
OTHER SOURCE(S):	CASREACT 136:216592; MARPAT 136:216592			
GI				




I



AB The present invention describes new 6-alkenyl- and 6-alkynylepothilone derivs., e.g., I [R1a, R1b = H, C1-10-alkyl, aryl, C7-20-aralkyl; R1aR1b = (CH2)_r, CH2OCH2; r = 1 - 5; R2a = H, C1-10-alkyl, aryl, C7-20-aralkyl, (CH2)_m-C.tplbond.C-(CH2)_pR₂₆, (CH2)_m-C:C-(CH2)_pR₂₆, X1, X2; n = 0 - 5; p = 0 - 3; m = 0 - 4; R2b = (CH2)_m-C.tplbond.C-(CH2)_pR₂₆, (CH2)_m-C:C-(CH2)_pR₂₆, X1, X2; R3a = H, C1-10-alkyl, aryl, C7-20-aralkyl; R3b = O-protecting group; R4 = H, C1-10-alkyl, aryl, C7-20-aralkyl, halogen, OH, O-protecting group, CN; R5 = H, C1-10-alkyl, aryl, C7-20-aralkyl, (CH2)_s-T; S = 1 - 4; T = OH, O-protecting group, halogen; R6R7 = C(R33)₂, NR32 AY = OC(:O), OCH2, CH2C(:O), NR29C(:O), NR29SO2; DE = CH2CH2, CH2O, OCH2; G = X:CR8-, bicyclic or tricyclic aryl; X = O, (O-alkyl)₂, etc.; Z = H, H,OH, H,O-protective group; R8 = H, halogen, CN, C1-20-alkyl, aryl, C7-20-aralkyl; R14 = H, OH, halogen, O-SO2-alkyl, O-SO2-aryl, O-SO2-aralkyl; R26 = H, C1-10-alkyl, aryl, C7-20-aralkyl, C1-10-acyl, OH, O-protecting group; R29 = H, C1-20-alkyl; R32 = H, C1-4-alkyl, C1-4-acyl; R33 = H, halogen], which interact with tubulins by stabilizing the formed microtubulins (no data). I are able specifically to affect cell division and are suitable, for example for the treatment of malignant tumors ovarian -, stomach -, colon -, adeno -, chest -, lungs -, head and neck carcinoma, malignant melanoma, acute lymphocytic and myelocytic leukemia. In addition I are suitable for the anti-angiogenesis therapy as well as for the treatment of chronic ignitable illnesses (psoriasis, arthritis). For the avoidance of uncontrolled cell rampant growths on as well as the better compatibility of medical implants I can be up and/or brought into polymers materials. According to invention, I can be used alone or for the achievement of additive or synergistic effects in combination with further principles and substance classes applicable in the tumor therapy. Exptl. data from patents PCT/EP00/01333 and PCT/IB00/00657 are reproduced here.

IT 303154-56-5P 303154-57-6P 303154-58-7P



I

AB A facile and efficient route to epothilone analogs has been developed from the natural product epothilone D (I). Degradation of I via an oxidative cleavage sequence provides acid intermediate II rapidly in six steps. From II, a variety of epothilone analogs have been prepared utilizing ring-closing metathesis to reconstruct the trisubstituted-12,13-double bond. Using this approach, we report a number of epothilone analogs with varying C-15 aromatic side chains and C-14 allylic substitutions and their antitumor activities.

IT **681259-79-0P**


RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(preparation of epothilone D analogs via semisynthetic degradation and ring-closing metathesis and their antitumor activity)

RN 681259-79-0 ZCAPLUS

CN Tridecanoic acid, 3,7-bis[[1,1-dimethylethyl]dimethylsilyl]oxy]-4,4,6,8-tetramethyl-5,12-dioxo-, methyl ester, (3S,6R,7S,8S)- (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 14 THERE ARE 14 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L25 ANSWER 8 OF 25 ZCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2002:157050 ZCAPLUS Full-text

DOCUMENT NUMBER: 136:216592

TITLE: Procedures for the production of 12,13-cyclopropylepothilone derivatives, as well as for their use in pharmaceutical preparations

PATENT ASSIGNEE(S): Schering Ag, Germany

SOURCE: Ger. Offen., 64 pp.


CODEN: GWXXBX

DOCUMENT TYPE: Patent

LANGUAGE: German

FAMILY ACC. NUM. COUNT: 1


PATENT INFORMATION:



RN 823203-24-3 ZCAPLUS

CN 3-Undecanone, 2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-10-[[[(1,1-dimethylethyl)diphenylsilyl]oxy]-5-hydroxy-2,4,6-trimethyl-, (4S,5R,6S)- (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE REFORMAT

L25 ANSWER 7 OF 25 ZCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2004:106102 ZCAPLUS [Full-text](#)

DOCUMENT NUMBER: 140:357084

TITLE: Rapid access to epothilone analogs via semisynthetic degradation and reconstruction of epothilone D

AUTHOR(S): Dong, Steven D.; Sundermann, Kurt; Smith, Karen M. J.; Petryka, Joseph; Liu, Fenghua; Myles, David C.

CORPORATE SOURCE: Department of Chemistry, Kosan Biosciences, Hayward, CA, 94545, USA

SOURCE: Tetrahedron Letters (2004), 45(9), 1945-1947

CODEN: TELEAY; ISSN: 0040-4039


PUBLISHER: Elsevier Science B.V.

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 140:357084


GI



RN 823203-04-9 ZCAPLUS

CN 3-Undecanone, 2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-10-[[[(1,1-dimethylethyl)diphenylsilyl]oxy]-5-hydroxy-2,6-dimethyl-4-(2-propenyl)-, (4R,5S,6S,10R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 823203-05-0 ZCAPLUS

CN 3-Undecanone, 2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-10-[[[(1,1-dimethylethyl)diphenylsilyl]oxy]-5-hydroxy-2,6-dimethyl-4-(2-propenyl)-, (4R,5S,6S,10S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 823203-23-2 ZCAPLUS

CN 3-Undecanone, 2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-10-[[[(1,1-dimethylethyl)diphenylsilyl]oxy]-5-hydroxy-2,4,6-trimethyl-, (4R,5S,6S)- (CA INDEX NAME)


Absolute stereochemistry.



RN 220774-59-4 ZCAPLUS

CN 3-Undecanone, 2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-10-[[[(1,1-dimethylethyl)diphenylsilyl]oxy]-5-hydroxy-2,6-dimethyl-4-(phenylmethyl)-, (4S,5R,6S)]-(9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 303154-56-5 ZCAPLUS

CN 3-Undecanone, 2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-10-[[[(1,1-dimethylethyl)diphenylsilyl]oxy]-5-hydroxy-2,6-dimethyl-4-[4-(trimethylsilyl)-3-butynyl]-, (4R,5S,6S)]-(9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 303154-57-6 ZCAPLUS

CN 3-Undecanone, 2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-10-[[[(1,1-dimethylethyl)diphenylsilyl]oxy]-5-hydroxy-2,6-dimethyl-4-[4-(trimethylsilyl)-3-butynyl]-, (4S,5R,6S)]-(9CI) (CA INDEX NAME)

Absolute stereochemistry.

823203-04-9P 823203-05-0P 823203-23-2P

823203-24-3P


RL: IMF (Industrial manufacture); RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(preparation and tetrahydropyranylation of; method for producing C1-C15 fragments of epothilones and derivs. thereof)

RN 220774-19-6 ZCAPLUS

CN 3-Undecanone, 2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-10-[[[(1,1-dimethylethyl)diphenylsilyl]oxy]-4-ethyl-5-hydroxy-2,6-dimethyl-, (4R,5S,6S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 220774-20-9 ZCAPLUS

CN 3-Undecanone, 2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-10-[[[(1,1-dimethylethyl)diphenylsilyl]oxy]-4-ethyl-5-hydroxy-2,6-dimethyl-, (4S,5R,6S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 220774-58-3 ZCAPLUS

CN 3-Undecanone, 2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-10-[[[(1,1-dimethylethyl)diphenylsilyl]oxy]-5-hydroxy-2,6-dimethyl-4-(phenylmethyl)-, (4R,5S,6S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




IT 823203-07-2P

RL: IMF (Industrial manufacture); RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(preparation and perruthenate oxidation of; method for producing C1-C15 fragments of epothilones and derivs. thereof)

RN 823203-07-2 ZCAPLUS

CN 3-Undecanone, 2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-10-hydroxy-2,6-dimethyl-4-(2-propenyl)-5-[(tetrahydro-2H-pyran-2-yl)oxy]-, (4R,5S,6S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




IT 823203-19-6P

RL: IMF (Industrial manufacture); RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(preparation and silylation of; method for producing C1-C15 fragments of epothilones and derivs. thereof)

RN 823203-19-6 ZCAPLUS

CN 2,9-Tridecanedione, 7,11,13-trihydroxy-6,10,10-trimethyl-8-(2-propenyl)-, (6S,7S,8R,11S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



IT 220774-19-6P 220774-20-9P 220774-58-3P
220774-59-4P 303154-56-5P 303154-57-6P


preparation); PREP (Preparation); RACT (Reactant or reagent)

(preparation and detetrahydropyranylation of; method for producing C1-C15 fragments of epothilones and derivs. thereof)

RN 823203-17-4 ZCAPLUS

CN 3-Nonanone, 2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-5-hydroxy-2,6-dimethyl-9-(2-methyl-1,3-dioxolan-2-yl)-4-(2-propenyl)-, (4R,5S,6S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 823203-18-5 ZCAPLUS

CN 3-Nonanone, 2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-5-hydroxy-2,6-dimethyl-9-(2-methyl-1,3-dioxolan-2-yl)-4-(2-propenyl)-, (4S,5R,6S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



IT 220774-23-2P

RL: IMF (Industrial manufacture); RCT (Reactant); SPN (Synthetic


preparation); PREP (Preparation); RACT (Reactant or reagent)

(preparation and detetrahydropyranylation/deketalization of; method for producing C1-C15 fragments of epothilones and derivs. thereof)

RN 220774-23-2 ZCAPLUS

CN 2,9-Undecanedione, 10-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-8-ethyl-6,10-dimethyl-7-[(tetrahydro-2H-pyran-2-yl)oxy]-, (6S,7S,8R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 823203-06-1 ZCAPLUS

CN 3-Undecanone, 2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-10-[[[(1,1-dimethylethyl)diphenylsilyl]oxy]-2,6-dimethyl-4-(2-propenyl)-5-[(tetrahydro-2H-pyran-2-yl)oxy]-, (4R,5S,6S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry..



RN 823203-25-4 ZCAPLUS

CN 3-Undecanone, 2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-10-[[[(1,1-dimethylethyl)diphenylsilyl]oxy]-2,4,6-trimethyl-5-[(tetrahydro-2H-pyran-2-yl)oxy]-, (4R,5S,6S)- (CA INDEX NAME)

Absolute stereochemistry:



IT 823203-17-4P 823203-18-5P

RL: IMF (Industrial manufacture); RCT (Reactant); SPN (Synthetic)


IT 220774-21-0P 220774-60-7P 303154-58-7P
823203-06-1P 823203-25-4P

RL: IMF (Industrial manufacture); RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(preparation and desilylation of; method for producing C1-C15 fragments of epothilones and derivs. thereof)

RN 220774-21-0 ZCAPLUS

CN 3-Undecanone, 2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-10-[[[(1,1-dimethylethyl)diphenylsilyl]oxy]-4-ethyl-2,6-dimethyl-5-[(tetrahydro-2H-pyran-2-yl)oxy]-, (4R,5S,6S)- (9CI) (CA INDEX NAME)---


Absolute stereochemistry.



RN 220774-60-7 ZCAPLUS

CN 3-Undecanone, 2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-10-[[[(1,1-dimethylethyl)diphenylsilyl]oxy]-2,6-dimethyl-4-(phenylmethyl)-5-[(tetrahydro-2H-pyran-2-yl)oxy]-, (4R,5S,6S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 303154-58-7 ZCAPLUS

CN 3-Undecanone, 2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-10-[[[(1,1-dimethylethyl)diphenylsilyl]oxy]-2,6-dimethyl-5-[(tetrahydro-2H-pyran-2-yl)oxy]-4-[4-(trimethylsilyl)-3-butynyl]-, (4R,5S,6S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

trimethyl-5-[(tetrahydro-2H-pyran-2-yl)oxy]-, (4R,5S,6S)- (CA INDEX NAME)

Absolute stereochemistry.



IT 823203-08-3P 823203-20-9P


RL: IMF (Industrial manufacture); RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(preparation and Wittig reaction of, with (benzothiazolylpropyl)phosphonium iodide derivative; method for producing C1-C15 fragments of epothilones and derivs. thereof)

RN 823203-08-3 ZCAPLUS

CN 2,9-Undecanedione, 10-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-6,10-dimethyl-8-(2-propenyl)-7-[(tetrahydro-2H-pyran-2-yl)oxy]-, (6S,7S,8R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 823203-20-9 ZCAPLUS

CN 2,9-Tridecanedione, 7,11,13-tris[[(1,1-dimethylethyl)dimethylsilyl]oxy]-6,10,10-trimethyl-8-(2-propenyl)-, (6S,7S,8R,11S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 220774-22-1 ZCAPLUS

CN 3-Undecanone, 2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-4-ethyl-10-hydroxy-2,6-dimethyl-5-[(tetrahydro-2H-pyran-2-yl)oxy]-, (4R,5S,6S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 220774-61-8 ZCAPLUS

CN 3-Undecanone, 2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-10-hydroxy-2,6-dimethyl-4-(phenylmethyl)-5-[(tetrahydro-2H-pyran-2-yl)oxy]-, (4R,5S,6S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 303154-59-8 ZCAPLUS

CN 3-Undecanone, 4-(3-butynyl)-2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-10-hydroxy-2,6-dimethyl-5-[(tetrahydro-2H-pyran-2-yl)oxy]-, (4R,5S,6S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 823203-27-6 ZCAPLUS

CN 3-Undecanone, 2-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-10-hydroxy-2,4,6-

RN 220775-76-8 ZCAPLUS

CN 2,9-Undecanedione, 10-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-6,8,10-trimethyl-7-[(tetrahydro-2H-pyran-2-yl)oxy]-, (6S,7S,8R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 303154-60-1 ZCAPLUS

CN 2,9-Undecanedione, 8-(3-butynyl)-10-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-6,10-dimethyl-7-[(tetrahydro-2H-pyran-2-yl)oxy]-, (6S,7S,8R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 823203-02-7 ZCAPLUS

CN 2,9-Tridecanedione, 8-ethyl-7,11,13-trihydroxy-6,10,10-trimethyl-, (6S,7S,8R,11S)- (CA INDEX NAME)

Absolute stereochemistry.



IT 220774-22-1P 220774-61-8P 303154-59-8P
823203-27-6P

RL: IMF (Industrial manufacture); RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent).

(preparation and Swern oxidation of; method for producing C1-C15 fragments

of

epothilones and derivs. thereof)

alkyl, aryl, C7-20-aralkyl; R6, R7 = H; R6R7 = bond, O; G = X:CR8, bi- or tricyclic aryl; R8 = H, halogen, (un)substituted C1-20-alkyl, aryl, C7-20-aralkyl; X = O, (OR23)2, C2-10-alkylene- α,ω -dioxy, H(OR9), CR10R11; R23 = C1-20-alkyl; R9 = H, protecting group; R10, R11 = H, C1-10-alkyl, aryl, C7-20-aralkyl; CR10R11 = 5 - to 7-membered carbocycle; R13 = CH2OR13a, CH2-halo, CHO, CO2R13b, CO-halo; R13a, R14a = H, SO2alkyl, SO2-aryl, SO2-aralkyl; R13aR14a = (CH2)o, CR15aR15b; o = 2 - 4; R13b, R14b = H, C1-10-alkyl, aryl, C7-20-aralkyl; R15a, R15b = H, C1-10-alkyl, aryl, C7-20-aralkyl; R15aR15b = (CH2)q; q = 3 - 6; R20 = O-PG, NHR29, N3; Z = O, H(OR12); R12 = H, PG] of epothilones and derivs. The procedure comprises the bonding of a C1-C6 fragment, R13CH2CHR14CR1aR1bC(:O)CHR2aR2b, to a C7-C12 fragment, R5C(:V)(CH2)3CR4aR4bC(:W)R3a [V, W = O, (OR23)2, C2-10-alkylene- α,ω -dioxy, H(OR9)], to form a C1-C12 fragment, R5C(:V)(CH2)3CR4aR4bCR3a(O-PG14)CR2aR2bC(:Z)CR1aR1bCHR14CH2R13 [PG = H, protecting group], which is then treated with a C13-C15 fragment, G-CR20'CH2CHR7'R21 [R7' = H; R20' = halogen, N3, NHR29, OH, O-PG, NR29-PG, C1-20-(perfluoro)alkylsulfonyloxy, (C1-4-alkyl, NO2, Cl, Br-substituted) benzyloxy, NR29SO2Me, NR29C(:O)Me, CH2C(:O)Me; R21 = OH, halo, O-PG, P+Ph3Hal- (Hal = F, Cl, Br, I), P(O)(OQ)2 (Q = C1-10-alkyl, Ph), P(:O)Ph2; R29 = H, C1-6-alkyl], to form the C1-C15 epothilone intermediate product I. Thus, I [R1a = R1b = R5 = Me, R2a = CH2CH:CH2- β , R2b = R4b = H- α , R3 = H- β , R4a = Me- β , R6R7 = bond, R13 = CO2H, R14 = OSiMe2CMe3- β , R20 = OSiMe2CMe3- α , G = 2-methylbenzothiazol-5-yl, PG = SiMe2CMe3, Z = O] was prepared from (S)-4-(2-methyl-3-oxohept-6-en-2-yl)-2,2-dimethyl-1,3-dioxane via lithiation and reaction with (2S,6RS)-2-methyl-6-[(tert-butyl)dimethylsilyl]oxy]heptanal, tetrahydropyranlation, desilylation with Bu4NF in THF, oxidation in CH2Cl2 containing N-methylmorpholine N-oxide and catalytic tetrapropylammonium perruthenate, Wittig reaction with [(3S)-3-(2-methylbenzothiazol-5-yl)propyl]triphenylphosphonium iodide, deisopropylidenation/detetrahydropyranlation with catalytic 4-MeC6H4SO3H in EtOH, silylation with CF3SO2SiMe2CMe3, regioselective desilylation with (\pm)-camphor-10-sulfonic acid, Swern oxidation with DMSO/(COCl)2 in CH2Cl2 and carbonyl oxidation with NaOCl2 in aqueous THF/Me3COH. The produced C1-C15 epothilone intermediate products can be converted into the intrinsically active ingredients II [AK = OC(:O), OCH2, CH2C(:O), NR29C(:O), NR29SO2; R29 = H, C1-6-alkyl] according to known methods. The invention also relates to the corresponding C1-C12 fragments.


IT 220774-62-9P 220775-76-8P 303154-60-1P
823203-02-7P

RL: IMF (Industrial manufacture); SPN (Synthetic preparation); PREP (Preparation)
(method for producing C1-C15 fragments of epothilones and derivs. thereof)

RN 220774-62-9 ZCAPLUS


CN 2,9-Undecanedione, 10-[(4S)-2,2-dimethyl-1,3-dioxan-4-yl]-6,10-dimethyl-8-(phenylmethyl)-7-[(tetrahydro-2H-pyran-2-yl)oxy]-, (6S,7S,8R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




Skuballa, Werner
 PATENT ASSIGNEE(S): Schering Aktiengesellschaft, Germany
 SOURCE: PCT Int. Appl., 48 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: German
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2005003071	A1	20050113	WO 2004-EP6685	20040619
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
DE 10331004	A1	20050224	DE 2003-10331004	20030703
AU 2004254200	A1	20050113	AU 2004-254200	20040619
CA 2531078	A1	20050113	CA 2004-2531078	20040619
EP 1641734	A1	20060405	EP 2004-740122	20040619
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK, HR				
CN 1816514	A	20060809	CN 2004-80019005	20040619
BR 2004012179	A	20060822	BR 2004-12179	20040619
IN 2006DN00056	A	20070824	IN 2006-DN56	20060103
MX 2006PA00172	A	20060427	MX 2006-PA172	20060105
NO 2006000554	A	20060403	NO 2006-554	20060202
US 2007142675	A1	20070621	US 2006-563058	20060619
PRIORITY APPLN. INFO.:			DE 2003-10331004	A 20030703
			WO 2004-EP6685	W 20040619
OTHER SOURCE(S):		CASREACT 142:113814; MARPAT 142:113814		
GI				



I



II

AB The invention relates to a method for preparing C1-C15 fragments I [R1a, R1b = H, C1-10-alkyl, aryl, C7-20-aralkyl; R1aR1b = (CH2)m; m = 2 - 5; R2a, R2b = H, C1-10-alkyl, C2-10-alkenyl, C2-10-alkynyl, aryl, C7-20-aralkyl; R2aR2b = (CH2)n; n = 2 - 5; R3 = H, C1-10-alkyl, aryl, C7-20-aralkyl; R4a, R4b = H, C1-10-alkyl, aryl, C7-20-aralkyl; R4aR4b = (CH2)p; p = 2 - 5; R5 = H, C1-10-

AB Synthesis of C1-C12 segment of epothilones A and B was achieved via diastereo- and regioselective opening of a trisubstituted epoxy ketone at the more substituted carbon. Epoxide I (R = SiMe₂CMe₃, R₁ = benzyl) was cleaved selectively at the more substituted carbon using SmI₂ in MeOH/THF at -90° to form the silyl protected β-hydroxyketone II which contains the C5-C7 epothilone aldol moiety.

IT **201683-59-2P**


RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(synthesis of C1-C12 segment of epothilones A and B via radical induced opening of trisubstituted epoxides)

RN 201683-59-2 ZCAPLUS

CN Dodecanoic acid, 7-[[[(1,1-dimethylethyl)dimethylsilyl]oxy]-3-hydroxy-4,4,6,8-tetramethyl-5-oxo-12-(phenylmethoxy)-, 1,1-dimethylethyl ester, [3S-(3R*,6S*,7R*,8R*)]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



IT **201683-49-0P**


RL: SPN (Synthetic preparation); PREP (Preparation)

(synthesis of C1-C12 segment of epothilones A and B via radical induced opening of trisubstituted epoxides)

RN 201683-49-0 ZCAPLUS

CN Dodecanoic acid, 3,7-bis[[[(1,1-dimethylethyl)dimethylsilyl]oxy]-4,4,6,8-tetramethyl-5-oxo-12-(phenylmethoxy)-, 1,1-dimethylethyl ester, [3S-(3R*,6S*,7R*,8R*)]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT:

28 THERE ARE 28 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE REFORMAT


L25 ANSWER 6 OF 25 ZCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2005:29293 ZCAPLUS Full-text

DOCUMENT NUMBER: 142:113814

TITLE: Method for producing C1-C15 fragments of epothilones and derivatives thereof


INVENTOR(S): Klar, Ulrich; Buchmann, Bernd; Schwede, Wolfgang;



RN 346652-91-3 ZCAPLUS

CN Dodecanal, 6,10,12-tris[[(1,1-dimethylethyl)dimethylsilyl]oxy]-5,7,9,9-tetramethyl-8-oxo-, (5S,6S,7R,10S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 117 THERE ARE 117 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L25 ANSWER 5 OF 25 ZCAPLUS COPYRIGHT 2007 ACS on STN DUPLICATE 5

ACCESSION NUMBER: 1998:14400 ZCAPLUS Full-text

DOCUMENT NUMBER: 128:114806

TITLE: Radical-induced opening of trisubstituted epoxides: application in the synthesis of C1-C12 segment of epothilones

AUTHOR(S): Chakraborty, T. K.; Dutta, S.

CORPORATE SOURCE: Indian Inst. Chem. Technol., Hyderabad, 500 007, India

SOURCE: Tetrahedron Letters (1998), 39(1/2), 101-104

CODEN: TELEAY; ISSN: 0040-4039


PUBLISHER: Elsevier Science Ltd.



DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 128:114806

GI





AB Naturally occurring epothilones have been synthesized starting from enantiomerically pure aldol compds. I and II, which were obtained by antibody catalysis. Aldolase antibody 38C2 catalyzed the resolution of (\pm)-I by enantioselective retro-aldol reaction to afford I in 90% ee at 50% conversion. Compds. II (R = Me, CH₂OH) were obtained in more than 99% ee at 50% conversion by resolution of their racemic mixts. using newly developed aldolase antibodies 84G3, 85H6 or 93F3. Compds. I and II were resolved in multigram quantities and then converted to the epothilones by metathesis processes, which were catalyzed by Grubbs' catalysts.

IT **346651-96-5P**


RL: BPN (Biosynthetic preparation); RCT (Reactant); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); RACT (Reactant or reagent)

(total synthesis of epothilones A-F)

RN 346651-96-5 ZCAPLUS

CN 2,9-Tridecanedione, 7,11,13-tris[[1,1-dimethylethyl]dimethylsilyl]oxy]-6,8,10,10-tetramethyl-, (6S,7S,8R,11S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



IT **346652-88-8P 346652-91-3P**

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(total synthesis of epothilones A-F)

RN 346652-88-8 ZCAPLUS

CN 4,12-Dioxo-3,13-disilapentadecan-7-one, 9-[[1,1-dimethylethyl]dimethylsilyl]oxy]-5-[(1S)-5-hydroxy-1-methylpentyl]-2,2,3,3,6,8,8,13,13,14,14-undecamethyl-, (5S,6R,9S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

CORPORATE SOURCE: Abteilung Naturstoffchemie, Gesellschaft fuer
Biotechnologische Forschung mbH, Braunschweig,
D-38124, Germany

SOURCE: Journal of the Chemical Society, Perkin Transactions 1
(2002), (22), 2490-2503
CODEN: JCSPCE; ISSN: 1472-7781

PUBLISHER: Royal Society of Chemistry

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 138:187542

AB: Novel and unique chiral building blocks of high structural diversity were obtained by selective chemical fragmentation of natural products from myxobacteria. Subsequent modification reactions provided primary alc. and carboxylic acid derivs., which are suitable for the construction of combinatorial chemical libraries. The single SPOT synthesis of a hybrid structure on a polypropylene membrane was employed to demonstrate the chemical recombination of such rare building blocks on a micro-scale.


IT 498580-02-2P

RL: SPN (Synthetic preparation); PREP (Preparation)
(fragmentation of natural products from myxobacteria as building blocks for combinatorial synthesis)

RN 498580-02-2 ZCAPLUS

CN Dodecanoic acid, 3,7,12-trihydroxy-4,4,6,8-tetramethyl-5-oxo-, methyl ester, (3S,6R,7S,8S)- (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 50 THERE ARE 50 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L25 ANSWER 4 OF 25 ZCAPLUS COPYRIGHT 2007 ACS on STN DUPLICATE 4

ACCESSION NUMBER: 2001:316603 ZCAPLUS Full-text

DOCUMENT NUMBER: 135:76707

TITLE: Catalytic antibody route to the naturally occurring
epothilones: total synthesis of epothilones A - F

AUTHOR(S): Sinha, Subhash C.; Sun, Jian; Miller, Gregory P.;
Wartmann, Markus; Lerner, Richard A.

CORPORATE SOURCE: Department of Molecular Biology and the Skaggs
Institute for Chemical Biology, The Scripps Research
Institute, La Jolla, CA, 92037, USA

SOURCE: Chemistry--A European Journal (2001), 7(8), 1691-1702
CODEN: CEUJED; ISSN: 0947-6539

PUBLISHER: Wiley-VCH Verlag GmbH


DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 135:76707

GI

PUBLISHER: CODEN: ACIEF5; ISSN: 1433-7851
 DOCUMENT TYPE: Wiley-VCH Verlag GmbH & Co. KGaA
 LANGUAGE: Journal
 OTHER SOURCE(S): English
 CASREACT 139:197285
 GI



I

AB A total synthesis of epothilone C (I) with concomitant formal synthesis of epothilone A is described, using immobilized reagents and scavengers to effect multistep synthetic transformations and purifications.

IT **346652-91-3P**


RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(total synthesis of epothilones using solid-supported reagents and scavengers)

RN 346652-91-3 ZCAPLUS

CN Dodecanal, 6,10,12-tris[[(1,1-dimethylethyl)dimethylsilyl]oxy]-5,7,9,9-tetramethyl-8-oxo-, (5S,6S,7R,10S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 73 THERE ARE 73 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT



L25 ANSWER 3 OF 25 ZCAPLUS COPYRIGHT 2007 ACS on STN DUPLICATE 3

ACCESSION NUMBER: 2002:863402 ZCAPLUS Full-text

DOCUMENT NUMBER: 138:187542

TITLE: Natural product-derived building blocks for combinatorial synthesis. Part 1. Fragmentation of natural products from myxobacteria

AUTHOR(S): Niggemann, Jutta; Michaelis, Katrin; Frank, Ronald; Zander, Norbert; Hoefle, Gerhard



AB The total synthesis of the cytotoxic antitumor natural product epothilone C has provided a stage for the exploitation and further development of immobilized reagent methods. A stereoselective convergent synthetic strategy was applied, incorporating polymer-supported reagents, catalysts, scavengers and catch-and-release techniques to avoid frequent aqueous work-up and chromatog. purification. The enantioselective preparation of 3 key fragments heptanone I, (S)-2-methyl-6-heptenal, and thiazole II along with their elaboration via diastereoselective coupling into epothilone C is presented.

IT 346652-91-3P


RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(total synthesis of epothilone C via asym. synthesis and stereoselective coupling of heptanone, methylheptenal, and thiazole fragments using immobilized reagents and scavengers)

RN 346652-91-3 ZCAPLUS

CN Dodecanal, 6,10,12-tris[[(1,1-dimethylethyl)dimethylsilyl]oxy]-5,7,9,9-tetramethyl-8-oxo-, (5S,6S,7R,10S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 122 THERE ARE 122 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L25 ANSWER 2 OF 25 ZCAPLUS COPYRIGHT 2007 ACS on STN DUPLICATE 2

ACCESSION NUMBER: 2003:494861 ZCAPLUS Full-text

DOCUMENT NUMBER: 139:197285

TITLE: A total synthesis of epothilones using solid-supported reagents and scavengers

AUTHOR(S): Storer, R. Ian; Takemoto, Toshiyasu; Jackson, Philip S.; Ley, Steven V.

CORPORATE SOURCE: University Chemical Laboratories, University of Cambridge, Cambridge, CB2 1EW, UK

SOURCE: Angewandte Chemie, International Edition (2003), 42(22), 2521-2525

licensed to Beilstein GmbH and MDL Information Systems GmbH

FILE LAST UPDATED: 25 JUN 2007 <20070625/UP>

FILE COVERS 1980 TO DATE.

=> d stat que L14

L14 7 SEA FILE=BABS ABB=ON PLU=ON (6300090/BABSAN OR 6630563/BABSAN
OR 6085475/BABSAN OR 6376421/BABSAN OR 6410256/BABSAN OR
6473119/BABSAN OR 6597156/BABSAN)

=> dup rem L6 L19 L14

DUPLICATE IS NOT AVAILABLE IN 'BEILSTEIN'.

ANSWERS FROM THESE FILES WILL BE CONSIDERED UNIQUE

FILE 'ZCAPLUS' ENTERED AT 15:47:21 ON 11 OCT 2007

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2007 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'BEILSTEIN' ENTERED AT 15:47:21 ON 11 OCT 2007

COPYRIGHT (c) 2007 Beilstein-Institut zur Foerderung der Chemischen Wissenschaften
licensed to Beilstein GmbH and MDL Information Systems GmbH

FILE 'BABS' ENTERED AT 15:47:21 ON 11 OCT 2007

COPYRIGHT (c) 2007 Beilstein-Institut zur Foerderung der Chemischen Wissenschaften
licensed to Beilstein GmbH and MDL Information Systems GmbH

PROCESSING COMPLETED FOR L6

PROCESSING COMPLETED FOR L19

PROCESSING COMPLETED FOR L14

L25 25 DUP REM L6 L19 L14 (7 DUPLICATES REMOVED)

ANSWERS '1-18' FROM FILE ZCAPLUS

ANSWERS '19-25' FROM FILE BEILSTEIN

=> d ibib abs hitstr L25 1-18; d ide allref L25 19-25

L25 ANSWER 1 OF 25 ZCAPLUS COPYRIGHT 2007 ACS on STN DUPLICATE 1

ACCESSION NUMBER: 2004:454851 ZCAPLUS Full-text

DOCUMENT NUMBER: 141:140221

TITLE: Multi-step application of immobilized reagents and
scavengers: A total synthesis of epothilone C

AUTHOR(S): Storer, R. Ian; Takemoto, Toshiyasu; Jackson, Philip
S.; Brown, Dearg S.; Baxendale, Ian R.; Ley, Steven V.

CORPORATE SOURCE: Department of Chemistry, University of Cambridge,
Cambridge, CB2 1EW, UK

SOURCE: Chemistry--A European Journal (2004), 10(10),
2529-2547

CODEN: CEUJED; ISSN: 0947-6539


PUBLISHER: Wiley-VCH Verlag GmbH & Co. KGaA

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 141:140221

GI



chain nodes :

12 13 14 15 16 17 18 19 20 21 22

ring/chain nodes :

1 2 3 4 5 6 7 8 9 10 11

chain bonds :

2-12 2-13 3-14 3-15 4-16 4-17 6-18 8-19 10-20 11-21 11-22

ring/chain bonds :

1-2 2-3 3-4 4-5 5-6 6-7 7-8 8-9 9-10 10-11

exact/norm bonds :

1-2 2-3 3-4 4-5 5-6 6-7 6-18 7-8 8-9 8-19 9-10 10-11

exact bonds :

2-12 2-13 3-14 3-15 4-16 4-17 10-20 11-21 11-22

Match level :

1:CLASS 2:CLASS 3:CLASS 4:CLASS 5:CLASS 6:CLASS 7:CLASS 8:CLASS 9:CLASS

10:CLASS 11:CLASS 12:CLASS 13:CLASS 14:CLASS 15:CLASS 16:CLASS 17:CLASS

18:CLASS 19:CLASS

20:CLASS 21:CLASS 22:CLASS

L3 SCR 1008
 L11 38 SEA FILE=BEILSTEIN SSS FUL L1 AND L3
 L12 26 SEA FILE=BEILSTEIN ABB=ON PLU=ON L11/COM
 L13 5 SEA FILE=BEILSTEIN ABB=ON PLU=ON L12 AND BABSAN/FA
 L15 21 SEA FILE=BEILSTEIN ABB=ON PLU=ON L12 NOT L13
 L16 14 SEA FILE=BEILSTEIN ABB=ON PLU=ON L15 AND RN/FA
 L19 7 SEA FILE=BEILSTEIN ABB=ON PLU=ON L15 NOT L16

=> file babs

FILE 'BABS' ENTERED AT 15:47:02 ON 11 OCT 2007

COPYRIGHT (c) 2007 Beilstein-Institut zur Foerderung der Chemischen Wissenschaften

FILE COVERS 1771 TO 2007.

*** FILE CONTAINS 10.119,480 SUBSTANCES ***

>>>PLEASE NOTE: Reaction Data and substance data are stored in separate documents and can not be searched together in one query. Reaction data for BEILSTEIN compounds may be displayed immediately with the display codes PRE (preparations) and REA (reactions). A substance answer set retrieved after the search for a chemical name, a compounds with available reaction information by combining with PRE/FA, REA/FA or more generally with RX/FA. The BEILSTEIN Registry Number (BRN) is the link between a BEILSTEIN compound and belonging reactions. For more detailed reaction searches BRNs can be searched as reaction partner BRNs Reactant BRN (RX.RBRN) or Product BRN (RX.PBRN).<<<

>>> FOR SEARCHING PREPARATIONS SEE HELP PRE <<<

 * PLEASE NOTE THAT THERE ARE NO FORMATS FREE OF COST. *
 * SET NOTICE FEATURE: THE COST ESTIMATES CALCULATED FOR SET NOTICE *
 * ARE BASED ON THE HIGHEST PRICE CATEGORY. THEREFORE; THESE *
 * ESTIMATES MAY NOT REFLECT THE ACTUAL COSTS. *
 * FOR PRICE INFORMATION SEE HELP COST *

NEW


- * **PATENT NUMBERS (PN) AND BABS ACCESSION NUMBERS (BABSAN) CAN NOW BE SEARCHED, SELECTED AND TRANSFERRED.**
- * **NEW DISPLAY FORMATS ALLREF, ALLP AND BABSAN SHOW ALL REFERENCES, ALL PATENT REFERENCES, OR ALL BABS ACCESSION NUMBERS FOR A COMPOUND AT A GLANCE.**

=> d stat que L19

L1 STR

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

Structure attributes must be viewed using STN Express query preparation:
 Uploading L1.str



chain nodes :

12 13 14 15 16 17 18 19 20 21 22

ring/chain nodes :

1 2 3 4 5 6 7 8 9 10 11

chain bonds :

2-12 2-13 3-14 3-15 4-16 4-17 6-18 8-19 10-20 11-21 11-22

ring/chain bonds :

1-2 2-3 3-4 4-5 5-6 6-7 7-8 8-9 9-10 10-11

exact/norm bonds :

1-2 2-3 3-4 4-5 5-6 6-7 6-18 7-8 8-9 8-19 9-10 10-11

exact bonds :

2-12 2-13 3-14 3-15 4-16 4-17 10-20 11-21 11-22

Match level :

1:CLASS 2:CLASS 3:CLASS 4:CLASS 5:CLASS 6:CLASS 7:CLASS 8:CLASS 9:CLASS

10:CLASS 11:CLASS 12:CLASS 13:CLASS 14:CLASS 15:CLASS 16:CLASS 17:CLASS

18:CLASS 19:CLASS

20:CLASS 21:CLASS 22:CLASS

L3

SCR 1008

L5

68 SEA FILE=REGISTRY SSS FUL L1 AND L3

L6

18 SEA FILE=ZCAPLUS ABB=ON PLU=ON L5

=> file beilstein

FILE 'BEILSTEIN' ENTERED AT 15:46:51 ON 11 OCT 2007

COPYRIGHT (c) 2007 Beilstein-Institut zur Foerderung der Chemischen Wissenschaften
licensed to Beilstein GmbH and MDL Information Systems GmbH

FILE LAST UPDATED ON September 26, 2007

=> file registry

FILE 'REGISTRY' ENTERED AT 15:46:28 ON 11 OCT 2007
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2007 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file
provided by InfoChem.

STRUCTURE FILE UPDATES: 10 OCT 2007 HIGHEST RN 950149-06-1
DICTIONARY FILE UPDATES: 10 OCT 2007 HIGHEST RN 950149-06-1

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH June 29, 2007

Please note that search-term pricing does apply when
conducting SmartSELECT searches.

REGISTRY includes numerically searchable data for experimental and
predicted properties as well as tags indicating availability of
experimental property data in the original document. For information
on property searching in REGISTRY, refer to:

<http://www.cas.org/support/stngen/stndoc/properties.html>

=> file zcaplus

FILE 'ZCAPLUS' ENTERED AT 15:46:31 ON 11 OCT 2007
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2007 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is
held by the publishers listed in the PUBLISHER (PB) field (available
for records published or updated in Chemical Abstracts after December
26, 1996), unless otherwise indicated in the original publications.
The CA Lexicon is the copyrighted intellectual property of the
American Chemical Society and is provided to assist you in searching
databases on STN. Any dissemination, distribution, copying, or storing
of this information, without the prior written consent of CAS is
strictly prohibited.

FILE COVERS 1907 - 11 Oct 2007 VOL 147 ISS 16
FILE LAST UPDATED: 10 Oct 2007 (20071010/ED)

New CAS Information Use Policies, enter HELP USAGETERMS for details.

This file contains CAS Registry Numbers for easy and accurate
substance identification.


'OBI' IS DEFAULT SEARCH FIELD FOR 'ZCAPLUS' FILE

=> d stat que L6

L1 STR

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

Structure attributes must be viewed using STN Express query preparation:
Uploading L1.str



chain nodes :

12 13 14 15 16 17 18 19 20 21 22

ring/chain nodes :

1 2 3 4 5 6 7 8 9 10 11

chain bonds :

2-12 2-13 3-14 3-15 4-16 4-17 6-18 8-19 10-20 11-21 11-22

ring/chain bonds :

1-2 2-3 3-4 4-5 5-6 6-7 7-8 8-9 9-10 10-11

exact/norm bonds :

1-2 2-3 3-4 4-5 5-6 6-7 6-18 7-8 8-9 8-19 9-10 10-11

exact bonds :

2-12 2-13 3-14 3-15 4-16 4-17 10-20 11-21 11-22

Match level :

1:CLASS 2:CLASS 3:CLASS 4:CLASS 5:CLASS 6:CLASS 7:CLASS 8:CLASS 9:CLASS

10:CLASS 11:CLASS 12:CLASS 13:CLASS 14:CLASS 15:CLASS 16:CLASS 17:CLASS

18:CLASS 19:CLASS

20:CLASS 21:CLASS 22:CLASS

L3

SCR 1008

L11 38 SEA FILE=BEILSTEIN SSS FUL L1 AND L3

L12 26 SEA FILE=BEILSTEIN ABB=ON PLU=ON L11/COM

L13 5 SEA FILE=BEILSTEIN ABB=ON PLU=ON L12 AND BABSAN/FA

L15 21 SEA FILE=BEILSTEIN ABB=ON PLU=ON L12 NOT L13

L16 14 SEA FILE=BEILSTEIN ABB=ON PLU=ON L15 AND RN/FA

L19 7 SEA FILE=BEILSTEIN ABB=ON PLU=ON L15 NOT L16

=> file babs

FILE 'BABS' ENTERED AT 15:47:02 ON 11 OCT 2007

COPYRIGHT (c) 2007 Beilstein-Institut zur Foerderung der Chemischen Wissenschaften

FILE COVERS 1771 TO 2007.

*** FILE CONTAINS 10.119,480 SUBSTANCES ***

>>>PLEASE NOTE: Reaction Data and substance data are stored in separate documents and can not be searched together in one query. Reaction data for BEILSTEIN compounds may be displayed immediately with the display codes PRE (preparations) and REA (reactions). A substance answer set retrieved after the search for a chemical name, a compounds with available reaction information by combining with PRE/FA, REA/FA or more generally with RX/FA. The BEILSTEIN Registry Number (BRN) is the link between a BEILSTEIN compound and belonging reactions. For more detailed reaction searches BRNs can be searched as reaction partner BRNs Reactant BRN (RX.RBRN) or Product BRN (RX.PBRN).<<<

>>> FOR SEARCHING PREPARATIONS SEE HELP PRE <<<

 * PLEASE NOTE THAT THERE ARE NO FORMATS FREE OF COST. *
 * SET NOTICE FEATURE: THE COST ESTIMATES CALCULATED FOR SET NOTICE *
 * ARE BASED ON THE HIGHEST PRICE CATEGORY; THEREFORE; THESE *
 * ESTIMATES MAY NOT REFLECT THE ACTUAL COSTS. *
 * FOR PRICE INFORMATION SEE HELP COST *

NEW

* **PATENT NUMBERS (PN) AND BABS ACCESSION NUMBERS (BABSAN) CAN NOW BE SEARCHED, SELECTED AND TRANSFERRED.**
 * **NEW DISPLAY FORMATS ALLREF, ALLP AND BABSAN SHOW ALL REFERENCES, ALL PATENT REFERENCES, OR ALL BABS ACCESSION NUMBERS FOR A COMPOUND AT A GLANCE.**

=> d stat que L19

L1 STR

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

Structure attributes must be viewed using STN Express query preparation:
 Uploading L1.str

INTERNATIONAL SEARCH REPORT

Intern al Application No
PCT/EP 2004/006685A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C07C49/17 C07D417/06 C07D493/04

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 C07C C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EPO-Internal, PAJ, BEILSTEIN Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 99/07692 A (KLAR ULRICH ; SCHERING AG (DE); BUCHMANN BERND (DE); SKUBALLA WERNER ()) 18 February 1999 (1999-02-18) cited in the application page 49, line 1 - page 50, line 15; claim 9 US 2003/0144523	1-5

☐ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- *8* document member of the same patent family

Date of the actual completion of the international search

13 October 2004

Date of mailing of the international search report

20/10/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel (+31-70) 340-2040, Tx. 31 651 epo nl
Fax (+31-70) 340-3016

Authorized officer

Seelmann, I

Abstract

This invention describes a process for the production of C₁-C₁₅-fragments of epothilones and derivatives thereof, in which a C₁-C₆-fragment is linked with a C₇-C₁₂-fragment to a C₁-C₁₂-fragment, and the latter then is reacted with a C₁₃-C₁₅-fragment to form the C₁-C₁₅ initial epothilone product that is to be produced.

The thus obtained C₁-C₁₅ initial epothilone products can be reacted according to known methods to form the actual active ingredients.

In addition, the invention relates to the corresponding C₁-C₁₂-fragments.

Translation

PATENT COOPERATION TREATY

PCT

INTERNATIONAL PRELIMINARY REPORT ON PATENTABILITY
(Chapter II of the Patent Cooperation Treaty)

(PCT Article 36 and Rule 70)

10/563058
ISR

Applicant's or agent's file reference 53110AWO		FOR FURTHER ACTION		See Form PCT/IPEA/416
International application No. PCT/EP2004/006685		International filing date (day/month/year) 19.06.2004		Priority date (day/month/year) 03.07.2003
International Patent Classification (IPC) or national classification and IPC C07C49/17, C07D417/06, C07D493/04				
Applicant SCHERING AKTIENGESELLSCHAFT				

<p>1. This report is the international preliminary examination report, established by this International Preliminary Examining Authority under Article 35 and transmitted to the applicant according to Article 36.</p> <p>2. This REPORT consists of a total of <u>5</u> sheets, including this cover sheet.</p> <p>3. This report is also accompanied by ANNEXES, comprising:</p> <p>a. <input type="checkbox"/> (sent to the applicant and to the International Bureau) a total of _____ sheets, as follows:</p> <p><input type="checkbox"/> sheets of the description, claims and/or drawings which have been amended and are the basis for this report and/or sheets containing rectifications authorized by this Authority (see Rule 70.16 and Section 607 of the Administrative Instructions).</p> <p><input type="checkbox"/> sheets which supersede earlier sheets, but which this Authority considers contain an amendment that goes beyond the disclosure in the international application as filed, as indicated in item 4 of Box No. I and the Supplemental Box.</p> <p>b. <input type="checkbox"/> (sent to the International Bureau only) a total of (indicate type and number of electronic carrier(s)) _____, containing a sequence listing and/or tables related thereto, in computer readable form only, as indicated in the Supplemental Box Relating to Sequence Listing (see Section 802 of the Administrative Instructions).</p>				
<p>4. This report contains indications relating to the following items:</p> <p><input checked="" type="checkbox"/> Box No. I Basis of the report</p> <p><input type="checkbox"/> Box No. II Priority</p> <p><input type="checkbox"/> Box No. III Non-establishment of opinion with regard to novelty, inventive step and industrial applicability</p> <p><input type="checkbox"/> Box No. IV Lack of unity of invention</p> <p><input checked="" type="checkbox"/> Box No. V Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement</p> <p><input type="checkbox"/> Box No. VI Certain documents cited</p> <p><input type="checkbox"/> Box No. VII Certain defects in the international application</p> <p><input type="checkbox"/> Box No. VIII Certain observations on the international application</p>				

Date of submission of the demand		Date of completion of this report	
Name and mailing address of the IPEA/EP		Authorized officer	
Facsimile No.		Telephone No.	

INTERNATIONAL PRELIMINARY REPORT ON PATENTABILITY

International application No.

PCT/EP2004/006685

Box No. I Basis of the report

1. With regard to the language, this report is based on the international application in the language in which it was filed, unless otherwise indicated under this item.

☐ This report is based on translations from the original language into the following language _____ which is the language of a translation furnished for the purposes of:

- ☐ international search (Rule 12.3 and 23.1(b))
☐ publication of the international application (Rule 12.4)
☐ international preliminary examination (Rule 55.2 and/or 55.3)

2. With regard to the elements of the international application, this report is based on (replacement sheets which have been furnished to the receiving Office in response to an invitation under Article 14 are referred to in this report as "originally filed" and are not annexed to this report):

☐ the international application as originally filed/furnished

☒ the description:

pages 1-36 _____ as originally filed/furnished

pages* _____ received by this Authority on _____

pages* _____ received by this Authority on _____

☒ the claims:

nos. 1-5 _____ as originally filed/furnished

nos.* _____ as amended (together with any statement) under Article 19

nos.* _____ received by this Authority on _____

nos.* _____ received by this Authority on _____

☐ the drawings:

sheets _____ as originally filed/furnished

sheets* _____ received by this Authority on _____

sheets* _____ received by this Authority on _____

☐ a sequence listing and/or any related table(s) - see Supplemental Box Relating to Sequence Listing.

3. ☐ The amendments have resulted in the cancellation of:

☐ the description, pages _____

☐ the claims, nos. _____

☐ the drawings, sheets/figs _____

☐ the sequence listing (specify): _____

☐ any table(s) related to sequence listing (specify): _____

4. ☐ This report has been established as if (some of) the amendments annexed to this report and listed below had not been made, since they have been considered to go beyond the disclosure as filed, as indicated in the Supplemental Box (Rule 70.2(c)).

☐ the description, pages _____

☐ the claims, nos. _____

☐ the drawings, sheets/figs _____

☐ the sequence listing (specify): _____

☐ any table(s) related to sequence listing (specify): _____

* If item 4 applies, some or all of those sheets may be marked "superseded."

INTERNATIONAL PRELIMINARY REPORT ON PATENTABILITY

International application No.

PCT/EP2004/006685

Box No. V	Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement		
1. Statement			
Novelty (N)	Claims		YES
	Claims	1-5	NO
Inventive step (IS)	Claims		YES
	Claims	1-5	NO
Industrial applicability (IA)	Claims	1-5	YES
	Claims		NO
2. Citations and explanations (Rule 70.7)			
<p>The present application appears not to satisfy the requirements of PCT Article 33(2) because the subject matter of the claims is not novel. Claim 9 and pages 49-50 of the description of document D1 (WO 99/07692 A) concern, <i>inter alia</i>, a method for producing epothilone derivatives from the fragments $A+B = A-B$ and $A-B + C = A-B-C$, wherein all three fragments structurally overlap the A, B and C claimed in the present application. The formula AB in claim 9 of document D1 therefore appears to be prejudicial to the novelty of the present claim 5. The method of claim 9 of document D1 likewise appears to be prejudicial to the novelty of present claims 1-4. In particular, in the C fragment $U=C-R$ appears to overlap with G in document D1, and in fragment AB CH-CH versus D-E in AB of document D1 does not result in a new selection, since D-E form a unit, that is to say, they cannot be selected independently of each other. Consequently, this is considered no more than a selection from a list.</p> <p>Document D1 is the closest prior art. It discloses the production of epothilone derivatives from the fragments $A+B = A-B$ and $A-B + C = A-B-C$. The problem to be solved</p>			

INTERNATIONAL PRELIMINARY REPORT ON PATENTABILITY

International application No.

PCT/EP2004/006685

Box No. V

Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement

by the present invention is understood to be that of providing an alternative method for producing epothilones. In the light of the experimental part it can be assumed that this problem is solved by the application. However, insofar as the subject matter of the present application can be considered novel, the fragments A, B and C are similar to those of document D1 to such an extent that the solution is obvious to a person skilled in the art. The problem to be solved by the present application must therefore be considered that of an alternative method having unexpected or surprising properties with respect to the closest prior art (D1). Without comparative test results or other arguments demonstrating the patentability of the invention it is not possible to assess whether the invention satisfies the requirements of PCT Article 33(3). The present application does not appear to meet the requirements of PCT Article 33(2) because the subject matter of the claims is not novel. Claim 9 and pages 49-50 of the description of document D1 (WO 99/07692 A) concern, *inter alia*, a method for producing epothilone derivatives from the fragments $A+B = A-B$ and $A-B + C = A-B-C$, wherein all three fragments structurally overlap with the fragments A, B and C claimed in the present application. The formula AB in claim 9 of document D1 therefore appears to be prejudicial to the novelty of the present claim 5. The method of claim 9 of document D1 likewise appears to be prejudicial to the novelty of the present claims 1-4. In particular, in the C fragment, U=C-R appears to overlap with G in document D1, and in fragment AB, CH-CH as opposed to D-E in AB of document D1 does not lead to a novel selection, since D-E form a unit, that is to say,

INTERNATIONAL PRELIMINARY REPORT ON PATENTABILITY

International application No.

PCT/EP2004/006685

Box No. V

Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability;
citations and explanations supporting such statement

they cannot be selected independently of each other.
Consequently, this is considered no more than a selection from a list.

Document D1 is the closest prior art. It discloses the production of epothilone derivatives from the fragments $A+B = A-B$ and $A-B + C = A-B-C$. The problem to be solved by the present invention is understood to be that of providing an alternative method for the production of epothilones. In the light of the experimental part, it can be assumed that this problem is solved in the application. However, insofar as the subject matter of the present application can be considered novel, the fragments A, B and C are similar to those of document D1 to such an extent that the solution is obvious to a person skilled in the art. The problem to be solved by the present application must therefore be considered that of making available an alternative method having unexpected or surprising properties with respect to the closest prior art document (D1). Without comparative test results or other arguments demonstrating the patentability of the invention it is not possible to assess whether the invention satisfies the requirements of PCT Article 33(3).

Lao, MariaLouisa

From: DiNatale, John
Sent: Thursday, October 11, 2007 4:01 PM
To: Lao, MariaLouisa
Subject: 10/563058

Examiner Lao,

Your search results for serial number 10/563058 are complete and have been submitted to SCORE for posting. Routinely these results will be posted as early as tomorrow. Please see the instructions at the bottom of this email for retrieving search results from eDan 2.2.1.

****Search-specific notes:**

The search results are located in 2 RTF files.

The organization of the search results within the RTF file called 20071011-10563058-str1.rtf is sequential, divided by page breaks:

- 1) author search**
- 2) Claim 1 reaction search,**
- 3) Claim 4 reaction search,**
- and**
- 4) search history.**

The organization of the search results within the RTF file called 20071011-10563058-str2.rtf is sequential, divided by page breaks:

- 1) Claim 5 structure search**
- and**
- 2) search history.**

It may be helpful to save this memo as an index to the search results.

Please contact me if you have any questions.

Thank you,
John DiNatale
X2-2557

To access your search results via eDAN:

- 1) Enter Application number**
- 2) Click on Supplemental Content Tab**
- 3) STN results (structure and text searches) are under **Other Content** (click on version listed). ABSS Sequence results are under the **Search Results** (click on version listed).**

```

-3/BI OR 220774-19-6/BI OR 220774-20-9/BI OR 220774-21-0/BI OR
220774-22-1/BI OR 220774-23-2/BI OR 220774-57-2/BI OR 220774-58
-3/BI OR 220774-59-4/BI OR 220774-60-7/BI OR 220774-61-8/BI OR
220774-62-9/BI OR 220775-18-8/BI OR 220775-76-8/BI OR 289477-70
-9/BI OR 289477-71-0/BI OR 289477-72-1/BI OR 289477-73-2/BI OR
289477-74-3/BI OR 303154-55-4/BI OR 303154-56-5/BI OR 303154-57
-6/BI OR 303154-58-7/BI OR 303154-59-8/BI OR 303154-60-1/BI OR
3035640-13-5/BI OR 823203-01-6/BI OR 823203-02-7/BI OR 823203-03
-8/BI OR 823203-04-9/BI OR 823203-05-0/BI OR 823203-06-1/BI OR
823203-07-2/BI OR 823203-08-3/BI OR 823203-09-4/BI OR 823203-10
-7/BI OR 823203-11-8/BI OR 823203-12-9/BI OR 823203-13-0/BI OR
823203-14-1/BI OR 823203-15-2/BI OR 823203-16-3/BI OR 823203-17
-4/BI OR 823203-18-5/BI OR 823203-19-6/BI OR 823203-20-9/BI OR
823203-23-2/BI OR 823203-24-3/BI OR 823203-25-4/BI OR 823203-27
-6/BI)
2 SEA ABB=ON PUJ=ON L66 AND L39
D SCA
L67

FILE 'CAPIUS' ENTERED AT 11:50:53 ON 11 OCT 2007
1 SEA ABB=ON PUJ=ON L67 AND L65
D SCA
L68

FILE 'REGISTRY' ENTERED AT 11:54:40 ON 11 OCT 2007
E EPOTHILONE C/GN
1 SEA ABB=ON PUJ=ON EPOTHILONE C/GN
D SCA
1 SEA ABB=ON PUJ=ON EPOTHILONE D/GN
D SCA
D SEA L67 1-2
L69

FILE 'CASREACT' ENTERED AT 11:56:56 ON 11 OCT 2007
21 SEA ABB=ON PUJ=ON 152044-54-7/PRO
14 SEA ABB=ON PUJ=ON 152044-53-6/PRO
7 SEA ABB=ON PUJ=ON L42 (L) L71
7 SEA ABB=ON PUJ=ON L42 (L) L72
14 SEA ABB=ON PUJ=ON (L73 OR L74)
13 SEA ABB=ON PUJ=ON L75 NOT L60
L70

FILE 'CAPIUS' ENTERED AT 11:58:45 ON 11 OCT 2007
11 SEA ABB=ON PUJ=ON L45 AND PY<2000
SEL AN
L71
L72
L73
L74
L75
L76
L77

```

```

FILE 'CASREACT' ENTERED AT 12:00:01 ON 11 OCT 2007
11 SEA ABB-ON PUJ-ON ('126:251010"/AN OR "127:108793"/AN OR
"127:293040"/AN OR "128:101936"/AN OR "129:189151"/AN OR
"131:195935"/AN OR "131:286299"/AN OR "131:318191"/AN OR
"131:318291"/AN OR "131:351125"/AN OR "132:49832"/AN OR
"1997:206419"/AN OR "1997:430309"/AN OR "1997:665094"/AN OR
"1997:787450"/AN OR "1998:378435"/AN OR "1999:1766999"/AN OR
"1999:372044"/AN OR "1999:383492"/AN OR "1999:444724"/AN OR
"1999:606636"/AN OR "1999:819379"/AN)
11 SEA ABB-ON PUJ-ON L78 AND L43
16 SEA ABB-ON PUJ-ON L79 OR L52
1 SEA ABB-ON PUJ-ON L80 AND L73
1 SEA ABB-ON PUJ-ON L80 AND L74
27 SEA ABB-ON PUJ-ON L52 OR L79 OR (L81 OR L82) OR L76
15 SEA ABB-ON PUJ-ON L61 NOT L63
D FHIT 7
59 SEA ABB-ON PUJ-ON L43 (L) 1/NS
45 SEA ABB-ON PUJ-ON L43 (L) 2/NS

```

169

37 SEA ABB=ON PIU=ON L43 (L) 3/NS
 31 SEA ABB=ON PIU=ON L43 (L) 4/NS
 21 SEA ABB=ON PIU=ON L43 (L) 5/NS
 28 SEA ABB=ON PIU=ON L85 NOT L88

 FILE 'CAPLUS' ENTERED AT 12:09:08 ON 11 OCT 2007
 80 SEA ABB=ON PIU=ON KLAR U7/4U
 116 SEA ABB=ON PIU=ON BUCHMANN B7/4U
 60 SEA ABB=ON PIU=ON SCHWEDE W7/4U
 186 SEA ABB=ON PIU=ON SKUBALLA W7/4U
 32 SEA ABB=ON PIU=ON L91 AND (L92 OR L93 OR L94)
 68 SEA ABB=ON PIU=ON L91 AND (L92 OR L93 OR L94)
 24 SEA ABB=ON PIU=ON L93 AND L94
 25 SEA ABB=ON PIU=ON L95 AND (L96 OR L97)
 24 SEA ABB=ON PIU=ON L96 AND L97
 24 SEA ABB=ON PIU=ON L98 AND L99
 1 SEA ABB=ON PIU=ON L100 AND L43

 FILE 'REGISTRY' ENTERED AT 12:11:27 ON 11 OCT 2007

 FILE 'CAPLUS' ENTERED AT 12:11:29 ON 11 OCT 2007
 D STAT QUE L100
 D IBIB ABS L100 1-24
 D IBIB ABS L100 8-24
 24 SEA ABB=ON PIU=ON L91 AND L92 AND L93 AND L94
 D COST FULL
 D IBIB ABS L102 TOT
 D IBIB L102 10
 D IBIB L102 9
 D ABS L102 8
 D ABS L102 8
 D IBIB ABS L102 9-24

 FILE 'REGISTRY' ENTERED AT 12:16:30 ON 11 OCT 2007

 FILE 'CASREACT' ENTERED AT 12:16:34 ON 11 OCT 2007
 D STAT QUE L33
 D IBIB ABS FHIT L33 1-23

 FILE 'CASREACT' ENTERED AT 12:18:14 ON 11 OCT 2007
 D STAT QUE L90
 D IBIB ABS FHIT L90 1-28
 3 SEA ABB=ON PIU=ON L77 NOT L90
 3 SEA ABB=ON PIU=ON L78 NOT L90

 FILE 'CAPLUS' ENTERED AT 12:40:57 ON 11 OCT 2007
 21 SEA ABB=ON PIU=ON L45 AND PY<2001
 SEL AN

 FILE 'CASREACT' ENTERED AT 12:41:26 ON 11 OCT 2007
 21 SEA ABB=ON PIU=ON ("126:251010"/AN OR "127:108793"/AN OR
 "127:293040"/AN OR "128:101936"/AN OR "129:189151"/AN OR
 "131:199535"/AN OR "131:286299"/AN OR "131:31819"/AN OR
 "131:31829"/AN OR "131:351125"/AN OR "132:251011"/AN OR
 "132:49832"/AN OR "133:266631"/AN OR "133:266634"/AN OR
 "133:321737"/AN OR "133:362657"/AN OR "134:178371"/AN OR
 "134:29228"/AN OR "134:47959"/AN OR "134:56502"/AN OR "135:37156/
 6"/AN OR "1997:206419"/AN OR "1997:430309"/AN OR "1997:665094"/
 AN OR "1997:787450"/AN OR "1998:378435"/AN OR "1999:176999"/AN
 OR "1999:372044"/AN OR "1999:383492"/AN OR "1999:444724"/AN OR

170.

"1999:606636"/AN OR "1999:819379"/AN OR "2000:514132"/AN OR
 "2000:523875"/AN OR "2000:597944"/AN OR "2000:624043"/AN OR
 "2000:701228"/AN OR "2000:719579"/AN OR "2000:733774"/AN OR
 "2000:842116"/AN OR "2000:853645"/AN OR "2001:843887"/AN
 7 SEA ABB=ON PIL=ON L106 NOT L90
 21 SEA ABB=ON PIL=ON L106 AND L43
 7 SEA ABB=ON PIL=ON L108 NOT L90
 D STAT QUE L109
 D IBIB ABS PHIT L109 1-7

FILE HOME

FILE REGISTRY

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 10 OCT 2007 HIGHEST RN 950149-06-1
 DICTIONARY FILE UPDATES: 10 OCT 2007 HIGHEST RN 950149-06-1

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH June 29, 2007

Please note that search-term pricing does apply when conducting SmartSELECT searches.

REGISTRY includes numerically searchable data for experimental and predicted properties as well as tags indicating availability of experimental property data in the original document. For information on property searching in REGISTRY, refer to:

<http://www.cas.org/support/stngen/stdoc/properties.html>

FILE ZCAPLUS

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS is strictly prohibited.

FILE COVERS 1907 - 11 Oct 2007 VOL 147 ISS 16
 FILE LAST UPDATED: 10 Oct 2007 (20071010/ED)

New CAS Information Use Policies, enter HELP USAGETERMS for details.

This file contains CAS Registry Numbers for easy and accurate substance identification.

FILE CASREACT

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications.

FILE CONTENT:1840 - 6 Oct 2007 VOL 147 ISS 16

New CAS Information Use Policies, enter HELP USAGETERMS for details.

 * CASREACT now has more than 13.8 million reactions *
 * *****

Some CASREACT records are derived from the ZIC/VINITI database (1974-1999) provided by InfoChem, INPI data prior to 1986, and Biotransformations database compiled under the direction of Professor Dr. Klaus Kieselich.

This file contains CAS Registry Numbers for easy and accurate substance identification.

FILE CAPLUS

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 11 Oct 2007 VOL 147 ISS 16
 FILE LAST UPDATED: 10 Oct 2007 (20071010/ED)

Effective October 17, 2005, revised CAS Information Use Policies apply. They are available for your review at:

<http://www.cas.org/infopolicy.html>

=>